



ECOLOGY
AUSTRALIA

Edithvale-Seafood Wetlands Ramsar Site Management Plan



Prepared for: Melbourne Water

©2016 Ecology Australia Pty Ltd and Melbourne Water Corporation

Copyright: Information and data recorded in this document is the property of Melbourne Water Corporation and Ecology Australia. This document and the information are solely for the use of the authorised recipient and this document may not be used, copied or reproduced in whole or part for any purpose other than that for which it was supplied by Ecology Australia or Melbourne Water Corporation. Melbourne Water Corporation and Ecology Australia make no representation, undertake no duty and accept no responsibility to any third party who may use or rely upon this document or the information.

Document information

This is a controlled document. Details of the document ownership, distribution, status and revision history are listed below.

All comments or requests for changes to content should be addressed to the document owner.

Owner	Ecology Australia Pty Ltd		
Project	15-088		
Author	Darren G. Quin, Fiona M. Sutton, Jennifer Hale and Andrew R.G. McMahon		
File	Edithvale Seaford Management Plan Final 26 Sep 2016a.docx		
Bioregion	Gippsland Plain		
Distribution	Paul Rees	Melbourne Water	

Document History			
Status	Changes	By	Date
Draft 1	First Draft	D Quin, F Sutton, J Hale, A McMahon	14/06/2016
Draft 2	Comments received from Melbourne Water	D Quin and F Sutton	16/09/2016
Final	Final Melbourne Water comments incorporated	D Quin and F Sutton	26/09/2016

Cover photo: Constructed wetlands in the north of Edithvale Wetlands, 6 June 2016.



88B Station Street, Fairfield 3078 VIC
T: (03) 9489 4191
E: admin@ecologyaustralia.com.au
W: ecologyaustralia.com.au

Contents

Acknowledgments	10
Summary Management Plan	11
1 Introduction	19
1.1 Background	19
1.2 Objectives of the Management Plan	19
1.3 Ramsar Management Policies and Guidelines	20
1.4 Consultative Framework	20
2 Ramsar Site Description	22
2.1 Location	22
2.2 Wetland Description	22
2.3 Wetland Types	24
2.4 Wetland Significance and Listings	24
2.4.1 Ramsar Criteria	24
2.4.2 Other listings/classifications	26
2.5 Land Tenure and Management	29
2.6 Landscape Setting	32
2.7 Catchment Setting	33
2.8 Local Government	33
3 Legislation and Policy Framework	34
3.1 International Legislation and Policy	34
3.1.1 The Ramsar Convention	34
3.1.2 Migratory Bird Bilateral Agreements and Conventions	35
3.1.3 East Asian-Australasian Flyway Partnership	36
3.2 National Legislation and Policy	38
3.2.1 <i>Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act)</i>	38
3.2.2 National Strategy for the Management of Coastal Acid Sulfate Soils 2000	38
3.3 Victorian Legislation and Policy	38
3.3.1 <i>Environment Protection Act 1970</i>	38
3.3.2 <i>Environment Effects Act 1978</i>	39
3.3.3 <i>Flora and Fauna Guarantee Act 1988</i>	39
3.3.4 <i>Water Act 1989</i>	39
3.3.5 <i>Planning and Environment Act 1986</i>	40
3.3.6 <i>Catchment and Land Protection Act 1994</i>	41
3.3.7 <i>Crown Land (Reserves) Act 1978</i>	41
3.3.8 <i>Coastal Management Act 1995</i>	42
3.3.9 <i>Wildlife Act 1975 and Wildlife Regulations 2002</i>	42
3.3.10 <i>Country Fire Authority Act 1958</i>	42
3.3.11 Victorian Waterway Management Strategy	43

3.3.12	Victorian Coastal Strategy 2014	43
3.3.13	Victorian Coastal Acid Sulfate Soils Strategy	43
3.3.14	<i>Public Health and Wellbeing Act 2008</i>	43
3.3.15	Framework for Mosquito Management in Victoria (DSE 2004)	44
3.4	Regional and Local Plans and Policy	45
3.4.1	Port Phillip and Western Port Regional Catchment Strategy	45
3.4.2	Better Bays and Waterways	45
3.4.3	Healthy Waterways Strategy	45
4	Ecological Character and Values - Critical Components, Processes and Services	47
4.1	Waterbird Diversity and Abundance	47
4.2	Waterbird Breeding	49
4.3	Physical Habitat for Waterbirds	49
4.4	Threatened Wetland Species	49
4.5	Flora	50
4.5.1	Vegetation	50
4.5.2	Significant Flora Species	54
4.6	Other Fauna	57
4.6.1	Threatened Bird Species	57
4.6.2	Mammals	60
4.6.3	Frogs	62
4.6.4	Reptiles	63
4.6.5	Fish	63
4.6.6	Invertebrates	64
4.7	Function	65
4.8	Cultural Heritage	65
4.8.1	Edithvale Wetlands	65
4.8.2	Seaford Wetlands	66
4.9	Social and Amenity Values	67
4.10	Scientific Research and Long-term Monitoring	67
4.11	Buffer Opportunities	68
4.12	'Downs Estate'	68
5	Status of Ecological Character and Limits of Acceptable Change (LAC)	70
6	Site Condition	72
6.1	Vegetation	72
6.2	Water Quality	72
6.2.1	Water Quality Sampling 2009/2010	73
6.2.2	Water Quality Monitoring 2009 Onwards	73
6.3	Hydrology	77
6.3.1	Edithvale South Wetlands	79
6.3.2	Edithvale North Wetlands	79
6.3.3	Seaford Wetlands	80

7	Risk Assessment	85
8	Threatening Processes	86
8.1	Water Management: Urban Stormwater – Altered Water Regimes	86
8.2	Water Management: Urban Stormwater – Increased Salinity	86
8.3	Pollution: Urban Stormwater – Increased Nutrients, Sediments and Toxicants	87
8.4	Urban Development and Recreation – Litter	87
8.5	Disturbance of Potential Acid Sulfate Soils – Acidity and Liberation of Heavy Metals	87
8.6	Invasive Species – Exotic Plants	88
8.7	Invasive Species – Exotic Grazing Animals (European Rabbit and Hares)	88
8.8	Invasive Species – Exotic Terrestrial Predators (foxes, cats and rats)	88
8.9	Invasive Species – Exotic Predatory Fish	89
8.10	Out of Balance/Invasive Native Species	89
8.10.1	Common Reed and Cumbungi	89
8.10.2	Coast Tea-tree and Coast Wattle	89
8.10.3	Eastern Grey Kangaroo	90
8.11	Recreational Activities – Human Disturbance to Native Wildlife	90
8.12	Climate Change	92
8.13	Fire and Fuel Management	92
9	Management	94
9.1	Management Categories	95
9.2	Management Zones and Objectives	95
9.3	Management Priorities	100
9.4	Priority 1 Management Actions (Legislative Obligations)	107
9.4.1	Grooming of Common Reed (Program Maintenance)	107
9.4.2	Monitoring of Common Reed Management Program Performance (Monitoring and Assessment)	108
9.4.3	Maintain Current Hydrology Regime (Program Maintenance)	109
9.4.4	Cost-benefit Analysis of Management Actions Associated with Climate Change (Investigation)	110
9.4.5	Monitoring of Abiotic and Biotic Changes with Climate Change (Monitoring and Assessment)	110
9.4.6	Manage CaLP Listed ‘Regionally Controlled’ Weeds (Program Maintenance)	111
9.4.7	Control CaLP Listed Pest Animals (Program Maintenance)	115
9.4.8	Mosquito Monitoring and Control (Program Maintenance)	115
9.4.9	Monitoring of Bird Populations (Biodiversity Monitoring and Assessment)	116
9.4.10	Potential Acid Sulfate Soil Management	117
9.4.11	Fire Management (Program Maintenance)	117
9.5	Priority 2 Management Actions (Best Practice Management)	118
9.5.1	Incorporate ‘Downs Estate’ into Ramsar Site (Capital/Grants Program)	118
9.5.2	Manage High Threat Weeds (Program Maintenance)	119
9.5.3	Maintain Dog Restrictions and Increase Signage (capital works)	119

9.5.4	Control of Other Pest Animals (Program Maintenance)	120
9.5.5	Undertake Nest Box Monitoring (Program Maintenance)	120
9.5.6	Revegetation after Disturbance (Program Maintenance)	120
9.5.7	Consolidate Melbourne Water and Frankston City Council Management Boundaries at Seaford Wetland (Capital/Grants Program)	121
9.5.8	Investigate Path Extension at Seaford Wetland (Investigation)	121
9.5.9	Fencing (Program Maintenance)	122
9.5.10	Monitoring (Program Maintenance)	122
9.5.11	Monitoring (Biodiversity Monitoring and Assessment)	123
9.6	Priority 3 Management Actions (Other Management Priorities)	123
9.6.1	Manage Medium Threat Weeds (Program Maintenance)	123
9.6.2	Revegetation and Supplementary Plantings in Woodland, Scrub and Non-Remnant Vegetation (Capital/Grants Program)	123
9.6.3	Manage User-related Issues: New Tracks and Disturbances (Program Maintenance)	124
9.6.4	Manage Lower Priority User Related Issues (Program Maintenance)	124
9.6.5	Monitoring (Biodiversity Monitoring and Assessment)	124
9.7	Implications for Cultural and Heritage Values	125
10	References	126
11	Glossary	136

Tables

Table 1	Assessment against the nine Ramsar listing criteria. Criteria that the site meets are also highlighted in blue.	25
Table 2	Palaeartic migratory waders recorded in the Ramsar site and their frequency of occurrence (percentage). The eight species that the site is considered to regularly support are shaded.	48
Table 3	Numbers of species recorded at Edithvale and Seaford Wetlands and listed as threatened or migratory/marine-overfly species under the Commonwealth Environment Protection and Biodiversity Act 1999 (EPBC Act) and/ or threatened under the Victorian Flora and Fauna Guarantee Act 1988 (FFG Act) and/or classified as threatened in Victoria by DSE (2013).	57
Table 4	Number of months during which BirdLife Australia surveys at Edithvale Wetlands from September 1989 to September 2015 and Seaford Wetlands from June 1994 to September 2015 have recorded 18 Latham's Snipe or more (number of snipe recorded are given), Edithvale–Seaford Wetlands.	60
Table 5	Summary of assessment against Limits of Acceptable Change for the Edithvale-Seaford Wetlands Ramsar Site.	71
Table 6	Water quality monitoring data summary for salinity, 2009–2015, Edithvale-Seaford Wetlands.	74

Table 7	Water quality monitoring data summary for copper and zinc, 2009–2015, Edithvale-Seaford Wetlands.	76
Table 8	Summary of risk assessment for the Edithvale-Seaford Wetlands Ramsar Site.	85
Table 9	Management zones and objectives for Edithvale Wetlands.	96
Table 10	Management zones and objectives for Seaford Wetlands.	97
Table 11	Management actions to protect and enhance the ecological values of Edithvale-Seaford Wetlands. CPS = critical components, processes and services are highlighted in bold type.	101
Table 12	Summary of estimated costs to undertake priority management actions at each wetland, Edithvale-Seaford Wetlands.	106
Table 13	Environmental weeds requiring management within Edithvale Wetlands.	112
Table 14	Environmental weeds requiring management within Seaford Wetlands.	113
Table 15	Significant flora species recorded (or modelled to possibly occur) within 5 km of the Edithvale Wetlands (excludes species with negligible likelihood of occurrence).	141
Table 16	Significant flora species recorded (or modelled to possibly occur) within 5 km of the Seaford Wetlands (excludes species with negligible likelihood of occurrence).	142
Table 17	Hydrology of the wetland cells at Edithvale Wetlands (summarised from GHD 2006, KBR 2009, SKM 2011, DSE 2012 and Jacobs 2016 and updated from data provided by Paul Rees, Melbourne Water).	169
Table 18	Hydrology of the wetland cells at Seaford Wetlands (summarised from GHD 2006, KBR 2009, SKM 2011, DSE 2012 and Jacobs 2016).	171
Table 19	Likelihood	175
Table 20	Consequence	175
Table 21	Risk matrix	176
Table 22	Risk assessment for the Edithvale-Seaford Wetlands Ramsar Site. Cells highlighted in blue provide a description of the pressure / stressor that is applicable to the relevant impact pathways below	177

Figures

Figure 1	The location of the Edithvale Wetlands near Chelsea and the Seaford Wetlands to the north of Frankston.	23
Figure 2	The location of the Carrum Important Bird Area comprising Edithvale Wetlands, Seaford Wetlands, Braeside Park Wetlands, Wannarkladdin Wetlands, Boundary Road Wetland, PARCS Wetland and the Eastern Treatment Plant	28
Figure 3	Plan of the Seaford Wetlands section of the Ramsar site, showing the land owned by Melbourne Water and Frankston City Council (figure provided by Melbourne Water).	30

Figure 4	Plan of the Seaford Wetlands section of the Ramsar site, showing the proposed management areas of Melbourne Water and Frankston City Council (figure provided by Melbourne Water).	31
Figure 5	The Edithvale-Seaford Wetlands Ramsar Site Management Plan in context of other requirements for the management of Ramsar sites (adapted from DEWHA 2008).	36
Figure 6	Maximum annual abundance of waterbird guilds at Edithvale-Seaford Wetlands 1994 to 2015 (data from BirdLife Australia). Guild membership as defined by Kingsford et al. (2012).	48
Figure 7	Maximum annual abundance of Australasian Bitterns at Edithvale-Seaford Wetlands 1994 to 2015 (data from BirdLife Australia).	50
Figure 8	Ecological Vegetation Class mapping for Edithvale Wetlands.	55
Figure 9	Ecological Vegetation Class mapping for Seaford Wetland.	56
Figure 10	Map of land that has the potential to contain coastal acid sulfate soils from DSE (2010).	75
Figure 11	Former Carrum Carrum Swamp, including remnant cells (from GHD 2006).	82
Figure 12	Location of hydrological cells, drains, weirs and sedimentation ponds and direction of flows for Edithvale Wetlands (reproduced from GHD 2006).	83
Figure 13	Location of hydrological cells, drains, pipes and weirs and direction of flows for Seaford Wetlands (reproduced from GHD 2006).	84
Figure 14	The location of management zones and key monitoring features at Edithvale Wetlands. Refer to Figure 5 for detail of Ecological Vegetation Class (EVC) mapping.	98
Figure 15	The location of management zones and key monitoring features at Seaford Wetlands. Refer to Figure 6 for detail of Ecological Vegetation Class (EVC) mapping.	99
Figure 16	Seasonal water level fluctuations for Edithvale-Seaford Wetlands (source: SKM 2011)	110
Figure 17	Conceptual cross section of the embankment for revegetating wetland environments, showing terrestrial vegetation and wetland vegetation zones 1, 2 and 3.	121
Figure 18	Edithvale Wetlands revegetation zones.	194
Figure 19	Seaford Wetlands revegetation zones.	195
Plates		
Plate 1	Ungroomed (left) and groomed (right) Common Reed <i>Phragmites australis</i> at Edithvale North Wetland.	52

Appendices

Appendix 1	Edithvale-Seaford Wetlands Ramsar site land parcels, land status and land managers (from KBR 2009, DSE 2012; updated by DEPI 2013a and DELWP 2016d).	137
Appendix 2	Significant flora species potentially occurring within the Edithvale-Seaford Wetlands.	140
Appendix 3	Lists of species recorded and frequency of records of bird species recorded at Edithvale Wetlands from 1989 to 2015 and at Seaford Wetlands from 1994 to 2015 during monthly surveys undertaken by Bird Life Australia (summary of surveys from BirdLife Australia Atlas database).	143
Appendix 4	List of threatened and/or migratory species recorded at Edithvale Wetlands from 1989 to 2015 and at Seaford Wetlands from 1994 to 2015 during monthly surveys undertaken by Bird Life Australia (summary of surveys from BirdLife Australia Atlas database).	162
Appendix 5	Non-avian fauna species recorded within the Edithvale-Seaford Wetlands Ramsar Site (see Ecology Australia 2001; KBR 2009; DELWP 2016).	165
Appendix 6	Details of microbat surveys undertaken by Parson Brinckerhoff (2009 and 2010), Edithvale-Seaford Wetlands Ramsar Site.	168
Appendix 7	Hydrology of wetland cells at Edithvale-Seaford Wetlands (see Section 6.3).	169
Appendix 8	The risk assessment for threats at Edithvale-Seaford Wetlands Ramsar Site.	174
Appendix 9	Works undertaken by Melbourne Water since the production of the last Ramsar Management Plan (KBR 2009).	184
Appendix 10	Mapping of weeds of management concern at Edithvale and Seaford Wetlands, undertaken by Australian Ecosystems (2016).	186
Appendix 11	Additional weed control and revegetation information.	188
Appendix 12	Plant species suitable for revegetation in the various habitat types of Edithvale-Seaford Wetlands Ramsar Site (brackish wetlands, freshwater wetlands, Swamp Scrub and Damp Sands Herb-rich Woodland).	189
Appendix 13	Revegetation prescriptions for Edithvale-Seaford Wetlands from TBLA and Australian Ecosystems (2005).	196
Appendix 14	Edithvale-Seaford Ramsar Wetlands Desktop Cultural Heritage Assessments for the Edithvale Section (Heritage Insight 2016a) and Seaford Section (Heritage Insight 2016b).	199

Acknowledgments

We gratefully acknowledge the assistance of:

- Paul Rees from Melbourne Water;
- William Steele from Melbourne Water;
- Deane Anderson from Melbourne Water;
- Alan Clements from Melbourne Water;
- Sharon Ismale from Melbourne Water;
- Mary Catus-Wood from Melbourne Water;
- Lee James from Kingston City Council;
- David Fairbridge from Frankston City Council;
- Alan Wallis from Frankston City Council;
- Clare Warren from Frankston City Council;
- Andrew Silcocks from BirdLife Australia;
- Sean Dooley from BirdLife Australia;
- Phillippa Bailey from Friends of Edithvale-Seaford Wetlands;
- Robin Clarey from Friends of Edithvale-Seaford Wetlands;
- Alison Kuitert from Seaford Community Group; and
- Matt Dell, Katie Stevenson, Chris Bloink, John Kershaw and Jamie McMahon from Ecology Australia.

Summary Management Plan

The following table outlines prioritised management actions required to maintain and enhance biodiversity values within the Edithvale-Seaford Wetlands Ramsar Site. A map for each site displaying management zones referred to in the table is provided below. Specific details pertaining to values, threats and associated management are provided in the body of the report.

Key: MZ Management Zone

EPBC *Environment Protection and Biodiversity Conservation Act 1999*

MW Melbourne Water

KCC Kingston City Council

CPS Critical components, process and services (highlighted in **bold type**)

CaLP *Catchment and Land Protection Act 1994*

FCC Frankston City Council

Action	Legislation applicable	Management Zone	Risk (based on risk assessment)	Values/CPS	Management Category	Target	Edithvale Wetlands Estimated cost	Seaford Wetlands Estimated cost	Management Responsibility (Seaford)	Timing
Priority 1 – Legislative Obligations										
Where possible, grooming of Common Reed to maintain Ecological Character	<i>EPBC Act 1999</i>	Edith: MZ 1, 2 Seaf: MZ 1, 2	High	Waterbird diversity and abundance; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Program maintenance	Edithvale: MZ 1 groomed annually (c. 33 ha); MZ 2 groomed in a rotational pattern with return frequency determined based on fauna habitat utilisation monitoring (c. 5 ha per year if on five year rotation) Seaford: MZ 1 groomed every two (to five) years depending on access due to seasonal conditions that may prevent slashing for one or more years (c. 61 ha); MZ 2 groomed in a rotational pattern with return frequency determined based on fauna habitat utilisation monitoring (c. 8 ha per year if on five year rotation)	\$114,000 per year (\$3,000 per hectare)	\$115,500 per year (\$3,000 per hectare)	MW	Autumn, after Common Reed has flowered, but before seed set.
Conduct a machinery trial for options to cut Common Reed at Seaford Wetland where soft sediments are restricting grooming machinery	<i>EPBC Act 1999</i>	Seaf: MZ 1	High	Waterbird diversity and abundance; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Investigation	Trial of machinery options completed by Year 3 to determine if alternative mechanically grooming options are viable. This is to occur in conjunction with the above grooming regime and is expected to involve trialing various machinery to cut the Common Reed in locations with soft sediments	NA	\$12,000 one-off cost (over three years) Includes additional investigation and hire costs beyond the grooming regime above.	MW	Autumn, after Common Reed has flowered, but before seed set.
Monitor Common Reed management program performance to maintain Ecological Character	<i>EPBC Act 1999</i>	Edith: MZ 1, 2 Seaf: MZ 1, 2	-	Waterbird diversity and abundance; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Monitoring and assessment	Flora: Quadrats established/assessed in Year 1, and reassessed annually Fauna: Waterbird abundance, diversity and visitation rates assessed across grooming regimes three times per year by year 3	Flora: \$9,000 per year Fauna: \$8,000 per year	Flora: \$9,000 per year Fauna: \$8,000 per year	MW	Flora: Mar/Apr Fauna: Sep–Dec
Where possible, maintain current hydrological regime in wetlands to maintain Ecological Character	<i>EPBC Act 1999</i>	Edith: MZ 1, 2 Seaf: MZ 1, 2	Medium	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Program maintenance	Where possible, wetlands inundated in winter and early spring, with gradual drawdown to the end of December (noting that this is largely driven by rainfall and evaporation with Melbourne Water able to input minimal influence). Ephemeral areas ideally dry between January and April while deeper pools remain inundated year round.	Running costs to be determined	Running costs to be determined	MW	All year

Action	Legislation applicable	Management Zone	Risk (based on risk assessment)	Values/CPS	Management Category	Target	Edithvale Wetlands Estimated cost	Seaford Wetlands Estimated cost	Management Responsibility (Seaford)	Timing
Undertake a cost-benefit analysis of management option associated with climate change to: <ul style="list-style-type: none"> - prolong the system as a freshwater to brackish environment; and - manage and facilitate the inevitable change of the system to a more saline environment. 	<i>EPBC Act 1999</i>	Edith: MZ 1, 2 Seaf: MZ 1, 2	Extreme	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Investigation	Build upon the Jacobs (2016b) impacts of climate change assessment to develop a cost-benefit analysis, completed by Year 6 with recommendations incorporated into the revised Ramsar management plan in Year 7.	\$10,000 for one-off assessment	\$10,000 for one-off assessment	MW	Any time
Monitor abiotic and biotic changes associated with climate change: <ul style="list-style-type: none"> - Salinity and water quality - Groundwater and surface water level - Bird surveys - Floristic surveys 	<i>EPBC Act 1999</i>	Edith: MZ 1, 2 Seaf: MZ 1, 2	-	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Monitoring and assessment	Monthly surveys undertaken to document surface water levels, water quality, salinity and bird populations, Quarterly surveys undertaken to record ground water levels, Annual surveys to monitor floristic changes	Abiotic: \$12,000 per year Floristic: \$6,000 per year (note monthly bird monitoring cost is addressed below)	Abiotic: \$12,000 per year Floristic: \$6,000 per year (note monthly bird monitoring cost is addressed below)	MW	All year
Manage CaLP listed "Regionally Controlled" weeds (Tables 13 and 14), particularly *Spiny Rush	<i>CaLP Act 1994</i>	All	High	Waterbird diversity and abundance; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities	Program maintenance	Targets (eliminate, control, contain) as specified in Tables 13 and 14	\$24,000 per year	\$24,000 per year	MW and FCC	All year, 12 visits/yr
Control foxes with soft-jaw traps, maintaining fencing and destroying dens and other harbour (if present)	<i>CaLP Act 1994</i>	All	Extreme	Waterbird diversity and abundance; and threatened waterbird species. Small mammals, frogs and reptiles.	Program maintenance	Soft-jaw traps checked daily for two weeks between April and May, and data on trap success rates collected and submitted to Melbourne Water Maintain predator-proof fence at Edithvale Wetlands No active warrens/harbour, new dens/harbour fumigated and destroyed within 3 months	\$4,000 per year \$2,000–\$4,000 per year \$0–\$3,000 per year	\$4,000 per year NA \$0–\$3,000 per year	MW and FCC	April-May All year
Control rabbits and hares by destroying warrens and other harbour (if present)	<i>CaLP Act 1994</i>	All	Medium	Flora and vegetation communities.	Program maintenance	No active warrens/harbour, new warrens/harbour fumigated and destroyed within 3 months	\$0–\$3,000 per year	\$5,000–\$10,000 per year (more for the first three years)	MW and FCC	Sep
Undertake mosquito monitoring and control (if necessary)	<i>Health (Infectious Diseases) Regulations 2001 and Public Health and Wellbeing Regulations 2009</i>	Seaf: MZ1	-	Human health.	Program maintenance	Monthly dip netting for larvae and CO ₂ light trapping undertaken in spring and summer (Sept-March/April) If monitoring data indicate that larvae are in high abundance, control with ProLink Briquettes will be undertaken	NA	\$8,000 per year	MW	Sept-Feb

Action	Legislation applicable	Management Zone	Risk (based on risk assessment)	Values/CPS	Management Category	Target	Edithvale Wetlands Estimated cost	Seaford Wetlands Estimated cost	Management Responsibility (Seaford)	Timing
Monthly monitoring of bird populations and water level to detect limits of acceptable change to maintain Ecological Character	<i>EPBC Act 1999</i>	All	-	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species.	Monitoring and assessment	Bird surveys and monitoring of water levels undertaken monthly and annual reporting to Melbourne Water	\$15,000 per year (this is an existing program)	\$15,000 per year (this is an existing program)	MW	All year
Potential Acid Sulfate Soil Management	<i>EPBC Act 1999, CaLP Act 1994, Environmental Protection Act 1970, Coastal Management Act 1995, Planning and Environment Act 1986 and Crown Land (Reserves) Act 1978</i>	All, especially Seaford	Medium	Waterbird diversity and abundance; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Program maintenance	Ensure an Acid Sulfate Management Plan is prepared prior to any excavation works	\$6,000 per one-off assessment	\$6,000 per one-off assessment	MW	Any time
Implement fire management recommendations provided by Terramatrix (2013)	<i>Country Fire Authority Act 1958</i>	Edith: MZ3 Seaf: MZ3, 4	High	Flora and vegetation communities.	Program maintenance	Vegetation within the recommended Asset Protection Zones and Bushfire Moderation Zone managed accordingly from Year 1.	NA	\$8,000 per year	MW and FCC	All year
Priority 2 – Best Practice Management										
Incorporate “Downs Estate” into the Ramsar boundary	-	Seaf: MZ 5	-	Physical habitat for waterbirds;	Capital/Grants Program	Down’s Estate incorporated into Ramsar site by Year 2	NA	\$5,000 per one-off investigation	MW and FCC	Any time
Manage high threat weeds (Tables 13 and 14)	-	All	High	Flora and vegetation communities.	Program maintenance	Targets (eliminate, control, contain) as specified in Tables 13 and 14	\$8,000 per year	\$12,000 per year	MW and FCC	All year, 4-6 visits/yr
Maintain current dog access restrictions (i.e. no dogs in wetlands, on-leash in buffers), install interpretative signage to educate the public of dog and cat impacts	-	All	-	Waterbird diversity and abundance; waterbird breeding; and threatened waterbird species.	Capital/Grants program	Signage installed by Year 2. These can be in the form of regulatory and educational interpretative signage.	\$5,000–\$10,000 for signage/letter drop	\$5,000–\$10,000 for one-off signage/letter drop	MW, KCC and FCC	Any time
Distribute letters to landholders in the local area regarding cat and dog impacts	-	All	Extreme	Waterbird diversity and abundance; waterbird breeding; and threatened waterbird species. Small mammals, frogs and reptiles.	Program maintenance	Letter distributed in Years 2, 5 and 8 to all registered cat and dog owners within 500 m of either wetland.	\$5,000 per year	\$6,000 per year	MW, KCC and FCC	All year
If incursions occurs in future, undertake pig trapping to remove released pigs (as had occurred at Seaford Wetland)	-	Seaford: All	Medium	Flora and vegetation communities	Program maintenance	Pig(s) removed within 6 months of confirming occurrence	NA	NA	MW and FCC	All year
Check nest boxes, document occupants and remove Common Myna nests and bee hives	-	All	-	Waterbird breeding.	Program maintenance	All nest boxes check and cleaned in May each year. Occupancy of nest boxes recorded and submitted to Melbourne Water.	\$1,600 per year	\$1,600 per year	MW and FCC	May

Action	Legislation applicable	Management Zone	Risk (based on risk assessment)	Values/CPS	Management Category	Target	Edithvale Wetlands Estimated cost	Seaford Wetlands Estimated cost	Management Responsibility (Seaford)	Timing
Revegetation of areas disturbed by management works (e.g. weed control, den destruction)	-	All	-	Flora and vegetation communities	Program maintenance	Bare ground revegetated within two months; all plant losses replaced with 90% survival to year 7; all guards removed once plants are established (c. 1-3 years depending on species and health of plant). To be undertaken in accordance with the species lists provided in Appendix 12 and the revegetation prescriptions prepared by TBLA and Australian Ecosystems (2005; Appendix 13)	Guarded: \$11/plant Unguarded: \$3/plant Maintenance: \$2/plant	Guarded: \$11/plant Unguarded: \$3/plant Maintenance: \$2/plant	MW and FCC	Ideally May-Oct
Consolidate Frankston City Council and Melbourne Water boundaries at Seaford Wetland (as per Section 2.5)	-	Seaf: All	-	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Capital/Grants program	Divide Seaford Wetland manager boundaries so that Melbourne Water manages all wetland areas and Frankston City Council manages all the exterior areas.	NA	\$5,000 for one-off investigation	MW and FCC	Anytime
At Seaford Wetland, investigate feasibility of extending the path through Downs Estate and along the northern boundary to create a complete loop track	-	Seaf: MZ 3, 5	-	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Investigation	Investigation undertaken by year 1, and findings implemented by year 2	NA	\$4,000 for one off assessment	MW and FCC	Anytime
Maintain fencing around the wetlands	-	Edith: MZ3 Seaf: MZ2, 3, 4, 5	-	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Program maintenance	Undertake fence checks twice per year and repair breaches.	\$1,000 per year (plus repairs)	\$1,000 per year (plus repairs)	MW and FCC	Anytime
Consider the potential need for fencing or seasonal closure of the internal path at Seaford Wetland, and fencing at Down's Estate if included in the Ramsar site (see proposed fencing on Figure 4).	-	Seaf: MZ 1, 2, 5	-	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Investigation	Investigation undertaken by year 2, and findings completed by year 4.	NA	\$3,000 for one-off investigation	MW and FCC	Anytime
Photo-point monitoring	-	All	-	Flora and vegetation communities	Program maintenance	Monitoring of photo-points undertaken every year	\$1,000 per year	\$1,000 per year	MW	In line with previous monitoring (October and March)
Native vegetation condition monitoring	-	All	-	Flora and vegetation communities	Monitoring and assessment	Habitat Hectare Assessment or other suitable monitoring approach collected at Year 5	\$9,000 for one-off assessment	\$9,000 for one-off assessment	MW	Sep-Dec
Monitor pest plant control performance	-	All	-	Flora and vegetation communities	Program maintenance	Distribution and cover of weeds of management concern assessed each year	\$2,000 per year	\$2,000 per year	MW and FCC	Sep-Nov

Action	Legislation applicable	Management Zone	Risk (based on risk assessment)	Values/CPS	Management Category	Target	Edithvale Wetlands Estimated cost	Seaford Wetlands Estimated cost	Management Responsibility (Seaford)	Timing
Monitor pest animal (rabbit/hare/fox) control performance	-	All	-	Waterbird diversity and abundance; and threatened waterbird species. Flora and vegetation communities	Program maintenance	Inspection for warrens / dens at least once per year (if active warrens found undertake more frequent inspections) BirdLife Australia to continue to record foxes and cats observed during monthly bird surveys	\$1,000 per year (more if active warrens found)	\$1,000 per year (more if active warrens found)	MW and FCC	Sep–Oct
Monitor Eastern Grey Kangaroo population	<i>Wildlife Act 1975 and Wildlife Regulations 2002</i>	Edith (south): MZ 1–3	Low-medium	Flora and vegetation communities	Program maintenance	Undertake annual monitoring of the Eastern Grey Kangaroo population using the sweep count procedure to ensure population is maintained between 16-27 kangaroos (see EcoPlan Australia 2015)	\$5,000 per year	NA	MW	March-April
Priority 3 – Other Management Priorities										
Manage medium threat weeds (Tables 13 and 14)	-	All	High-Medium	Flora and vegetation communities	Program maintenance	Targets (eliminate, control, contain) as specified in Tables 13 and 14	\$8,000 per year	\$8,000 per year	MW and FCC	All year, 4 visits/yr
Investigate potential for using fire as a management tool at Seaford Wetlands for #Coast Tea-tree control in MZ3.	-	Seaf: MZ 1–3	High	Flora and vegetation communities	Program maintenance	Feasibility of using fire as a management tool at Seaford for #Coast Tea-tree investigated by Year 3 and implemented accordingly.	NA	\$5,000 for one-off investigation	MW and FCC	Any time
Revegetation and supplementary plantings in Damp Sands Herb-rich Woodland and Swamp Scrub vegetation	-	Edith: MZ3 Seaf: MZ 3, 4	-	Flora and vegetation communities	Capital/Grants program	- Clustered plantings of shrubby and robust ground layer species in the Damp Sands Herb-rich Woodland vegetation with primarily mown lawn understorey - Supplementary and expansion plantings of Swamp Scrub - Revegetation of non-remnant areas opposite Seaford North Primary School - All plant losses replaced with 90% survival to year 7; all guards removed once plants are established (c. 1-3 years depending on species and health of plant). - To be undertaken in accordance with the species lists provided in Appendix 12 and the revegetation prescriptions prepared by TBLA and Australian Ecosystems (2005; Appendix 13)	Guarded: \$11/plant Unguarded: \$3/plant Maintenance: \$2/plant	Guarded: \$11/plant Unguarded: \$3/plant Maintenance: \$2/plant	FCC	May-Oct
User related issues: rehabilitate unauthorised tracks and disturbances	-	All	-	Flora and vegetation communities	Program maintenance	Negligible unauthorised tracks or disturbances by year 2, new disturbances rehabilitated within 2 months of identification	\$1,000–\$5,000 per incursion (depends on extent of damage)	\$1,000–\$5,000 per incursion (depends on extent of damage)	MW and FCC	All year
User related issues: undertake general litter collection and clear litter traps	-	All	Medium	Physical habitat for waterbirds. Flora and vegetation communities	Program maintenance	Negligible litter observed onsite and in litter traps	\$6,000–\$8000 per year	\$6,000–\$8000 per year	MW and FCC	All year
Mapping and monitoring of significant flora species (Appendix 2)	-	All	-	Flora and vegetation communities	Monitoring and assessment	Distribution and population size mapped and compared in Year 5	\$7,000 for one-off assessment	\$7,000 for one-off assessment	MW and FCC	Oct–Nov
Targeted surveys for reptiles (Swamp Skink)	-	Edith: MZ 1, 2 Seaf: MZ 1, 2	-	Reptiles.	Monitoring and assessment	One targeted survey completed by Year 3	\$8,000 per one-off assessment	\$8,000 per one-off assessment	MW and FCC	Nov-Mar
Fish survey to document fish species occurrence	-	Edith: MZ 1, 2 Seaf: MZ 1, 2	-	Fish.	Monitoring and assessment	Fish survey completed by Year 4	\$6,000 per one-off assessment	\$6,000 per one-off assessment	MW and FCC	Dec–May


Action	Legislation applicable	Management Zone	Risk (based on risk assessment)	Values/CPS	Management Category	Target	Edithvale Wetlands Estimated cost	Seaford Wetlands Estimated cost	Management Responsibility (Seaford)	Timing
Invertebrate survey, with the first priority on aquatic invertebrates and the lower priority on terrestrial invertebrates	-	All	-	Invertebrates	Monitoring and assessment	Invertebrate survey undertaken by Year 6	\$6,000 per one-off assessment	\$6,000 per one-off assessment	MW	Spring


Costs do not consider CPI.

Summary of estimated costs to undertake priority management actions at Edithvale and Seaford Wetlands:

Site	Priority	Estimated costs per year ^	Estimated costs for one-off assessments
Edithvale Wetland	Priority 1 management actions	\$194,000-\$202,000	\$16,000
	Priority 2 management actions	\$29,600-\$34,600	\$9,000
	Priority 3 management actions	\$15,000-\$21,000	\$27,000
Seaford Wetland	Priority 1 management actions	\$214,500-\$222,500	\$28,000
	Priority 2 management actions	\$29,600-\$34,600	\$26,000
	Priority 3 management actions	\$15,000-\$21,000	\$32,000

^ Does not include revegetation costs of \$11 / guarded plant or \$3 / unguarded plant, and \$2 / plant for maintenance.





1 Introduction

1.1 Background

The Edithvale-Seaford Wetlands represent remnant ecosystems of the once extensive Carrum Carrum Swamp which was located on the eastern shoreline of Port Phillip Bay, separated from the sea by a beach ridge-dune barrier system. The wetlands have long been recognised as supporting important habitat for birds (Carter 1975; Watkins 1993; ANCA 1996; Ecology Australia 2000; GHD 2006). The significance of the wetlands was acknowledged by their inclusion in the Directory of Important Wetlands in Australia (ANCA 1996). Further recognition of the diversity of values led to their listing as wetlands of international significance in 2001 under the Ramsar Convention (KBR 2009), shortly after the preparation of the first management plan for the site (Ecology Australia 2000).

Edithvale Wetlands are owned and managed by Melbourne Water, whilst Seaford Wetlands are owned and managed jointly by Melbourne Water and Frankston City Council (KBR 2009; DSE 2012). In accordance with the Australian Ramsar Management Principles, Melbourne Water needs to conduct seven year reviews of existing Ramsar Wetland Management Plans. The existing, 2009, Edithvale-Seaford Ramsar Site Management Plan (KBR 2009) is due for a comprehensive review with preparation of this new Management Plan to guide best practice management of these important wetland sites. This management plan will guide management at the site from 2016 until a new plan is prepared in 2023.

Furthermore, the sites contain a number of important biodiversity values that sees them included in Melbourne Water's Sites of Biodiversity Significance (SoBS) register (Melbourne Water 2013a and 2013b). Thus, the Management Plan is cognisant of Melbourne Water's requirements under the SoBS program.

1.2 Objectives of the Management Plan

Consistent with requirements of the Convention, the primary purpose of the Edithvale-Seaford Wetlands Ramsar Management Plan is to maintain ecological character and promote wise use of the site. Wise use is defined by the Convention as (Ramsar Convention 2005):

- *“the maintenance of their ecological character, achieved through the implementation of ecosystem approaches, within the context of sustainable development”.*

More plainly, this means balancing conservation with sustainable use of wetlands that benefits both people and nature. The primary objective of this Edithvale-Seaford Wetlands Ramsar Site Management Plan is:

- ***“To maintain, and where possible improve, the ecological character of the Edithvale-Seaford Wetlands Ramsar site and promote wise use”.***

More specifically, the objectives of the Management Plan are to:

- fulfil international, federal, state and local government requirements for Ramsar management plans;
- incorporate the views of local community groups;

- provide a comprehensive description of the Ramsar site;
- discuss the legislative and policy framework that may affect management of the Ramsar site;
- describe the values of the site, including those pertaining to the “Ecological Character” of the wetlands, those that satisfy the criteria for Ramsar listing, and other values, such as vegetation, wetland types, drainage function, cultural heritage and social amenity;
- identify management zones associated with the wetlands, including the values of the different zones;
- identify threats associated with the different management zones;
- undertake a risk assessment of the threats;
- propose prioritised management actions to address the threats and maintain the unique biodiversity values of the wetlands, according to zones;
- outline management requirements of the different zones; and
- protect cultural heritage values.

1.3 Ramsar Management Policies and Guidelines

The plan structure and contents of this Ramsar Site Management Plan is in-line with relevant Ramsar legislation, government policies and strategies, including the requirements of:

- the Australian Government (2000) - Australian Ramsar Management Principles;
- Department of Natural Resources and Environment (DNRE) (2002) - Management of Victoria’s Ramsar Wetlands Strategic Directions Statement;
- Ramsar Contracting Parties (2002) - New Guidelines for Management Planning for Ramsar Sites and Other Wetlands. Adopted by Resolution VIII.14 (2002) of the Ramsar Convention;
- Thomas and Middleton (2003) - Guidelines for Management Planning of Protected Areas; and
- Chatterjee A, Phillips B and Stroud D (2008) - Wetland Management Planning: A Guide for Site Managers.

1.4 Consultative Framework

This Management Plan has been developed collaboratively in order to represent the interests of the agencies and community groups, involve interested parties in the decision making process regarding future management of the wetlands and identify areas or actions for continuing community input and involvement with implementation of the Plan. An inception meeting (19 November 2015) and site familiarisation visit with staff from Melbourne Water (23 February 2016) were undertaken. Ecology Australia also met the Edithvale-Seaford Wetlands Community Liaison Committee on 24 February 2016. A workshop was run by Jenny Hale with stakeholder involvement on 8 March 2016 and a site visit to areas identified during the workshop was undertaken with the stakeholders on 16 March 2016.

Stakeholders comprised of staff or members from Melbourne Water, Kingston City Council, Frankston City Council, BirdLife Australia, Friends of Edithvale-Seaford Wetlands, Seaford Community Group and Aspendale Gardens Residents Association.

All comments received during the consultative phase were considered in finalising the Management Plan.

The draft Ramsar Management Plan was distributed by Melbourne Water to all of the stakeholders mentioned above as well as several additional stakeholders - Downs Estate Community Group, Department of Environment, Land, Water and Planning (DELWP). All of these stakeholders were encouraged to provide feedback on the plan and were given six weeks to do so.

The Management Plan also provides for continuing community involvement in implementation of the Plan through identification of areas of management that can be undertaken by the community in Section 9.

2 Ramsar Site Description

2.1 Location

The Edithvale-Seafood Wetlands Ramsar Site is located in Melbourne's south eastern suburbs, c. 30 km from the Melbourne Central Business District (CBD). The site is comprised of two separate wetlands: Edithvale (104 ha) and Seafood (158 ha) (Figure 1). The site is owned predominantly by Melbourne Water with some sections of Seafood Wetlands, including "Downs Estate", owned by the City of Frankston (KBR 2009; DSE 2012; DEPI 2013a).

The Edithvale-Seafood Wetlands Ramsar Site is located in the Gippsland Plain bioregion, which extends from Melbourne's CBD in the west to Lakes Entrance in the west and to Moe in the north (DSE 2012). The bioregion comprises coastal and lowland alluvial plains and is characterised by generally flat to undulating terrain, vegetated in parts with open grassy forest and areas of Swamp Scrub (DSE 2012).

The area experiences a temperate climate with an average annual rainfall of 709 mm recorded at Moorabbin Airport (Station Number 86077; Lat. 37.98° S; Lon. 145.10° E; Elevation: 12 m asl; data 1950-current), the closest weather station with long-term rainfall data. The wettest month is May with an average of 69.5 mm and the driest month is January with an average of 44.2 mm. The monthly mean maximum temperatures at Moorabbin Airport (from 1971 to current) range from 26.1°C in February to 13.7°C in July. The monthly mean minimum temperatures at Moorabbin Airport range from 14.4°C in February to 6.2°C in July (Bureau of Meteorology, 2016).

2.2 Wetland Description

The Edithvale-Seafood Wetlands Ramsar Site is the only Victorian Ramsar site located in an urban landscape. The Ramsar site has experienced a long history of disturbance, and subsequently, the condition of the vegetation, hydrology and water quality has been modified (Ecology Australia 2001; KBR 2009; DSE 2012; Section 6).

The distribution and occurrence of Ecological Vegetation Classes (EVCs) identified at Edithvale and Seafood Wetlands is in a constant state of flux primarily responding to water levels and salinity; the following EVCs have been identified (TBLD and Australian Ecosystems 2005; Australian Ecosystems 2011a, 2015; Section 4.1.5):

- Damp Sands Herb-rich Woodland (EVC 3);
- Swamp Scrub (EVC 53);
- Tall Marsh (EVC 821);
- Brackish Aggregate Wetland (EVC 656);
- Brackish Aquatic Herbland (EVC 537)
- Plains Sedgy Wetland (EVC 647);
- Aquatic Herbland (EVC 653);
- Brackish Herbland (EVC 538); and
- Wet Saltmarsh (EVC 9).




Figure 1 The location of the Edithvale Wetlands near Chelsea and the Seaford Wetlands to the north of Frankston.

Tall Marsh is the dominant wetland vegetation throughout both Edithvale and Seaford Wetlands. It is highly invasive of other habitats, especially of important mudflat areas for migratory waders, but Melbourne Water management is attempting to reduce the extent of invasion (Australian Ecosystems 2015; Greet and Rees 2015). It provides habitat for a number of waterbird species and marshland-dependent passerine species (Sections 4.1.3, 4.1.4 and 4.1.6; Birds Australia 2004; BirdLife Australia 2015a). Damp Sands Herb-rich Woodland surrounds Seaford Wetlands and provides habitat for terrestrial birds, mammals and reptiles (Birds Australia 2004; KBR 2009; Australian Ecosystems 2015).

2.3 Wetland Types

Victorian wetlands have been mapped and classified according to a Victorian Wetland Classification (DELWP, in prep.) in the WETLAND_CURRENT spatial layer. This identifies four wetland types in the Ramsar site:

- Fresh, intermittent, sedge/grass/forb – 116 hectares
- Fresh seasonal/episodic unknown vegetation – 11 hectares
- Fresh seasonal / episodic, forest/woodland – 4 hectares
- Fresh, intermittent, open water – 1.4 hectares.

However, knowledge of the site, and mapped EVCs indicate that there are areas of saline / brackish wetland on site (EVC 656 - Brackish wetland Aggregate; EVC 538 - Brackish Herbland; and EVC 537 - Brackish Aquatic Herbland).

Using all available information, this translates into four wetland types according to the Ramsar wetland classification, within the Edithvale-Seaford Wetlands Ramsar site. In order of dominance, these are:

- Ts - Seasonal / intermittent freshwater marshes/pools on inorganic soils;
- Ss - Seasonal/intermittent saline/brackish/alkaline marshes/pools;
- Xf - Freshwater, tree-dominated wetlands;
- P - Seasonal/ intermittent freshwater lakes.

2.4 Wetland Significance and Listings

2.4.1 Ramsar Criteria

When the Edithvale-Seaford Wetlands were first nominated in 2001, the original documents considered that the site met three of the then eight criteria (see Brett Lane and Associates 2001). The criteria under which a Ramsar site can be designated have gone through a series of changes, with the most recent major revisions occurring at the 9th Ramsar Conference in Kampala, Uganda 2005, when a ninth criterion was added. The most recent assessment of the site against Ramsar criteria (Hale 2016) indicated that there is insufficient evidence to suggest that the site met criteria 1 and 3, but continues to meet three criteria (2, 4 and 6; Table 1).

Table 1 Assessment against the nine Ramsar listing criteria. Criteria that the site meets are also highlighted in blue.

Criteria	Justification	Meets Criterion (Yes/No)
1. Contains a representative, rare, or unique example of a natural or near-natural wetland type found within the appropriate biogeographic region.	<p>The appropriate bioregion for the site is the South-east Coastal Drainage Division which includes all of the coastline of Victoria and New South Wales and a small portion of the South Australian coast (Department of the Environment, Water, Heritage and the Arts 2008). There is no comprehensive wetland inventory for this bioregion. As such the application of the terms “representative” and “rare” are difficult. In terms of “representative” advice from the Convention (Ramsar 2009) is that contracting parties should select the “best examples” of each wetland type within a bioregion and that the wetlands should be in “near-natural” condition.</p> <p>The Edithvale-Seaford wetlands are remnants of what was once the Carrum Carrum Swamp, a large freshwater wetland, largely drained in the late 19th century. The wetland types within the site may be locally important, however, they are currently highly modified and part of a regional drainage system for storing stormwater from surrounding urban landscapes. It is difficult to make the argument that these sites are rare, representative or near-natural. As such, the site does not meet this criterion and did not meet it at the time of listing.</p>	No
2. Supports vulnerable, endangered, or critically endangered species or threatened ecological communities.	<p>The site regularly supports two wetland dependent fauna species listed under the EPBC Act and / or IUCN Red List:</p> <ul style="list-style-type: none"> • Australasian Bittern <i>Botaurus poiciloptilus</i> – Endangered • Curlew Sandpiper <i>Calidris ferruginea</i> – Critically Endangered 	Yes
3. Supports populations of plant and/or animal species important for maintaining the biological diversity of a particular biogeographic region.	<p>Guidance from the Convention indicates that this criterion should be applied to “hotspots” of biological diversity and centres of endemism within a biogeographical region. The Edithvale-Seaford Wetlands is within a very large bioregion (southeast coast drainage division) and there is no evidence to suggest it is more species rich than other wetlands and Ramsar sites in this region.</p>	No
4. Supports plant and/or animal species at a critical stage in their life cycles, or provides refuge during adverse conditions.	<p>The Edithvale-Seaford Ramsar Site regularly supports eight international migratory waterbirds, provides habitat for many breeding birds and acts as a drought refuge in an urban landscape.</p>	Yes
5. Regularly supports 20,000 or more waterbirds.	<p>Total annual maximum counts from Edithvale-Seaford Wetlands are less than 20,000 (average is approximately 5000).</p>	No
6: Regularly supports 1% of the individuals in a population of one species or subspecies of waterbird.	<p>Counts above 1% of the estimated population of two species are regularly recorded within the site: Australasian Bittern <i>Botaurus poiciloptilus</i> and Sharp-tailed Sandpiper <i>Calidris acuminata</i>.</p>	Yes

Criteria	Justification	Meets Criterion (Yes/No)
7. Supports a significant proportion of indigenous fish subspecies, species or families, life-history stages, species interactions and/or populations that are representative of wetland benefits and/or values and thereby contributes to global biological diversity.	The site is not considered important for native fish.	No
8: An important source of food for fishes, spawning ground, nursery and/or migration path on which fish stocks, either within the wetland or elsewhere, depend.	The site is not considered important for native fish.	No
9: Regularly supports one percent of the individuals in a population of one species or subspecies of wetland-dependent non-avian animal species.	No evidence to suggest that this criterion is met.	No

2.4.2 Other listings/classifications

Edithvale-Seaford Wetlands are also:

- Considered to be a site of State and International Treaty Zoological Significance in the southeast of Melbourne and Mornington Peninsula by DSE (2004);
- included in the Directory of Important Wetlands (ANCA 1996; KBR 2009);
- subject to a number of migratory bird bilateral agreements and conventions (Section 3.1.2);
- identified as a high value Melbourne Water SoBS (Section 3.4.3);
- part of the Carrum Important Bird Area (IBA) program led by BirdLife Australia (see below).

Seaford Wetlands was also identified as:

- a site of International importance for the Sharp-tailed Sandpiper by Watkins (1993), in that the area supports (or supported) 1% or more of the individuals in the East Asian – Australasian Flyway population.

Carrum Wetlands Important Bird and Biodiversity Area

The Important Bird and Biodiversity Areas (IBAs) program is an international non-governmental conservation scheme lead by BirdLife International Partners, such as BirdLife Australia. IBAs are (BirdLife Australia 2016b):

- sites of international importance for bird conservation; and
- small enough to be practical targets for conservation management, but large enough to meet the global IBA criteria.

The Australian IBA program aims to protect a network of sites critical for the conservation of Australia's birds by:

- promoting IBAs as a tool for biodiversity conservation planning;
- encouraging government to prioritise conservation at IBAs (e.g. in grant-giving schemes); and
- encouraging and facilitating local community-based groups and land-owners to manage land sustainably and conserve key bird species.

Edithvale and Seaford Wetlands, together with Braeside Park Wetlands, Wannarkladdin Wetlands, Boundary Road Wetland, PARCS Wetland and the Eastern Treatment Plant constitute the Carrum Wetlands IBA (Figure 2; BirdLife Australia 2016b). The Carrum Wetlands IBA regularly supports more than 1% of the world populations of Sharp-tailed Sandpiper, Blue-billed Duck and Chestnut Teal and is an important non-breeding refuge for the endangered Australasian Bittern (BirdLife Australia 2016b).

Globally-important bird populations in this IBA identified from monitoring between 1989 and 2008 recorded (BirdLife Australia 2016b):

- Chestnut Teal 0 - 1962 individuals;
- Blue-billed Duck 0 - 510 individuals;
- Australasian Bittern 0 - 14 individuals; and
- Sharp-tailed Sandpiper 0 - 5839 individuals.




Figure 2 The location of the Carrum Important Bird Area comprising Edithvale Wetlands, Seaford Wetlands, Braeside Park Wetlands, Wannarkladdin Wetlands, Boundary Road Wetland, PARCS Wetland and the Eastern Treatment Plant

2.5 Land Tenure and Management

The Edithvale Wetlands site comprises freehold land parcels (allotments) owned and managed by Melbourne Water (KBR 2009; DSE 2012; DEPI 2013a).

Seaford Wetlands consists of freehold land parcels owned and managed by Melbourne Water and a Crown Land conservation reserve to the east of Seaford Wetlands that is managed by Melbourne Water under a Committee of Management (KBR 2009; DSE 2012). Frankston City Council owns a significant portion of the parcels of land surrounding Seaford Wetlands, particularly the dryland areas (see Figure 3). There is an informal arrangement in terms of management with Melbourne Water notionally managing inside the internal fence and Frankston City Council notionally managing outside the internal fence (see Figure 4; Paul Rees, Melbourne Water, pers. comm.).

Land tenure and management is described in KBR (2009) and DSE (2012) with land parcel/lot numbers updated by DELWP (2016d) provided in Appendix 1.

'Downs Estate' (Lot 93\PP3025), in the northeast of Seaford Wetlands, was acquired by Frankston City Council just prior to the preparation of the last Management Plan for the Ramsar site (KBR 2009). It is freehold land managed by Frankston City Council and is situated outside of the Ramsar boundary. However, Council's Environment and Planning Department and Melbourne Water are interested in including the land within the Ramsar boundary.




Figure 3 Plan of the Seaford Wetlands section of the Ramsar site, showing the land owned by Melbourne Water and Frankston City Council (figure provided by Melbourne Water).




Figure 4 Plan of the Seaford Wetlands section of the Ramsar site, showing the proposed management areas of Melbourne Water and Frankston City Council (figure provided by Melbourne Water).

2.6 Landscape Setting

The Edithvale-Seaford Wetlands Ramsar Site is the only Victorian Ramsar site and one of a few Australian Ramsar sites located in an urban landscape and support important areas of open space used by the public for recreation (Ecology Australia 2000; KBR 2009; DSE 2012).

Edithvale Wetlands are surrounded by freehold urban development to the north, east and south that form the suburbs of Aspendale Heights and Chelsea Heights (KBR 2009; DSE 2012). Centre Swamp Drain and an associated drainage reserve occur along the western boundary of the Edithvale Wetlands and runs from north to south between Mordialloc Creek in the north and the Patterson River in the south. A recreation reserve, the Edithvale Common, adjoins the Drain and drainage reserve to the west of Edithvale North Wetlands and supports two wetlands which supported good habitat for waterbirds (e.g. Latham's Snipe and ducks) and frogs on an annual basis prior to alterations/improvements to drainage; it now supports good waterbird and frog habitat only during very wet years (Ecology Australia 2001; KBR 2009; SKM 2011; Section 6.3.2; Paul Rees and Will Steele, Melbourne Water, pers. comm.). Kingston City Council is investigating alterations to the local drainage system to alleviate flooding in nearby streets, by providing water more frequently to the low lying areas in Edithvale Common (Paul Rees, Melbourne Water, pers. comm.). Rossdale Golf Course adjoins the Drain and drainage reserve to the northwest corner of Edithvale North Wetlands, whilst Chelsea Public Golf Course adjoins the northern half of Edithvale South Wetlands. Residential development adjoins the Drain and drainage reserve to the southwest of Edithvale South Wetlands (i.e. south of Chelsea Public Gold Course). The block of land to the south of Edithvale South Wetland and bordering the northern side of First Avenue and the western side of houses along Norman Avenue (c. 4.0 ha in size) may be developed and Melbourne Water will potentially acquire a portion that is covered by the LSIO (Paul Rees, Melbourne Water, pers. comm.).

Eel Race Creek borders the northern end of Seaford Wetlands (Ecology Australia 2000; KBR 2009). Freehold urban development and undeveloped land surrounds Seaford Wetlands, to the west, south, east and northeast, including the Patterson River Secondary College, residential development and Seaford North Primary School in the northwest, residential development to the west, Austin Road to the south, Francis Street and Old Wells Road to the east, and the Frankston City Motorcycle Park to the east in the northern half of the site (Ecology Australia 2000; KBR 2009).

'Downs Estate' in the northeast of Seaford Wetlands, is freehold land managed by Frankston City Council that is situated outside of the Ramsar boundary. However, Council's Environment and Planning Department and Melbourne Water are interested in including the land within the Ramsar boundary (see Sections 2.5 and 4.1.12).

Prohibitive fencing with access gates for maintenance vehicles and pedestrians have been installed to separate sensitive wetland habitats from publically accessible dryland and recreation areas at Edithvale and Seaford Wetlands (KBR 2009; Paul Rees, Melbourne Water pers. comm.).

The surrounding urban landscape poses considerable challenges for management and protection of the significant values of the Ramsar site. Management issues associated with the urban landscape include reduced stormwater quality, disturbance of wildlife, predation of wildlife by domestic animals, weed invasion and litter (e.g. Brett Lane and Associates 2001; Ecology Australia 2000; KBR 2009; DSE 2012).

2.7 Catchment Setting

The Edithvale-Seaford Wetlands are located in the Bunyip River Basin in the Southeast Coast Drainage Division (KBR 2009; DSE 2012). The Southeast Coast Drainage Division extends along the coast of Australia from south-eastern Queensland to south-eastern South Australia. The Bunyip River Basin occupies some 389,033 ha, with land use divided between urban development and agriculture (KBR 2009).

Within the Bunyip River Basin, Edithvale Wetlands and Seaford Wetlands occur in the Dandenong Catchment (c. 85,500 ha). Dandenong Creek is a major waterway within the catchment, with headwaters in the Dandenong Ranges National Park. At the time of European Settlement, Carrum Carrum Swamp received water largely from Dandenong Creek and Eumemmering Creek, with contributions from catchments that drain to Mordialloc Creek and Boggy Creek (GHD 2006). However, the hydrology of the Edithvale-Seaford Wetlands has been significantly altered since European Settlement, such that (GHD 2006; KBR 2009; SKM 2011; DSE 2012; Section 6.3):

- Seaford Wetlands has been disconnected from the Boggy Creek catchment and largely receive inflows from drains and groundwater intrusions; and
- Edithvale Wetlands receive water largely from local stormwater inlets, with groundwater intrusions also occurring.

2.8 Local Government

The Edithvale-Seaford Wetlands Ramsar site intersects both the southern extent of the Kingston City Council and the northern extent of Frankston City Council local government areas (Figures 1, 8 and 9; Brett Lane and Associates 2001; Ecology Australia 2000; KBR 2009; DSE 2012): Edithvale Wetlands are located within the Kingston City Council local government area; and Seaford Wetlands are situated within Frankston City Council local government area.

3 Legislation and Policy Framework

3.1 International Legislation and Policy

3.1.1 The Ramsar Convention

Ecological Character

The Convention on Wetlands of International Importance, otherwise known as the Ramsar Convention, was signed in Ramsar Iran in 1971 and came into force in 1975. It provides the framework for local, regional and national actions, and international cooperation, for the conservation and wise use of wetlands. Wetlands of International Importance are selected on the basis of their international significance in terms of ecology, botany, zoology, limnology and/or hydrology. Australia was one of the first countries to sign the Convention, and there are now 169 Contracting Parties and over 2000 designated wetlands of international importance. The mission of the Ramsar Convention is (Ramsar Convention 2016):

“Conservation and wise use of all wetlands through local and national actions and international cooperation, as a contribution towards achieving sustainable development throughout the world.”

In 1997 the Convention identified three pillars for the conservation and protection of wetlands:

- i) the wise use of all wetlands through national plans, policies and legislation, management actions and public education;
- ii) the designation and sustainable management of suitable wetlands for inclusion on the list of Wetlands of International Importance; and
- iii) international cooperation on transboundary wetlands and shared species.

Under the terms of the Convention contracting parties nominate wetlands to be designated as Wetlands of International Importance, with nominated sites required to meet at least one of nine listing criteria. The act of designating a wetland as a Ramsar site carries with it certain obligations, including managing the site to maintain its ‘ecological character’ and to have procedures in place to detect if any threatening processes are likely to, or have altered, the ‘ecological character’. The Ramsar Convention has defined “ecological character” and “change in ecological character” as (Ramsar Convention 2005):

- *“Ecological character is the combination of the ecosystem components, processes and benefits/services [CPS] that characterise the wetlands at a given point in time”; and*
- *“...change in ecological character is the human induced adverse alteration of any ecosystem component, process and or ecosystem benefit/service.”*

Ramsar documentation

Ramsar site management to maintain ecological character is reliant on a number of key documents and processes as illustrated in Figure 5. The three key documents are:

Ramsar Information Sheet (RIS) - compiled for each site it documents the essential information related to the site and its management. The Administrative Authority of each Contracting Party submits the RIS to the Ramsar Secretariat. In the case of Australia this is the Australian Government Department of the

Environment (DoE). The Parties have committed to providing updated RIS information for their Ramsar sites every six years, or on the occasion of any significant change in a site's ecological character. The most recent RIS for Edithvale-Seaford Wetlands was compiled in 2001 (Brett Lane and Associates 2001) and can be obtained from the DoE website (<http://www.environment.gov.au/cgi-bin/wetlands/ramsardetails.pl?refcode=57#>). This RIS is currently being updated (Hale, in pre.) with a revised RIS due in mid-2016. Once formally approved this revised RIS will be available from the Ramsar Information Service (<https://rsis.ramsar.org/>).

Ecological Character Description (ECD) – provides a more detailed and quantitative description of ecological character for a Ramsar site. The ECD for Edithvale-Seaford (DSE 2012) was written in 2008, prior to the development of guidelines for ECDs (DEWHA 2008). It can be accessed from the DoE website (<http://www.environment.gov.au/cgi-bin/wetlands/ramsardetails.pl?refcode=57#>).

An update to the ECD is being drafted (Hale 2016), which identifies critical components, processes and services and sets limits for acceptable change.

Management plan – documents the management strategies required to protect and restore the ecological character of a Ramsar site. In Australia, the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) establishes the framework for management of Australian Ramsar sites, and Schedule 6 of Environment Protection and Biodiversity Conservation Regulations 2000 outlines the principles relevant to the preparation of Ramsar site management plans (Text Box 1).

Ramsar Rolling Review – The Department of Environment has developed a three-year Ramsar Rolling Review program for reporting the status of the ecological character of Australia's Ramsar sites. The broad aims of the Ramsar Rolling Review (DELWP 2016e) program are to:

- Review and report on the status of the ecological character of Australia's Ramsar sites.
- Be a tool to assist managing sites in order to maintain their ecological character, improving links between ecological character, site management plans and monitoring programs for critical components, processes and services and associated threats.
- Provide input to a database of baseline and threat data.
- Record updates as knowledge gaps are addressed and refine Limits of Acceptable Change.
- Highlight issues and facilitate assessment of a potential change of character, focussing on proactive management before the situation requires notification.
- Identify broad trends or common threats across site and jurisdiction boundaries.

3.1.2 Migratory Bird Bilateral Agreements and Conventions

Australia is party to a number of bilateral agreements, initiatives and conventions for the conservation of migratory birds, which are relevant to the Edithvale-Seaford Wetlands Ramsar Site. The bilateral agreements are:

- *Japan-Australia Migratory Bird Agreement (JAMBA)* – The agreement between the Government of Australia and the Government of Japan for the Protection of Migratory Birds in Danger of Extinction and their Environment, 1974;

- *China-Australia Migratory Bird Agreement (CAMBA)* - The Agreement between the Government of Australia and the Government of the People's Republic of China for the Protection of Migratory Birds and their Environment 1986;
- *Republic of Korea-Australia Migratory Bird Agreement (ROKAMBA)* - The Agreement between the Government of Australia and the Republic of Korea for the Protection of Migratory Birds and their Environment, 2006; and
- *The Bonn Convention on Migratory Species (CMS)* - The Bonn Convention adopts a framework in which countries with jurisdiction over any part of the range of a particular species co-operate to prevent migratory species becoming endangered. For Australian purposes, many of the species are migratory birds.

3.1.3 East Asian-Australasian Flyway Partnership

The East Asian-Australasian Flyway Partnership (EAAFP) was launched in November 2006 and is an informal and voluntary initiative that aims to protect migratory waterbirds, their habitat and the livelihoods of people dependent upon them. The partnership covers the East Asia-Australasia flyway which covers 22 countries and extends from the Arctic Circle, through East and South-east Asia, to Australia and New Zealand.

Partners include governments, site managers, academic institutions, UN agencies, development agencies, industrial and private sector, non-government organisations and local people. The partnership provides a platform for dialogue and cooperation, supports the listing and maintenance of internationally recognised wetlands and supports a range of community education programs and activities.




Figure 5 The Edithvale-Seaford Wetlands Ramsar Site Management Plan in context of other requirements for the management of Ramsar sites (adapted from DEWHA 2008).

- 1 General principles
 - 1.01 The primary purpose of management of a declared Ramsar wetland must be, in accordance with the Ramsar Convention :
 - (a) to describe and maintain the ecological character of the wetland; and
 - (b) to formulate and implement planning that promotes:
 - (i) conservation of the wetland; and
 - (ii) wise and sustainable use of the wetland for the benefit of humanity in a way that is compatible with maintenance of the natural properties of the ecosystem.
 - 1.02 Wetland management should provide for public consultation on decisions and actions that may have a significant impact on the wetland.
 - 1.03 Wetland management should make special provision, if appropriate, for the involvement of people who:
 - (a) have a particular interest in the wetland; and
 - (b) may be affected by the management of the wetland.
 - 1.04 Wetland management should provide for continuing community and technical input.
- 2 Management planning
 - 2.01 At least one management plan should be prepared for each declared Ramsar wetland.
 - 2.02 A management plan for a declared Ramsar wetland should:
 - (a) describe its ecological character; and
 - (b) state the characteristics that make it a wetland of international importance under the Ramsar Convention; and
 - (c) state what must be done to maintain its ecological character; and
 - (d) promote its conservation and sustainable use for the benefit of humanity in a way that is compatible with maintenance of the natural properties of the ecosystem; and
 - (e) state mechanisms to deal with the impacts of actions that individually or cumulatively endanger its ecological character, including risks arising from:
 - (i) physical loss, modification or encroachment on the wetland; or
 - (ii) loss of biodiversity; or
 - (iii) pollution and nutrient input; or
 - (iv) changes to water regimes; or
 - (v) utilisation of resources; or
 - (vi) introduction of invasive species; and
 - (f) state whether the wetland needs restoration or rehabilitation; and
 - (g) if restoration or rehabilitation is needed--explain how the plan provides for restoration or rehabilitation; and
 - (h) provide for continuing monitoring and reporting on the state of its ecological character; and
 - (i) be based on an integrated catchment management approach; and
 - (j) include adequate processes for public consultation on the elements of the plan; and
 - (k) be reviewed at intervals of not more than 7 years.
- 3 Environmental impact assessment and approval
 - 3.01 This principle applies to the assessment of an action that is likely to have a significant impact on the ecological character of a Ramsar wetland (whether the action is to occur inside the wetland or not).
 - 3.02 Before the action is taken, the likely environmental impact of the action on the wetland's ecological character should be assessed under a statutory environmental impact assessment and approval process.
 - 3.03 The assessment process should:
 - (a) identify any part of the ecological character of the wetland that is likely to be affected by the action; and
 - (b) examine how the ecological character of the wetland might be affected; and
 - (c) provide adequate opportunity for public consultation.
 - 3.04 An action should not be approved if it would be inconsistent with:
 - (a) maintaining the ecological character of the wetland; or
 - (b) providing for the conservation and sustainable use of the wetland.
 - 3.05 Approval of the action should be subject to conditions, if necessary, to ensure that the ecological character of the wetland is maintained.
 - 3.06 The action should be monitored by the authority responsible for giving the approval (or another appropriate authority) and, if necessary, enforcement action should be taken to ensure compliance with the conditions.

Text Box 1. Australian Ramsar management principles.

3.2 National Legislation and Policy

3.2.1 *Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act)*

The EPBC Act regulates actions that will have or are likely to have a significant impact on any matter of national environmental significance, which includes the ecological character of a Ramsar wetland (EPBC Act 1999 s16(1)). An action that will have or is likely to have a significant impact on a Ramsar wetland will require an environmental assessment and approval under the EPBC Act. An 'action' includes a project, a development, an undertaking or an activity or series of activities (<http://www.environment.gov.au/epbc/index.html>).

The EPBC Act establishes a framework for managing Ramsar wetlands, through the Australian Ramsar Management Principles (EPBC Act 1999 s335), which are set out in Schedule 6 of the Environment Protection and Biodiversity Conservation Regulations 2000. These principles are intended to promote national standards of management, planning, environmental impact assessment, community involvement, and monitoring, for all of Australia's Ramsar wetlands in a way that is consistent with Australia's obligations under the Ramsar Convention. Some matters protected under the EPBC Act are not protected under local or state/territory legislation, for example, many migratory birds are not specifically protected under State legislation. Species listed under international treaties JAMBA, CAMBA and CMS have been included in the List of Migratory species under the Act. Threatened species and communities listed under the EPBC Act may also occur, or have habitat in the Ramsar site; some species listed under State legislation as threatened are not listed under the EPBC Act as threatened, usually because they are not threatened at the national (often equivalent to whole-of-population) level. The Regulations also cover matters relevant to the preparation of management plans, environmental assessment of actions that may affect the site, and the community consultation process.

3.2.2 *National Strategy for the Management of Coastal Acid Sulfate Soils 2000*

National Strategy for the Management of Coastal Acid Sulfate Soils 2000 has four objectives:

- Identify and define coastal acid sulfate soils;
- Avoid disturbance of coastal acid sulfate soils;
- Mitigate impacts when disturbance is unavoidable; and
- Rehabilitate disturbed coastal acid sulfate soils and acid drainage.

3.3 Victorian Legislation and Policy

3.3.1 *Environment Protection Act 1970*

This Act establishes the Environment Protection Authority and makes provision for the Authority's powers, duties and functions. These relate to improving the air, land and water environments by managing waters, acid sulfate soils, control of noise and control of pollution. State Environment Protection Policies (SEPPs) are subordinate legislation made under the provisions of the Act. The SEPP (Waters of Victoria) sets water quality objectives to protect the beneficial uses of waters and covers wetlands in a general nature. Both Edithvale Wetlands and Seafood Wetlands are subject to a Land Subject to Inundation Overlay (LSIO), of which one of its purposes is to protect water quality in

accordance with the provisions of relevant state environment protection policies, particularly in accordance with Clauses 33 and 35 of the SEPP (Waters of Victoria or WoV) (KBR 2009).

Both wetlands are potentially subject to acid sulfate soils. Policy has been established in Victoria to prevent and regulate pollution from acid sulfate soils. These include: the State Environment Protection Policy (Prevention and management of contamination of land); the Industrial Waste Management Policy (Waste acid sulfate soils 1999) which deal with onsite and offset (e.g. disposal) impacts; and the Best Practice Management Guidelines for Dredging which provide broad policy principles for what needs to be considered and a framework for assessing applications for dredging and specifically refer to coastal acid sulfate soils.

Policies emphasise the need to manage land in a manner that avoids adverse impacts from acid sulphate soils, including avoiding any off-site impacts.

3.3.2 Environment Effects Act 1978

This Act establishes the processes for assessment of proposed projects (works) that are capable of having a significant effect on the environment. The Act establishes the role of the Minister for Planning to decide whether an Environmental Effects Statement (EES) is required. The roles and responsibilities of the EES process are described in the Ministerial guidelines for Assessment of Environmental Effects under the *Environment Effects Act 1978* (DSE 2006). The guidelines specify the criteria for referring a project to the Minister for decision on the requirement for an EES. These include effects of potential long term change to the ecological character of a Ramsar site and the need to consider coastal acid sulfate soils in assessing proposed developments.

3.3.3 Flora and Fauna Guarantee Act 1988

This Act provides a legislative and administrative framework for the conservation of biodiversity in Victoria. The Act provides for the listing of threatened taxa, communities and potentially threatening processes. It requires the preparation of action statements for listed species, communities and potentially threatening processes and sets out the process for implementing interim conservation orders to protect critical habitats. The Act also seeks to provide programs for community education in the conservation of flora and fauna and to encourage co-operative management of flora and fauna. Ramsar sites provide habitat for many of Victoria's threatened species and support threatened communities. The Act and Victorian Biodiversity Strategy are currently under review.

3.3.4 Water Act 1989

This Act establishes rights and obligations in relation to water resources and provides mechanisms for the allocation of water resources. This includes the consideration of environmental water needs of rivers and wetlands as well as for human uses, such as urban water supply and irrigation.

The most relevant section of the Act for this Plan is s189 - Functions of Authorities:

- to identify and plan for state and local community needs relating to the use and to the economic, social and environmental values of land and waterways ; and
- to develop and to implement effectively schemes for the use, protection and enhancement of land and waterways;

- to maintain the environmental water reserve in accordance with the environmental water reserve objective;
- to improve the environmental values and health of water ecosystems, including their biodiversity, ecological functions, quality of water and other uses that depend on environmental condition;
- to investigate, promote and research any matter related to its functions, powers and duties in relation to waterway management;
- to educate the public about any aspect of waterway management;
- to identify and plan for State and local community needs relating to the application or use of water in the Water Holdings or other exercise of rights in the Water Holdings within the waterway management district of the Authority;
- to apply or use water in the Water Holdings or otherwise exercise rights in the Water Holdings within the waterway management district of the Authority in accordance with any relevant seasonal watering statement.

(see http://www.austlii.edu.au/au/legis/vic/consol_act/wa198983/s189.html)

3.3.5 *Planning and Environment Act 1986*

The *Planning and Environment Act 1986* sets out procedures for preparing and amending the Victoria Planning Provisions and planning schemes, obtaining permits under schemes, settling disputes, enforcing compliance with planning schemes, and other administrative procedures.

One main function of the Act is to define the roles and responsibilities of the Minister, councils, government departments, the community and other stakeholders in the planning system. This is in addition to setting the broad objectives for planning in Victoria and the main rules, procedures and statutory instruments for how the planning system is to work. The Act uses the planning schemes to set out how land may be used and developed.

The State Planning Policy Framework (SPPF), clause 15.08, stresses the need to avoid disturbing acid sulfate soils. The planning system provides further opportunity at the local level to include more accurate and confident mapping of coastal acid sulfate soils risk areas directly into planning schemes and to identify a planning trigger for implementing a coastal acid sulfate soils risk identification process when considering development proposals.

The SPPF 15.08-2 (gazetted December 2008) states that planning for coastal areas should:

- Avoid development in identified coastal hazard areas susceptible to coastal acid sulfate soils; and
- Ensure that development conserves, protects and seeks to enhance biological diversity and ecological values by avoiding the disturbance of coastal acid sulfate soils.

Zones and Overlays

The majority Edithvale Wetlands is zoned Public Use Zone – Service and Utility, with the exception of the south-western corner which is a Public Park and Recreation Zone. The site is covered by:

- an Environmental Significance Overlay to ensure that development is compatible with identified environmental values; and
- a Land Subject to Inundation Overlay to identify land within the 1 in 100 year flood level, protect water quality and river/wetland/floodplain health, and ensure that development maintains free passage and temporary storage of floodwaters to minimise potential flood damage/impacts.

Similarly, the majority of Seaford Wetlands is a Public Use Zone – Service and Utility, with the exception of the dryland area to the north which is a Public Conservation and Resource Zone. Overlays covering the site include:

- an Environmental Significance Overlay – Schedule 1 (Areas of Botanical or Zoological Significance) to ensure that development is compatible with identified environmental values, botanical and zoological values and protected and enhanced in the long term, and populations and communities of native plants and or fauna are protected;
- a Land Subject to Inundation Overlay to identify land within the 1 in 100 year flood level, protect water quality and river/wetland/floodplain health, and ensure that development maintains free passage and temporary storage of floodwaters to minimise potential flood damage/impacts; and
- Wildfire Management Overlay to ensure that the development of land considers bushfire risk (protection of life and assets) and bushfire protection measures are implemented where warranted.

3.3.6 *Catchment and Land Protection Act 1994*

The *Catchment and Land Protection Act 1994* establishes a framework for the integrated management and protection of catchments. It establishes processes to encourage and support community participation in the management of land and water resources and provides for a system of controls on noxious weeds and pest animals. Section 20 of this Act outlines the duty of land owners to take all reasonable steps to avoid causing or contributing to land degradation which causes or may cause damage to the land of another land owner, as well as to conserve soil and protect water resources.

3.3.7 *Crown Land (Reserves) Act 1978*

The *Crown Land (Reserves) Act 1978* provides for the reservation of Crown lands for public purposes, including: watersheds and gathering grounds for water supply purposes; the supply and distribution of water and works including reservoirs, aqueducts, pipe-lines channels and waterways; the protection of the beds or channels and the banks of waterways; and drainage and sewerage works.

Section 13 of the Act provides for regulating Crown lands for: its care, protection and management; the carrying out of works or improvements; the safety of persons on the land; and for issuing permits and licences and entering into agreements in relation to the land.

Section 14 of the Act provides for the appointment of Committees of Management.

While the Act does not make specific reference to coastal acid sulphate soils, there is scope under these provisions for any coastal acid sulphate soils risk identification process to be triggered or for regulations to be made.

3.3.8 Coastal Management Act 1995

Section 37 of the *Coastal Management Act 1995* requires the written consent of the Minister for proposals for use and development on coastal Crown land (Crown land 200 m from high tide). There is opportunity for a coastal acid sulfate soils risk identification process to be triggered under these consent provisions.

3.3.9 Wildlife Act 1975 and Wildlife Regulations 2002

The purposes of the *Wildlife Act 1975 and Wildlife Regulations 2002* are to:

- establish procedures in order to promote:
 - the protection and conservation of wildlife;
 - the prevention of taxa of wildlife from becoming extinct; and
 - the sustainable use of and access to wildlife.
- prohibit and regulate the conduct of persons engaged in activities related to wildlife.

The Act covers all wildlife, including all native vertebrate animals, all kinds of deer, non-indigenous quail, pheasants, and partridges, and all terrestrial invertebrate animals listed under the *Flora and Fauna Guarantee Act 1988*. The Act regulates the hunting, trading and taking of wildlife.

Research or Scientific Permits are issued under the Act (and *Flora and Fauna Guarantee Act 1988*) to carry out:

- surveys, research, management, protection, conservation and enhancement of wildlife populations and wildlife habitats;
- public education programmes for promoting and maintaining an appreciation of the value of wildlife;
- scientific or biological research, studies, or investigations in conjunction with other scientific organizations; and
- removal of wildlife from any particular locality.

Management of Eastern Grey Kangaroos at Edithvale South Wetland is covered under a Wildlife Permit issued to Melbourne Water under the *Wildlife Act 1975*. Wildlife licences are also issued under this Act to take or destroy wildlife, which would include removal of introduced species from the wetlands.

3.3.10 Country Fire Authority Act 1958

Fire prevention responsibilities are derived from Section 43 of the *Country Fire Authority Act 1958* which delegates fire suppression power and duties to public authorities (see Terramatrix 2013a):

- “in the country area of Victoria it is the duty of every municipal council and public authority to take all practicable steps (including burning) to prevent the occurrence of fires on, and minimise the danger of the spread of fires on and from (a) any land vested in it or under its control or management, and (b) any road under its care and management”.

Section 96A of this Act requires municipal councils to appoint a person to be the fire prevention officer for that council for the purposes of this Act.

Under Sections 20 and 46 of the Act, respectively, and regarding the country area of Victoria (CFA response area), it is the Country Fire Authority's (CFA's) role to superintend and enforce fire prevention and to report any failure by a public authority or municipal council to properly carry out their duties.

Under Sections 54 and 55 of the Act, respectively, the CFA may appoint a Municipal Fire Prevention Committee (MFPC) to undertake a range of functions, including advising Council on the existence and management of hazards, and making recommendations in the preparation of the Municipal Fire Prevention Plan. The key fire management responsibilities for the municipality rest with the (Frankston) Municipal Fire Management Planning Committee (MFMP), which is a sub-committee of the Municipal Seaford Wetlands – Fire Management Plan (see Terramatrix 2013a).

3.3.11 Victorian Waterway Management Strategy

The 2013 Victorian Waterway Management Strategy (VWMS) provides the framework for government—in partnership with the community—to maintain or improve the condition of rivers, estuaries and wetlands so that they can continue to provide environmental, social, cultural and economic values for all Victorians. The framework is based on regional planning processes and decision-making, within the broader system of integrated catchment management in Victoria.

3.3.12 Victorian Coastal Strategy 2014

Achievements of previous iterations of the Victorian Coastal Strategy (Victorian Coastal Council 2014) have included the establishment of a hierarchy of principles that guides effective coastal planning and decision-making in accordance with the objectives of the *Coastal Management Act 1995*. The first principle of this Strategy is to protect significant environmental and cultural values, based upon a sound understanding of coastal features, vulnerability and risks. This Strategy also emphasises the need to avoid disturbing coastal acid sulfate soils and ensure any development proposed near or on coastal acid sulfate soils demonstrates that it will avoid any disturbance.

3.3.13 Victorian Coastal Acid Sulfate Soils Strategy

The Victorian Coastal Acid Sulfate Soils Strategy (DSE 2009c) aims to protect the environment, humans and infrastructure from the harmful effects of disturbing coastal acid sulfate soils. It aims to build the capacity of land owners and land and water managers to plan and manage activities in coastal acid sulfate soils risk areas. It identifies principles and promotes a risk management approach to guide decision-making. The emphasis of the strategy is on avoiding disturbing coastal acid sulfate soils.

3.3.14 Public Health and Wellbeing Act 2008

The *Public Health and Wellbeing Act 2008* received Royal Assent in September 2008 and took effect on 1 January 2010. The Act replaces the previous Victorian *Health Act 1958* and is a key piece of legislation designed to protect the health of Victoria's population, with implications for:

- authorised officers within local councils and the Department of Health and Human Services;
- pest control operators;
- cooling tower operators;
- the governance and management of a range of consultative councils established under the Act;

- the management of infectious diseases, micro-organisms and medical conditions by medical and health practitioners, Victoria's Chief Health Officer and affected individuals; and
- the development of public health policy through providing for municipal public health and wellbeing plans, a State public health and wellbeing plan and in some circumstances, health impact assessments.

The *Public Health and Wellbeing Regulations 2009* were made by the Governor in Council on 15 December 2009 and took effect on 1 January 2010. The regulations replace the previous *Health (Infectious Diseases) Regulations 2001* made under the former *Health Act 1958* which empowered the Victorian Health Department and councils as their agents to direct owners and occupiers of land to undertake measures to reduce the incidence of mosquito breeding to prevent mosquito-borne diseases.

The *Public Health and Wellbeing Regulations 2009 S.R. No. 178/2009 (1)* states for the purposes of Section 235(a) of the Act, in order to prevent breeding of mosquitoes which may be vectors of arboviruses, an authorised officer may give written directions to the owner or occupier of any premises to remove, or take steps to avoid, any condition on those premises conducive to the breeding of mosquitoes. Thus, Melbourne Water has the responsibility to manage mosquitoes within the assets that it owns and maintain as a matter of public health as consistent with other managers of water assets in Victoria (also see DSE 2004; GHD 2015; Melbourne Water 2016, undated^a and undated^b).

3.3.15 Framework for Mosquito Management in Victoria (DSE 2004)

The objective of the Framework for Mosquito Management in Victoria (DSE 2004) is to provide a strategic framework to assist councils and State government agencies to implement local mosquito management programs in an effective and environmentally appropriate manner consistent with Victoria's commitments to sustaining and improving community health and protecting the environment. The Framework provides a legislative framework and discusses the potentially conflicting demands of legislation designed to protect human health (i.e. *Public Health and Wellbeing Regulations 2009*) via mosquito control and legislation designed to protect the environment, species and ecological processes that occur naturally, such as breeding sites for mosquitoes being located in high value wetlands (e.g. Ramsar sites and EPBC Act 1999), the role of mosquitoes in the food chain and identifying methods of mosquito control potentially impacting unintentionally and negatively on the environment (e.g. physical, chemical and biological control). The Framework also addresses the potential disturbance to cultural values (both indigenous and post-settlement heritage) associated with mosquito management (e.g. construction of drainage channels and changes to natural water flow) and identifies the relevant legislation which aims to protect cultural values.

The Framework also:

- identifies responsibilities of State Government, councils and community in mosquito management;
- provides a risk assessment framework to health and well-being based on the likelihood of being bitten by arbovirus-carrying or nuisance mosquitoes and the health and social impacts of being bitten (criteria for health risk are based on past outbreaks of diseases - the level of risk from an incidence or high potential of MVE is very high, the level of risk from a high incidence of RRV or BFV is high, the level of risk from an abundance of nuisance mosquitoes and some RRV or BFV in the past is moderate, the level of risk from a low density of nuisance mosquitoes is low); and

- gives a risk assessment framework to the environment from mosquito management intervention based on the environmental sensitivity of the site (criteria for environmental sensitivity are based on land status – Ramsar sites, sites supporting important habitat for nationally threatened species or migratory species have high environmental sensitivity, National, State or Coastal Parks have high sensitivity, etc.) and the environmental risk associated with mosquito management based on the Ramsar Convention Wetland Risk Assessment; and
- provides a framework for mosquito management incorporating an integrated and co-ordinated approach with various methods and delivery of management varying depending on environmental sensitivity of the site and the area to be covered.

Based on the Framework for Mosquito Management in Victoria (DSE 2004), Melbourne Water (undated^a) has prepared the Mosquito Management Decision Support Tool to guide mosquito management at Melbourne Water assets.

3.4 Regional and Local Plans and Policy

3.4.1 Port Phillip and Western Port Regional Catchment Strategy

The Port Phillip and Western Port Regional Catchment Strategy is a statutory document under the CaLP Act that provides the overarching framework for land, water and biodiversity management in the region.

3.4.2 Better Bays and Waterways

Better Bays and Waterways was developed by the Environment Protection Authority Victoria and Melbourne Water to achieve water quality improvement for the Port Phillip and Western Port region. The plan was aimed at reducing the amount of pollutants entering waterways and bays from rural, urban and coastal areas.

3.4.3 Healthy Waterways Strategy

The *Healthy Waterways Strategy* (Melbourne Water 2013a) outlines Melbourne Water's role in managing rivers, estuaries and wetlands in the Port Phillip and Westernport region. This strategy focuses on investing in areas that the community values and that will protect and improve environmental values and increase liveability.

Melbourne Water's (2013a) *Healthy Waterways Strategy 2013* recognises that Melbourne Water land supports important environment and social values and states that waterways:

- connect diverse and thriving communities of native plants and animals;
- provide amenity to urban and rural areas and engage communities with their environment; and
- are managed sustainably to balance environmental, economic and social values.

Sites on Melbourne Water land that support important biodiversity values may be considered a Site of Biodiversity Significance (SoBS). Under the *Waterways Sites of Biodiversity Significance Plan 2013* (Melbourne Water 2013b), Melbourne Water recognises 36 SoBS all of which are managed by Melbourne Water and support at least one of the following biodiversity values:

1. species or ecological communities protected under the Commonwealth EPBC Act;

2. species or ecological communities listed as threatened under the Victorian FFG Act;
3. Ecological Vegetation Class or Classes (EVCs) which are considered to be vulnerable or endangered within the bioregion;
4. species that are listed under the Advisory List of Rare or Threatened Plants in Victoria – 2014 (DEPI 2014), Advisory List of Threatened Vertebrate Fauna in Victoria – 2013, Advisory List of Threatened Invertebrate Fauna in Victoria – 2009; and
5. connecting habitat for threatened flora and fauna.

In addition to providing the listing criteria for SoBS, the *Waterways Sites of Biodiversity Significance Plan 2013* (Melbourne Water 2013b) gives directions on how SoBS should be managed in accordance with Melbourne Water's obligations under environmental legislation, and to achieve the objectives of the Healthy Waterways Strategy 2013.

Edithvale-Seaford Wetlands were identified as high value SoBS because they fulfilled a number of criteria, including the sites: are greater than 50 ha in size; support substantial vegetation remnants with several threatened Ecological Vegetation Classes; support large populations or representatives of several EPBC Act and FFG Act-listed items (e.g. Australasian Bittern; Ramsar-listed); support other threatened flora and fauna (e.g. VROTs, Advisory Listed fauna); support significant populations of Migratory and/or Marine Overfly bird species (e.g. migratory waders); and demonstrate connectivity for threatened species (Ecology Australia 2013).

4 Ecological Character and Values - Critical Components, Processes and Services

The Australian Government has developed and implemented a framework for describing the ecological character of Ramsar sites (DEWHA 2008). This framework requires the identification and description of critical components, processes and services. These are defined as characteristics of the Ramsar site (DEWHA 2008):

1. that are important determinants of the sites unique character;
2. that are important for supporting the Ramsar criteria under which the site was listed;
3. for which change is reasonably likely to occur over short to medium time scales (less than 100 years); and/or
4. that will cause significant negative consequences if change occurs.

The ECD for the Edithvale-Seafood Ramsar Site (2002) was drafted prior to the release of these national guidelines and so does not describe character in terms of critical components, processes and services. A recent review has developed an addendum to the ECD (Hale 2016) and identified four components, processes and services that are critical to the ecological character of the Ramsar site:

- Waterbird diversity and abundance;
- Waterbird breeding;
- Physical habitat for waterbirds; and
- Threatened wetland species.

4.1 Waterbird Diversity and Abundance

The Edithvale-Seafood Wetlands support a diversity and abundance of waterbirds, with average total annual maximum counts of around 5000. The site supports a diversity of birds and over 75 waterbird species have been recorded at the site (KBR 2009; DSE 2012). Twenty species of waterbirds listed under international migratory agreements have been recorded within the Ramsar site. This number includes species that, in Australia, are residents (e.g. Eastern Great Egret) and a number of migratory species that are only occasionally recorded at the site. There are eight species of international migratory shorebirds that are regularly supported (two thirds of seasons) by the Edithvale-Seafood Wetlands Ramsar site (Table 2).

The site supports a variety of different types of waterbirds, which is illustrated by looking at the functional guilds (Figure 6). In particular, the site is important for waders, which are the most numerous birds at the site. The site, however, also supports numbers of ducks, fish eating species (such as pelicans and cormorants), herbivores (such as swans and coots) and large wading birds (such as ibis and herons).

Table 2 Palaeartic migratory waders recorded in the Ramsar site and their frequency of occurrence (percentage). The eight species that the site is considered to regularly support are shaded.

Common name	Species name	JAMBA	CAMBA	ROKAMBA	Percent of years
Australian Painted Snipe	<i>Rostratula australis</i>		X		5
Common Greenshank	<i>Tringa nebularia</i>	X	X	X	86
Curlew Sandpiper	<i>Calidris ferruginea</i>	X	X	X	68
Latham's Snipe	<i>Gallinago hardwickii</i>	X	X	X	100
Long-toed Stint	<i>Calidris subminuta</i>	X	X	X	14
Marsh Sandpiper	<i>Tringa stagnatilis</i>	X	X	X	68
Pectoral Sandpiper	<i>Calidris melanotos</i>	X	X	X	68
Red Knot	<i>Calidris canutus</i>	X	X	X	5
Red-necked Stint	<i>Calidris ruficollis</i>	X	X	X	73
Sharp-tailed Sandpiper	<i>Calidris acuminata</i>	X	X	X	100
Wood Sandpiper	<i>Tringa glareola</i>	X	X	X	68




Figure 6 Maximum annual abundance of waterbird guilds at Edithvale-Seaford Wetlands 1994 to 2015 (data from BirdLife Australia). Guild membership as defined by Kingsford et al. (2012).

4.2 Waterbird Breeding

There are records of over 20 species of waterbird breeding within the Ramsar site (RAOU 2001; Birds Australia 2006, 2007, 2008, 2010, 2012; BirdLife Australia 2013a, 2013b, 2015b and 2016a). The most common breeding species are Black Swan *Cygnus atratus* and a range of duck species, such as Chestnut Teal *Anas castanea* and Blue-billed Duck *Oxyura australis* (FFG Act-listed, 10 breeding records at Edithvale Wetlands since 1998 and they occasionally breed at Seaford). There are also records of wetland dependent raptors (Swamp Harrier *Circus approximans*) and other wetland dependent birds (e.g. Australian Reed-Warbler *Acrocephalus australis*) breeding at the site (e.g. BirdLife Australia 2016a).

4.3 Physical Habitat for Waterbirds

Hydrology and vegetation type have been identified as the most important habitat components for supporting waterbirds at the Ramsar site (Birds Australia 2004). The mosaic nature of the habitat is what supports the broad range of species. The wetlands have been divided into habitat zones and two zones in each portion of the Ramsar site are considered most important for waterbirds (Birds Australia 2004):

- Edithvale (Figure 12):
 - Edithvale North (EN1) - deeper water for a number of duck species, surrounded by tall reeds; and
 - Edithvale South (ES1) - shallow wetlands that seasonally dry providing foraging habitat for shorebirds, grading to tall marsh at the fringes, providing cover for species, such as Australasian Bittern and Latham's Snipe.
- Seaford (Figure 13):
 - Downs Estate (Cell SN) - pasture which provides foraging habitat for large wading species, such as ibis; and
 - Seaford North 2 Pool, Seaford Central West 1 (SCW1) and Seaford East 2 - mosaic of deeper water, tall marsh, deeper saline ponds important for all wetland bird species.

4.4 Threatened Wetland Species

The Edithvale-Seaford Wetlands regularly support two waterbird species listed as threatened species under the Commonwealth EPBC Act 1999.

Curlew Sandpiper *Calidris ferruginea* is a small bird weighing just 60 grams (Higgins and Davies 1996). It is an international migratory species that spends the non-breeding season in the southern hemisphere. They arrive in late spring, spend the summer feeding on invertebrates in intertidal mudflats and depart for the northern hemisphere in February to March. Juveniles who arrive in the Ramsar site spend their first one or two winters at the site before heading to the northern hemisphere to breed.

Curlew Sandpiper is listed as Critically Endangered under the EPBC Act due to declines in their global populations. There have been a large number of investigations into the decline of shorebirds in the East Asian-Australasian Flyway, with habitat declines particularly at staging areas in the Yellow Sea

recognized as the most significant impact factors (MacKinnon et al. 2012, Murray et al. 2015, Hua et al. 2015).

Australasian bittern *Botaurus poiciloptilus* is a shy and cryptic wading species of waterbird. Habitat preferences are for permanent, densely vegetated freshwater wetlands (Menkhorst 2012). It forages mainly at night in shallow water up to 30 cm deep and feeds on frogs, fish and invertebrates as well as occasional plant material (Marchant and Higgins 1990). Within the Ramsar site the species inhabits emergent vegetation, but there is a balance between having sufficient cover and the vegetation being too dense for effective foraging. Control of emergent vegetation such as Common Reed *Phragmites australis* is important for this species at the site. The bitterns require reed beds and edges, but they benefit from mosaic reed control, as the re-growth Tall Marsh is less dense and therefore more suitable for foraging and habitat (Andrew Silcocks, BirdLife Australia, pers. comm.; Paul Rees, Melbourne Water, pers. comm.).

The site provides important winter feeding habitat for the species, which travels north to breed. Within the site, there have been declines in the species (Figure 7), but this may reflect more widespread population changes than characteristics of the Ramsar site (Andrew Silcocks, BirdLife Australia, pers. comm.).




Figure 7 Maximum annual abundance of Australasian Bitterns at Edithvale-Seaford Wetlands 1994 to 2015 (data from BirdLife Australia).

4.5 Flora

4.5.1 Vegetation

The large Carrum Carrum Swamp that formerly encompassed both Edithvale and Seaford Wetlands was dramatically altered when the Patterson Cut was constructed in the 1870's to drain the Swamp. Edithvale and Seaford Wetlands both occur in what would have been the deeper parts of the former Carrum Carrum Swamp. The drainage of the wetland irreversibly changed the hydrology and hence the

vegetation communities present within these two sites, whereby the current vegetation now more closely resembles what would have surrounded the edges of the Carrum Carrum Swamp prior to drainage.

Eight Ecological Vegetation Classes (EVCs) were identified at Edithvale and Seaford Wetlands in Australian Ecosystems' latest survey (Australian Ecosystems 2015): Brackish Wetland Aggregate, Brackish Aquatic Herbland, Brackish Herbland, Aquatic Herbland, Plains Sedge Wetland, Tall Marsh, Swamp Scrub and Damp Sands Herb-rich Woodland. The distribution and occurrence of these EVCs as mapped by Australian Ecosystems (2011a, 2015; TBLD and Australian Ecosystems 2005) is in a constant state of flux primarily responding to seasons, water levels and salinity. These vegetation communities and distributional changes are discussed below based on field observations and previous reports (TBLD and Australian Ecosystems 2005; Australian Ecosystems 2011a, 2015). Maps illustrating the distribution of EVCs at each site are provided in Figures 8 and 9.

Note: an asterisk (*) denotes exotic species while a hash (#) denotes native species that are not considered indigenous at the site.

Damp Sands Herb-rich Woodland (EVC 3) – Vulnerable in the Gippsland Plain Bioregion

According to Australian Ecosystems (2015), Damp Sands Herb-rich Woodland occurred at Seaford Wetlands in areas that had primarily been revegetated, or previously identified as Plains Grassy Woodland (Australian Ecosystems 2011a). This vegetation community occurs on the quaternary dunes that surrounded the Carrum Carrum Swamp. The dominant canopy species observed comprised River Red-gum *Eucalyptus camaldulensis* and Coast Manna Gum *Eucalyptus viminalis* subsp. *pryoriana*, along with Black Wattle *Acacia mearnsii* and Coast Banksia *Banksia integrifolia* in the sub-canopy. Understorey shrubs were sometimes included in the revegetation and some areas have been heavily invaded by Coast Tea-tree #*Leptospermum laevigatum*, a large shrub species that is only considered indigenous to the coastal dunes and headlands (Thompson 1989). The observed understorey was generally dominated by exotic vegetation, primarily mown or unmown grasses, with some areas of Bracken *Pteridium esculentum* or revegetated indigenous species.

Swamp Scrub (EVC 53) – Endangered in the Gippsland Plain Bioregion

Swamp Scrub was identified in small disjunct patches across Seaford Wetlands (including an artificial island) and three small islands in the north of Edithvale Wetlands. This EVC had previously been identified as Estuarine Scrub in 2010 (Australia Ecosystems 2011a) based on the understorey composition of species more tolerant of salt that were uncommon during the current field surveys. Swamp Paperbark *Melaleuca ericifolia* dominated this EVC which is situated in slightly more elevated areas than the aquatic wetland and marsh EVCs. Scattered Swamp Gums *Eucalyptus ovata* were also occasionally present. Understorey species included Water Ribbons *Cycnogeton procerum* and Common Reed *Phragmites australis* in wetter areas and Blackberry **Rubus fruticosus* spp. agg. and exotic grasses such as Paspalum **Paspalum dilatatum*, Kikuyu **Cenchrus clandestinum*, Couch **Cynodon dactylon* var. *dactylon* in drier areas. In some places the understorey has been revegetated to introduce diversity.

Tall Marsh (EVC 821) – not assigned a bioregional conservation status for the Gippsland Plain

Tall Marsh vegetation was dominated by thick swards of Common Reed *Phragmites australis* and occasionally Cumbungi (*Typha domingensis* and/or *Typha orientalis*). Other plant species, such as Water

Ribbons *Cyanogeton procerum*, Salt Club-sedge *Bolboschoenus caldwellii*, Common Duckweed *Lemna disperma*, Common Blown-grass *Lachnagrostis filiformis*, Hastate Orache *Atriplex prostrata*, Aster-weed *Symphotrichum subulatum*, Spiny Rush *Juncus acutus*, Couch, Kikuyu and Toowoomba Canary-grass *Phalaris aquatica*, were present though uncommon amongst the Common Reed and Cumbungi.

In many locations at the wetlands, this EVC has been expanding into the other wetland EVCs, which reduces floristic diversity and changes the structure of the vegetation from an open wetland to a dense, tall sward (sometimes to 3.5 m tall) of Common Reed or Cumbungi. Given the scarcity of the more open and diverse wetland communities, and the reduced fauna habitat suitability of the Tall Marsh, grooming of Common Reed has been trialled to limit the expansion of this EVC (Plate 1). The results have shown to be quite successful, as discussed further in Section 9.4.1.



Plate 1 Ungroomed (left) and groomed (right) Common Reed *Phragmites australis* at Edithvale North Wetland.

Brackish Wetland Aggregate (EVC 656) – Endangered in the Gippsland Plain Bioregion

The Brackish Wetland was variously dominated by Sea Rush *Juncus kraussii* subsp. *australiensis*, Salt Club-sedge *Bolboschoenus caldwellii*, Fine Twig-sedge *Baumea arthropphylla*, Common Spike-sedge *Eleocharis acuta*, Shiny Swamp-mat *Selliera radicans* and Creeping Monkey-flower *Mimulus repens* among others. Some areas have also been invaded by Spiny Rush *Juncus acutus* subsp. *acutus*. It is common at both wetlands, however since 2010, large areas of this EVC has been displaced by the encroachment of Common Reed which has resulted in a changed vegetation structure, reduced floristic diversity and reduced fauna habitat diversity. Particularly at Edithvale Wetlands, this EVC commonly appears as Unvegetated open water/bare soil/mud (EVC 990) for around six months of the year.

Brackish Aquatic Herbland (EVC 537) – not assigned a bioregional conservation status for the Gippsland Plain

Brackish Aquatic Herbland occurred in deeper water with aquatic herbs and filamentous algae dominating. Common species included Water Ribbons, Lake Water-milfoil *Myriophyllum salsgineum*,

and Fennel Pond-weed *Potamogeton pectinatus* in the deeper water, through to Slender Knotweed *Persicaria decipiens* and Swamp Crassula *Crassula helmsii* in the drier margins.

This EVC has contracted in extent at both Edithvale and Seaford Wetlands over the past decade where it was once abundant in both the northern and southern Edithvale Wetlands, and scattered in the south of Seaford Wetlands (TBLD and Australian Ecosystems 2005). This EVC now only occurs in the constructed northern wetlands at Edithvale (Australian Ecosystems 2011a, 2015) and is no longer represented at Seaford (seeming due to encroaching Tall Marsh).

Plains Sedgy Wetland (EVC 647) – not assigned a bioregional conservation status for the Gippsland Plain

Plains Sedgy Wetland occurs at both sites. The occurrence at Seaford was restricted to the perimeter of two small Aquatic Herbland dominated waterbodies in the south east of the site that are separated from the main Brackish Wetland dominated waterbody by an artificial levee. At Edithvale, it occurred in the lower lying areas of the large wetland just north of Edithvale Road. Plains Sedgy Wetland was dominated by Common Spike-sedge *Eleocharis acuta* and Fine Twig-sedge, Water Ribbons, Common Blown-grass *Lachnagrostis filiformis* and a variety of amphibious forbs. Common Reed and Cumbungi surround or are interspersed through all patches identified and their encroachment threaten the floristic diversity.

Aquatic Herbland (EVC 653) – not assigned a bioregional conservation status for the Gippsland Plain

Small discrete pockets of Aquatic Herbland exist at Seaford Wetlands, while at Edithvale the areas of Aquatic Herbland are replaced with Tall Marsh and Brackish Wetland Aggregate/Unvegetated (open water/bare soil/mud) as the water draws down in spring. Dominant species commonly included Water Ribbons, Water-milfoil *Myriophyllum* spp., Small River Buttercup *Ranunculus amphitrichus*, Streaked Arrowgrass *Triglochin striata*, Common Duckweed, Tall Spike-sedge *Eleocharis sphacelata*, and to a lesser extent Common Reed (Greet 2015; Australian Ecosystems 2011b).

Brackish Herbland (EVC 538) – not assigned a bioregional conservation status for the Gippsland Plain

Brackish Herbland occurred in two discrete areas on the outskirts of the Edithvale South wetland. It had previously also been identified as occurring in the middle of this wetland (Australian Ecosystems 2016), though was not identified there in the current survey. Instead, this area was dominated by open water, mudflat or Common Reed (resembling Brackish Wetland Aggregate), with occasional herbs tolerant of brackish conditions such as Beaded Glasswort *Sarcocornia quinqueflora* subsp. *quinqueflora* and Creeping Monkey-flower *Mimulus repens*. As per Aquatic Herbland, it is likely that this EVC will continue to fluctuate in distribution depending on the season and annual rainfall each year.

Wet Saltmarsh Herbland (Coastal Saltmarsh Aggregate, EVC 9) – Least Concern in the Gippsland Plain Bioregion, Vulnerable under the EPBC Act 1999

A small area of vegetation resembling Wet Saltmarsh Herbland has established in Seaford Wetland where the Tall Marsh has died back (possibly as a result of high salinity; Paul Rees, Melbourne Water, pers. comm.). This area was dominated by Beaded Glasswort *Sarcocornia quinqueflora* subsp. *quinqueflora* with a small scattering of other typical saltmarsh species such as Shiny Swamp-mat *Selliera radicans* and Creeping Monkey-flower *Mimulus repens*. Spiny Rush *Juncus acutus* subsp. *acutus* was present, though most plants had been killed as a result of management. The small extent of this EVC

within the site disqualifies it as part of the EPBC Act listed community *Subtropical and Temperate Coastal Saltmarsh*.

4.5.2 Significant Flora Species

Seven Victorian rare or threatened (VROT) plant species have been identified as having a moderate likelihood of occurrence within Edithvale Wetlands, and five VROT species have been identified for Seaford Wetland. Further information on this can be found in Appendix 2.



Figure 8 Ecological Vegetation Class mapping for Edithvale Wetlands.




Figure 9 Ecological Vegetation Class mapping for Seaford Wetland.

4.6 Other Fauna

4.6.1 Threatened Bird Species

Monthly surveys undertaken since 1989 at Edithvale Wetlands have recorded totals of:

- 157 species at Edithvale North wetlands (including 11 exotic species, and the Cockatiel *Nymphicus hollandicus* is probably an aviary escapee);
- 166 species at Edithvale South wetlands (including 14 exotic species); and
- 177 species over both wetlands (BirdLife Australia 2016a; BirdLife Atlas database; Appendix 3).

Monthly surveys undertaken since 1994 at Seaford Wetlands have recorded a total of 174 species (including 13 exotic species, and the Cockatiel *Nymphicus hollandicus* and Scaly-breasted Lorikeet *Trichoglossus chlorolepidotus*, both probably represent aviary escapees and the Pacific Black Duck – Mallard hybrid *Anas superciliosa* - *Anas platyrhynchos*) (BirdLife Australia 2016a; BirdLife Atlas database; Appendix 3).

Overall, 197 species have been recorded at Edithvale-Seaford Wetlands (Appendices 3 and 4).

A number of threatened or migratory species have been recorded at the wetlands (Appendices 3 and 4) and the totals are given in Table 3.

Table 3 Numbers of species recorded at Edithvale and Seaford Wetlands and listed as threatened or migratory/marine-overfly species under the Commonwealth Environment Protection and Biodiversity Act 1999 (EPBC Act) and/ or threatened under the Victorian Flora and Fauna Guarantee Act 1988 (FFG Act) and/or classified as threatened in Victoria by DSE (2013).

Wetlands	EPBC Act-listed threatened species	EPBC Act Migratory or Marine-overfly species	FFG Act listed threatened species	DSE (2013) classified threatened species
Edithvale North	1	48	10	28
Edithvale South	2	59	13	30
Seaford	2	58	12	28
Edithvale and Seaford combined	4	68	14	34

The wetlands support a high number of threatened bird species, including (from ANCA 1996; Ecology Australia 2000; RAOU 2001; Birds Australia 2004, 2006, 2007, 2008, 2010 and 2011; BirdLife Australia 2013a, 2013b, 2015a, 2015b and 2016a; KBR 2009; SKM 2011; Andrew Silcocks, Birds Australia, pers. comm.):

- populations of diving duck species which are threatened in Victoria, such as the Blue-billed Duck *Oxyura australis* (FFG and endangered under DSE 2013) and Hardhead *Aythya australis*

(vulnerable under DSE 2013) which are known to breed at the wetlands, and Musk Duck *Biziura lobata* (vulnerable under DSE 2013), which favour areas of deeper water;

- Australasian Shoveler *Anas rhynchos* (vulnerable under DSE 2013), a dabbling duck which strains insects and seeds from water (e.g. in the marshy areas);
- Little Bittern *Ixobrychus minutus dubius* (FFG and endangered under DSE 2013) is a summer visitor to Victoria from northern Australia which bred at Edithvale Wetlands over several seasons;
- populations of crakes and rails occur regularly at ESW, including Lewin's Rail *Rallus pectoralis* and Baillon's Crake *Porzana pusilla palustris* (both FFG and vulnerable under DSE 2013) which prefer dense vegetation (e.g. reed beds) for shelter and nesting and nearby mudflat edges for foraging;
- non-breeding individuals of FFG Act-listed Eastern Great Egret *Ardea modesta*, Little Egret *Egretta garzetta nigripes* and Intermediate Egret *Ardea intermedia*;
- A suite of other threatened species have been recorded less regularly at ESW, such as the Australian Painted Snipe *Rostratula australis* (EPBC and FFG), Freckled Duck *Stictinetta naevosa* (FFG and endangered under DSE 2013), Caspian Tern *Sterna caspia* (FFG and near threatened under DSE 2013) and Wood Sandpiper *Tringa glareola* (vulnerable under DSE 2013).
- Latham's Snipe *Gallinago hardwickii* is a medium-sized wader which breeds on Hokkaido in northern Japan and far eastern Russian Islands in the Japanese Archipelago during the boreal summer, and migrates to south-eastern Australia where it spends the austral summer. It arrives in Victoria from late August and leaves by late March or early April (Higgins and Davies 1996). Latham's Snipe is listed as a migratory species under the EPBC Act and is classified as near threatened in Victoria (DSE 2013). It used to occur in large numbers of up to 400 snipe at Seaford Wetlands in the 1970's and 1980's (Sean Dooley, BirdLife Australia, pers. comm.; ANCA 1996), but has more recently been recorded in smaller numbers during regular counts (up to 20–30 snipe, Table 4). DEWHA (2009) developed the EPBC Act Policy Statement 3.21 outlining the Significant Impact Assessment Guidelines for 36 Migratory Shorebird species. DEWHA (2009), with advice from experts, define 'important habitat' for Latham's Snipe as those sites that support at least 18 individuals of the species. Edithvale Wetlands have exceeded/equalled the 18 Snipe threshold during 17 months of the monthly surveys undertaken since 1989 (Table 4). Seaford Wetlands have exceeded/equalled 18 Snipe during six months since 1994. Together, the Edithvale-Seaford Wetlands Ramsar Site has exceeded the threshold on 21 occasions. Thus, Edithvale and Seaford Wetlands supports 'important habitat' for Latham's Snipe as defined under the EPBC Act.
- Pied Cormorant *Phalacrocorax varius* and Royal Spoonbill *Platalea regia* (near threatened), generally forage in small numbers at the wetlands (a pair of Royal Spoonbills bred in the southern section of Seaford Wetlands over the 2010/2011 season; this was the first and only breeding record for this species at the Ramsar site).
- For the terrestrial species, there are also records of Swift Parrot *Lathamus discolor* (EPBC and FFG) from Seaford Wetlands, the latest record being from March 2015 (Appendices 3 and 4).

- The Orange-bellied Parrot *Neophema chrysogaster* (EPBC and FFG) was last recorded in the area in 1980's, at Seaford Swamp in 1986 and 2 km south of Edithvale and at Wells Road in 1988 (VBA, DELWP 2016).
- A Powerful Owl *Ninox strenua* (FFG and vulnerable under DSE 2013) was recorded at Edithvale South Wetland over a two week period in November 2015 (Paul Rees, Melbourne Water, pers. comm.).
- Marshland Passerines (e.g. Australian Reed-warbler *Acrocephalus australis*, Little Grassbird *Megalurus garmineus*, Golden-headed Cisticola *Cisticola exilis*, White-fronted Chat *Epithanura albifrons*, Striated Fieldwren *Calamanthus fuliginosus*, White-browed Scrubwren *Sericornis frontalis*, Superb Fairy-wren *Malurus cyaneus* and Brown Thornbill *Acanthiza pusilla*) represent an important component of the margins of wetlands that support tall reedbeds (BirdLife Australia 2015b). Higher numbers occur at the wetlands during wet years, suggesting that lower invertebrate prey supplies during droughts force marshland passerines to seek wetter environments (Andrew Silcocks, BirdLife Australia, pers. comm.).
- A diversity of raptor species occur at the wetlands. Those commonly recorded include the Swamp Harrier *Circus approximans*, Black-shouldered Kite *Elanus axillaris*, Whistling Kite *Haliastur sphenurus*, Brown Goshawk *Accipiter fasciatus*, Peregrine Falcon *Falco peregrinus*, Brown Falcon *Falco berigora*, Australian Hobby *Falco longipennis*, Nankeen Kestrel *Falco cenchroides* and Eastern Barn Owl *Tyto javanica*. Little Eagle *Hieraaetus morphnoides* and White-bellied Sea-Eagle *Haliaeetus leucogaster* (FFG) also occasionally visit the wetlands. The mosaic of open wetlands, Short Marsh, Tall Marsh, Swamp Scrub, Woodland, indigenous and non-indigenous plantings and grassland provide a diversity of habitats for raptor species. Most species have been recorded foraging over all habitats at Edithvale Wetlands. At Seaford Wetlands, the habitats most used are Red Gum Woodland and associated wetlands and adjoining terrestrial areas in the east. The dependency of each species on the wetlands will vary (e.g. two pairs of Swamp Harriers occur and breed in and forage over wetlands at Edithvale and Seaford Wetlands: whereas Australian Hobbies breed in trees but forage over wetlands and both Black-shouldered Kites and Nankeen Kestrels may opportunistically take prey from reedbeds or adjacent habitats).

Table 4 Number of months during which BirdLife Australia surveys at Edithvale Wetlands from September 1989 to September 2015 and Seaford Wetlands from June 1994 to September 2015 have recorded 18 Latham’s Snipe or more (number of snipe recorded are given), Edithvale–Seaford Wetlands.

Month and Year	Edithvale-Seaford (highest count)	Edithvale North	Edithvale South	Seaford
November 1991	18	<18	18	No counts
December 1991	25	<18	25	No counts
January 1992	30	<18	30	No counts
November 1999	18	<18	18	<18
December 1999	30	30	25	<18
January 2000	54	54	<18	<18
February 2000	20	<18	20	<18
December 2000	24	<18	24	<18
January 2001	22	<18	22	<18
January 2002	22	22	<18	<18
February 2006	22	22	<18	22
December 2012	24	<18	<18	24
January 2013	22	<18	<18	22
November 2013	22	22	<18	<18
December 2013	27	27	<18	21
January 2014	22	<18	<18	22
February 2014	21	21	<18	<18
November 2014	22	22	<18	20
December 2014	28	28	<18	<18
February 2015	24	24	<18	<18
Total months	20 months out of a possible *209 survey months	10 months out of a possible *209 survey months	8 months out of a possible *209 survey months	6 months out of a possible *170 survey months

*Possible survey months for potentially detecting Latham’s Snipe at Edithvale-Seaford Wetlands August, September, October, November, December, January, February and March based on the species known occurrence in southern Australia.

4.6.2 Mammals

A total of 28 species of mammal has been recorded at the Ramsar site as listed in databases (Ecology Australia 2001; KBR 2009; Appendix 5). The Bush Rat *Rattus fuscipes* was also reported to occur at Seaford Wetlands during a Monash University study in the mid-1980’s, but hasn’t been recorded since then (Ecology Australia 2001). The Southern Freetail Bat *Mormopterus sp. 4* was definitively recorded at both Edithvale and Seaford Wetlands in 2009 by Parsons Brinckerhoff (2009) bringing the total number of mammals recorded to 29 (Appendices 5 and 6; see below).

There have been no records of the EPBC Act- and FFG Act-listed Southern Brown Bandicoot *Isodon obesulus obesulus* from the area since 1980 (DELWP 2016).

The EPBC Act- and FFG Act-listed Grey-headed Flying-fox *Pteropus poliocephalus* has been recorded in the region more recently and is likely to occur at the wetlands (see below). The Swamp Rat *Rattus lutreolus* was last recorded at Seaford Wetlands in 2005 (DELWP 2016; Will Steele, Melbourne Water, pers. comm.) and still potentially occurs there.

The Rakali or native Water Rat *Hydromys chrysogaster* has recently been identified from footprints as being present at the wetlands in early-2016 (Paul Rees, Melbourne Water, pers. comm.).

The Swamp Wallaby *Wallabia bicolor* has also recently been identified from footprints as being present at the wetlands in early-2016 (Paul Rees, Melbourne Water, pers. comm.).

Feral animals that occur include the Fox *Vulpes vulpes*, Cat *Felis catus*, Domestic Dog *Canis familiaris* and European Rabbit *Oryctolagus cuniculus* (Appendix 5). Domestic pigs *Sus scrofa domesticus* were recently illegally released at Seaford Wetlands, but have since been removed (Paul Rees, Melbourne Water, pers. comm.).

Eastern Grey Kangaroo

In 1985, seven Eastern Grey Kangaroos *Macropus giganteus* and three pouch young from Healesville Sanctuary were introduced to the Edithvale South Wetlands under permit to control grass growth and minimise fire risk and promote conservation and enjoyment of wildlife in an urban environment (Ecology Australia 2001; KBR 2009; Ecoplan Australia 2015). Edithvale South Wetland is enclosed by a 2.1 m high fence and the population is therefore enclosed (Ecoplan Australia 2015). Growth of the isolated population is initially thought to have been relatively slow as a result of environmental factors, such as Fox (*Vulpes vulpes*) predation (KBR 2009). In 2001, Ecology Australia (2001) noted that the population comprised 25–30 kangaroos. In 2005, the population was estimated to be 53 kangaroos (KBR 2009; Ecoplan Australia 2015). Melbourne Water installed drinking troughs and shelters for the kangaroos in 2009, as the organisation is obliged to ensure adequate resources for the population (KBR 2009). In March 2016, the captive population was estimated to be 52 kangaroos inside the enclosure, which exceeds the sustainable carrying capacity of 16–27 animals as recommended in the current Kangaroo Management Plan (Paul Rees, Melbourne Water, pers. comm.).

Historical management of the population included vasectomy of three males in 1995, and vasectomy of three males and treatment of 19 females with levonorgestrel contraceptive implants in 2009 (Ecoplan Australia 2015; VWSA 2016).

To reduce the population to the sustainable level, Melbourne Water commissioned Strathbogie Wildlife [including VWSA (VWSA 2016)] to remove a number of animals from the population and supply the females with medium- to long-term contraception. The work undertaken in April resulted in the removal of 32 kangaroos from the population, and implantation of nine females with contraception. This left 20 kangaroos at Edithvale comprising the nine implanted females and 11 kangaroos which were not captured, three of which are females that received contraceptive implants in 2009 (VWSA 2016). VWSA (2016) used population and sex ratio data to estimate that there may only be two to three female kangaroos capable of breeding.

Bat species

Microbat surveys have been undertaken at both Edithvale and Seaford Wetlands using Anabat call detectors (Parsons Brinckerhoff 2009) and follow-up targeted harp trapping at Edithvale Wetlands (Parsons Brinckerhoff 2010) (Appendix 6). These surveys recorded the following results:

- Eight species were positively identified at Edithvale Wetlands (Chocolate Wattled Bat *Chalinolobus morio*, Gould's Wattled Bat *Chalinolobus gouldii*, Southern Freetail Bat, Large Forest Bat *Vespadelus darlingtoni*, Southern Forest Bat *V. regulus*, Little Forest Bat *V. vulturnus*, White-striped Freetail Bat *Tadarida australis* and Lesser Long-eared Bat *Nyctophilus geoffroyi*);
- Five species were positively identified at Seaford Wetlands (Gould's Wattled Bat, Southern Freetail-Bat, Large Forest Bat, Southern Forest Bat and White-striped Freetail Bat);
- Species groups/complexes recorded at the Wetlands for which calls are indistinguishable and cannot be identified to species levels include:
 - Eastern Freetail Bat/Southern Freetail Bat *Mormopterus* sp. 2 and *Mormopterus* sp. 4, Forest Bat *Vespadelus* species and Long-eared Bat *Nyctophilus* species at both Edithvale and Seaford Wetlands; and
 - Gould's Wattled Bat/*Mormopterus* species and Large Forest Bat/Eastern Bent-wing Bat *Miniopterus schreibersii oceanensis* at Seaford Wetlands.
- 'Potential' calls of three other bat species were obtained during the Anabat surveys, including:
 - The Eastern Bent-wing Bat, listed under the FFG Act 1988, and Eastern Broad-nosed Bat *Scotorepens orion* (not threatened) at both Edithvale and Seaford Wetlands; and
 - Yellow-bellied Sheathtail Bat *Saccolaimus flaviventris*, listed under the FFG Act, at Seaford Wetlands.

Of these three species, Eastern Broad-nosed Bat and Yellow-bellied Sheathtail Bat could be expected to occur at the wetlands (Rob Gration, EcoAerial, pers. comm.). The Yellow-bellied Sheathtail Bat is only vagrant to southern Victoria so wouldn't occur regularly (Rob Gration, EcoAerial, pers. comm.). The Eastern Bent-wing Bat has a low likelihood of occurrence due to the lack of suitable roosting habitat i.e. caves, mines, aqueducts (Rob Gration, EcoAerial, pers. comm.).

The Grey-headed Flying-fox *Pteropus poliocephalus* (EPBC Act and FGG Act) and Eastern Freetail Bat *Mormopterus* sp. 2 (not threatened) were recorded at the Eastern Treatment Plant, and both species probably also occur at Edithvale-Seaford Wetlands (Parsons Brinckerhoff 2009, 2010; Rob Gration, EcoAerial, pers. comm.).

Therefore, 11 bat species potentially occur at the Edithvale-Seaford Wetlands, including two threatened species: Grey-headed Flying-fox (EPBC, FGG) and Yellow-bellied Sheathtail Bat (FFG).

4.6.3 Frogs

Ten species of frog have been recorded at the wetlands (Ecology Australia 2001; KBR 2009; Appendix 5). Threatened species recorded include the Growling Grass Frog *Litoria raniformis* (EPBC Act- and FFG Act), which was last recorded in 1988 to the south of Edithvale (DELWP 2016). The closest records of Southern Toadlet *Pseudophryne semimarmorata* (vulnerable under DSE 2013) occur at Braeside Park (in 1987 and 1990). There are more recent records (2013) from Langwarrin Flora and Fauna Reserve

(DELWP 2016). The Southern Toadlet is very cryptic and difficult to detect, but could still occur at Edithvale or Seaford Wetlands; particularly Edithvale Wetlands. Haswell's Froglet *Paracrinia haswelli* is not considered to be threatened, but it would be considered notable if it still occurred at the wetlands. It is an uncommon frog species that prefers dense, wet, heathy habitat near waterbodies such as swamps, ponds, dams and creeks but may also occur in forest, woodland and shrubland near waterbodies (Wildlife Profiles 1999, Ecology Australia 2005c, Ecology Partners 2007; DSE 2004, DELWP 2016).

Species recorded in recent times at the wetlands during the monthly bird surveys are the frog species that have persisted in urban development and commonly occur in Greater Melbourne: Common Froglet, Spotted Marsh Frog, Striped Marsh Frog, Southern Bullfrog and Southern Brown Tree Frog (see Birds Australia 2006, 2007, 2008, 2010 and 2011; BirdLife Australia 2013a, 2013b 2015 and 2016).

The most favourable habitats for frogs at the wetlands are (see e.g. BirdLife Australia 2016; and Andrew Silcocks, BirdLife Australia, pers. comm.):

- Most of Edithvale South (i.e. Zone 1 of BirdLife Australia or Cell ES1 of GHD 2006);
- BirdLife Australia Zones 5 and 7 of Edithvale North (Cells EN1 and EN3a of GHD 2006);
- The freshwater swamp on the eastern side (i.e. wetland in south-east of Zone 1 of BirdLife Australia, and Cell SSE2 of GHD 2006); and
- The wetland near Seaford North Primary School (wetland in northwest of Zone 1 of BirdLife Australia and northwest of Cell SCW1 of GHD 2006); and
- The wetland in northeast of Zone 1 of BirdLife Australia and east of Cell SN1 of GHD (2006).

Edithvale Common was reported to regularly support good frog habitat prior to drainage improvements. It still provides good habitat for frogs, but only during very wet periods (Paul Rees, Melbourne Water, pers. comm.).

4.6.4 Reptiles

There are no known reptile surveys undertaken at Edithvale-Seaford Wetlands Ramsar Site (no survey results are listed in the VBA, DSE 2016).

A total of 21 reptile species has been recorded in the region (Appendix 5). None of these species are considered to be threatened. The Eastern Long-necked Turtle is classified as data deficient in Victoria by DSE (2013) and is widespread in wetlands of Greater Melbourne.

SKM (2011) considered that potential habitat was available for the Swamp Skink *Lissolepis coventryi* (FFG Act and classified as vulnerable in Victoria by DSE 2013). Surveys using Elliott trapping, remote-sensing camera and artificial shelters would be needed to confirm the presence or otherwise of this species.

4.6.5 Fish

No fish surveys are known to have been undertaken at Edithvale-Seaford Wetlands Ramsar Site (no survey results are listed in the VBA, DSE 2016).

Four native fish species and one exotic species have been recorded in the region (Appendix 5). None of the native fish species are considered to be threatened. Common Galaxias and Short-finned Eels were

reported to be abundant in Seaford Wetlands in 1985 (see Ecology Australia 2001), and Short-finned Eels are still regularly seen at Seaford Wetlands (Paul Rees, Melbourne Water, pers. comm.). The introduced Mosquitofish or Plague Minnow/Eastern Gambusia *Gambusia holbrooki* is present in the wetlands and is highly invasive and capable of tolerating a wide range of environmental conditions. It is an aggressive competitor that potentially out-competes native fish species. It also preys on frog eggs and tadpoles and appears to differentially affect frog species (Morgan and Buttemer 1996; Pyke and White 1996 and 2000; Komak and Crossland 2000; Ecology Australia 2001; Pyke 2005).

SKM (2011) considered that further survey work was required to confirm the presence or otherwise of the Dwarf Galaxias *Galaxiella pusilla*, listed as Vulnerable under the EPBC Act 1999 and as a threatened taxon under the FFG Act 1988 and classified as endangered in Victoria by DSE (2013), which is known from the Boggy Creek catchment. We consider that due to the disconnection of the wetlands from their natural catchments, and factors, such as salinity in some wetlands (e.g. particularly Seaford), this species has a low likelihood of occurrence, but reiterate the point above that no survey data is available for the wetlands to enable a more confident assessment.

4.6.6 Invertebrates

Very limited information is available on the aquatic macro-invertebrate communities of the wetlands. Ecology Australia (2001) reported on benthos and plankton sampling undertaken by Monash University at Seaford Wetlands in 1985, which recorded Polychaete and Oligochaete worms and zooplankton; the latter comprising mostly copepods and cladocerans. A saline water tolerant species of mosquito *Aedes camphotrinus* is known from Seaford Wetlands (GHD 2006). There are no invertebrate species records listed in the VBA within a 5 km radius of the wetlands (DELWP 2016).

Mosquitoes occur in moderate abundance and require management at Seaford Wetland (see Section 9.4.8) as some species can transmit disease causing pathogens (GHD 2015; Melbourne Water 2016). In Victoria, there are three main mosquito-borne diseases: Ross River Virus (RRV), Murray Valley Encephalitis (MVE) and Barmah Forest Virus (BFV). Currently, Seaford Wetlands is the only high risk asset for mosquito outbreaks (DSE 2004; ALS 2010; GHD 2015; Melbourne Water undated^b) due to:

- A moderate average abundance of mosquitoes (based on median abundances of c. 80 adult mosquitoes collected in one light trap in 2010);
- Frequent mosquito control action;
- A very high abundance event of adults collected from the wetland;
- A high percentage composition of vector/pest species (*Aedes camptorhynchus* will bite during the day, night and after sunset and is considered to be a principal vector if RRV and BFV and is known to carry MVE in laboratory conditions, *Culex australicus* is thought to be an amplifier of arboviruses because it carries many arboviruses for different hosts although it does not preferentially bite humans and other species which were collected are potential vectors but in extremely low abundance);
- Numerous complaints from residents;
- RRV was detected in one tested mosquito; and
- A high average of vector larvae species have been found in the Wetland.

4.7 Function

Edithvale-Seaford Wetlands (Brett Lane and Associates 2001; GHD 2006; Ecology Australia 2001; KBR 2009; SKM 2011; DSE 2012):

- support a diversity of habitats for a diverse range of flora and fauna species;
- are an essential component of the regional drainage system, receiving, retaining and diverting stormwater and other surface run-off;
- have a critical flood storage capacity that protects surrounding and downstream properties from flooding; and
- contribute to protecting the water quality of Port Phillip Bay by retaining and treating stormwater and other run-off.

4.8 Cultural Heritage

Heritage Insight (2016a and 2016b) has completed a desktop cultural heritage assessment for both Edithvale and Seaford Wetlands (Appendix 14). Their findings are summarised below.

4.8.1 Edithvale Wetlands

A search of the Victorian Aboriginal Heritage Register identified 13 registered Aboriginal Places within a 2 km radius of the study area, comprising a total of 19 components. Of these 13 sites, four were located within 200 m of the study area. The majority of sites in the area are small deposits of stone artefacts located in both surface and sub-surface contexts. Artefact raw material in these sites is primarily quartz and quartzite, with some silcrete. These sites have all been located on the lowland plain on the former swamp landform and likely represent transitory use of the landscape rather than substantial long term campsites. It should also be noted that contour mapping shows all of these sites are located at a higher elevation than the study area.

Aboriginal sites are commonly found around the margins of the Currum Currum Swamp and on elevated ground within the area. However, recent archaeological work has also located sites underneath swamp deposits. A Cultural Heritage Management Plan by Kennedy, Wheeler and Foley (2012) challenged the assumption that Aboriginal sites will not be found within the former swamp, as they identified archaeological deposits within a sandy dune layer located underneath the swamp deposits which is believed to predate the inundation of Carrum Swamp, dating the finds to at least 7000 years old. However, this sandy layer has not been identified in other CHMPs conducted nearby. More broadly however, archaeological work conducted in Edithvale and surrounds has shown that Aboriginal archaeological sites located within the margins of the former Carrum Swamp are extremely rare.

The site prediction model developed concluded that there is an extremely low likelihood that Aboriginal cultural material will be present in the northern section of the study area, and a low likelihood in the southern section. If Aboriginal cultural heritage material is present it will most likely be in the form of low density stone artefact deposits.

4.8.2 Seaford Wetlands

A search of the Victorian Aboriginal Heritage Register identified that there are 33 registered Aboriginal Places within a 2 km radius of the study area, comprising a total of 34 components. These sites comprise artefact scatters (n=21), object collections (n=2), shell middens (n=4) and scarred trees (n=7). There are two sites located within the activity area and two sites within 200 m of the activity area.

It should be noted that historically, if it was unclear whether a scarred tree was cultural or not, it was still practice to register them as potential scarred trees. Five of the seven scarred trees are potentially culturally scarred trees. Two of the potentially scarred trees are located within 200 m of the study area.

Artefact scatters are the most common site type in the local region, and these sites are generally located around the margins of the former Carrum Swamp within sandy dune deposits and Cranbourne Sands deposits. These sites are located in both surface and sub-surface contexts. Quartz is the most common raw material identified however quartzite, silcrete, chert, basalt and glass artefacts have all been recovered.

One artefact scatter is located within the study area itself which is a scatter of stone artefacts located on a ridgeline around the edge of a swampy low-lying area. The recording archaeologist noted that it was likely that this site represented the remnants of a much larger scatter that had likely been collected previously.

There are four registered shell middens within the search region. Two of these shell middens are located on the foreshore dunes on Seaford Beach, one is associated with Kananook Creek and one is located within the study area which comprises three dense exposures of mussel and pipi shell deposits eroding from the side of an access track.

Within the wider Seaford/Frankston area, shell middens are commonly located along the coastal shoreline and deposits of stone artefacts are frequently located within sandy dune landforms and in deposits of Cranbourne Sands, on elevated ridge lines and in areas associated with the margins of Carrum Swamp. This archaeological record demonstrates that both Carrum Swamp and the coastal area were highly important resources for Aboriginal people and were frequently utilised.

The presence of two Aboriginal sites within the study area also show that despite the Seaford Swamp having previously been part of Carrum Swamp and frequently inundated, fluctuating water levels would have provided access both around and through the swamp. Registered Aboriginal places within the study area are evidence of previous Aboriginal activity within the area

The site prediction model developed concludes that there is potential for further deposits of Aboriginal cultural material within the study area. Areas at higher elevations, such as land along the eastern boundary and in the north western corner should be considered to contain moderate to high Aboriginal archaeological potential. Lower-lying areas should be considered to contain lower archaeological potential. The archaeological potential will also be dependent on the level of ground disturbance which has occurred within any given area.

A field survey of the study area would be able to provide a more detailed archaeological assessment and identify areas of archaeological potential along with any surface Aboriginal sites within the study area.

4.9 Social and Amenity Values

Edithvale and Seaford Wetlands are well renowned for bird watching given the abundance and diversity of species, and regular occurrence of many threatened species. At Edithvale, the state of the art Education Centre (built in 2011) and bird hide (which reopened in August 2016 after being refurbished) offer excellent facilities for the general public, particularly bird watchers and school students of all ages. At the Education Centre, school students, special interest groups and the general community can learn about wetland function, plants and animals that live in a wetland, indigenous and European history of the area and sustainable building design.

Both Edithvale and Seaford Wetlands provide open space with natural values amongst the residential land. Walking tracks traverse the reserves, and visitors can also utilise areas of mown lawn. These facilities provide access for casual walking, birdwatching and dog walking for nearby residents and visitors. There is also increasing pressure from the Down's Estate Community Working Group to utilise Seaford Wetland for running events and exercise, although this may have impacts on wildlife. In an effort to reduce disturbance of wildlife across Edithvale Wetland, some of the ponds in the northern section have been designated for dogs to swim and play in and are fenced out of the main wetland area.

The Friends of Edithvale-Seaford Wetlands volunteer community group holds regular working bees, field days and lectures at both wetlands. The group is involved in the promotion of public appreciation and awareness of the wetlands, ecological education, revegetation and general management of the sites.

4.10 Scientific Research and Long-term Monitoring

The Edithvale-Seaford Wetlands Ramsar site has provided significant opportunities for scientific research and long-term monitoring (Brett Lane and Associates 2001; Ecology Australia 2001; KBR 2009). The diversity of flora and fauna, in particular, has provided opportunities for ecological studies. Edithvale and Seaford Wetlands have been used by Birds Australia and BirdLife Australia for long-term monitoring of bird populations since 1989 and 1994 respectively (Birds Australia 2004, 2006, 2007, 2008, 2010 and 2011; BirdLife Australia 2013a, 2013b, 2015a and 2016a). A mist-netting and bird-banding project commenced at Edithvale Wetlands in 2006 (BirdLife Australia 2015b). Monitoring of frog populations has also been undertaken in association with the long-term bird monitoring.

For flora, research and long-term monitoring into the success of control of Common Reed has been undertaken at the wetlands using vegetation mapping, quadrat monitoring and photo-point monitoring (e.g. see Greet and Rees 2015). Monitoring of vegetation condition (i.e. 'Habitat Hectares' methodology) and weed mapping investigations have also been undertaken. (e.g. see Australian Ecosystems 2011a, 2011b and 2016).

The function of the wetlands in flood mitigation and water quality remediation has provided opportunities for research related to the water industry and hydrology (Ecology Australia 2001; KBR 2009; SKM 2011).

Personnel from Monash University have been using the site for blue carbon assessments for research.

Personnel from Chisholm Tafe have been conducting acid sulfate soil assessments for research.

Long-term monitoring and control of a saline water tolerant species of mosquito (*Aedes camphotrinus*) has been undertaken by Melbourne Water.

Water level monitoring has been undertaken at the wetlands since 2008.

Water quality monitoring has been undertaken at Seaford Wetlands since 2003 and at Edithvale Wetlands since 2009 (see SKM 2011; Section 6.2).

Groundwater monitoring commenced in 2014 with the installation of groundwater bores to monitor groundwater dependent ecosystems (GHD 2014).

4.11 Buffer Opportunities

Inside the Ramsar boundary, the wetlands at the Edithvale site are bounded on their eastern side by dryland habitats that offer a number of opportunities. The largest area of interest is an area of c. 8 ha on the north-eastern side of Edithvale South Wetlands, running from the rear of properties along Amaroo Drive in the south to Edithvale Road in the north and bordered in the east by the rear of properties along Enterprize Avenue and Isabella Close. The dryland area supports managed grassland with scattered non-indigenous plantings (e.g. eucalypts and *Melaleuca* sp.). Considerable opportunities exist to make use of this land to complement the wetland values by the installation of treatment wetlands that would treat water from urban drains prior to entering the Edithvale South Wetlands, plus provide additional habitat for waterbirds and frogs. Another opportunity would be to revegetate this area with appropriate woodland species that originally grew in the area.

As discussed in Section 2.6, the Edithvale Common—which formerly supported good waterbird (e.g. Latham’s Snipe) and frog habitat during most years, but now supports these habitat values intermittently during very wet years—adjoins the Ramsar boundary to the west of Centre Road Drain and the Edithvale North wetlands. It supports open space which is complementary to the Ramsar wetlands. The potential for inclusion of the Edithvale Common into the Ramsar boundary should be investigated. The City of Kingston is currently investigating whether it can re-flood the area to provide the habitat it formerly did prior to drainage works (Paul Rees, Melbourne Water, pers. comm.).

Should Melbourne Water acquire land associated with the Rossdale Golf Course and First Avenue developments, opportunities to include this land in the Ramsar boundary, or use of this land in a fashion that complements/buffers the Ramsar site, should be investigated (see Section 9).

4.12 ‘Downs Estate’

‘Downs Estate’ in the northeast of Seaford Wetlands, is freehold land managed by Frankston City Council that is situated outside of the Ramsar boundary. However, Council’s Planning and Environment Department and Melbourne Water are interested in including the land within the Ramsar boundary (see Sections 2.5 and 2.6).

Downs Estate (c. 20.8 ha) was acquired by Frankston City Council in 2006 prior to the preparation of the last Management Plan (KBR 2009). The dominant habitat type is introduced pasture. The land supports seasonal wetlands in a series of old watercourses, and these periodic wetlands fill only during a wet winter. When the site was grazed, it was favoured by flocks of Flame Robins *Petroica phoenicea* (30–40 robins) during their annual winter migration from their breeding areas at higher altitudes to the lowlands. It also favoured the internationally migrant Latham’s Snipe during the spring. Both of these

species prefer the short grass for foraging (Birds Australia 2004; TBLD and Australian Ecosystems 2005; Andrew Silcocks, BirdLife Australia, pers. comm.). Flocks of Straw-necked Ibis and Australian White Ibis also foraged in this area as they prefer foraging in an open habitat.

Planning zones over the property include (DELWP 2016c):

- A Green Wedge Zone (GWZ); and
- An Urban Floodway Zone (UFZ)

Planning overlays:

- An Environmental Significance Overlay (ESO);
- Land Subject to Inundation Overlay (LSIO); and
- Wildfire Management Overlay (WMO).

Small areas (<0.2 ha) of Aquatic Herbland EVC (EVC 653) were identified in the northwest, northeast and southeast of the property. A larger area (c. 0.8 ha) of Brackish Aquatic Herbland EVC (EVC 537) and a smaller area (<0.10 ha) of Tall Marsh EVC (EVC 821) occur in the southwest corner of the property (Aspect Studios 2012).

The Ephemeral wetlands have potential to be managed for late-winter and early-spring habitat for waders, by pumping water from Wadsley's Drain Pump Station into the 'old watercourse' which runs southwest across the land, including inundation of the adjoining low-lying areas (Paul Rees, Melbourne Water, pers. comm.). The land immediately adjoining the Seaford Wetlands (to the west of the walking trail) is undoubtedly the most important part of the 'Downs Estate', and flooding the low-lying areas would provide habitat for waders and open grassy areas for other bird species.

Frankston Council Officers support the notion to incorporate Downs Estate into the Ramsar boundary. However, the notion would need to be passed at a Council meeting. The area east of shared trail is likely to continue to have some community use, and therefore, it is appropriate for land west of the trail to be included in the Ramsar boundary as is discussed in Section 9.5.1. A recent Council resolution supported some community activities east of the shared trail. While these activities will be carefully controlled under a licence issued by Council, there may be potential for impacts on migratory and other birds due to some activities. This could put additional pressure on the wetlands, particularly if water is returned to western portion of Down Estate and birds start to utilise it more frequently.

5 Status of Ecological Character and Limits of Acceptable Change (LAC)

The mechanism against which change in ecological character is assessed is via comparison with Limits of Acceptable Change (LAC). LAC are defined by Phillips (2006) as:

“...the variation that is considered acceptable in a particular measure or feature of the ecological character of the wetland. This may include population measures, hectares covered by a particular wetland type, the range of certain water quality parameter, etc. The inference is that if the particular measure or parameter moves outside the ‘limits of acceptable change’ this may indicate a change in ecological character that could lead to a reduction or loss of the values for which the site was Ramsar listed. In most cases, change is considered in a negative context, leading to a reduction in the values for which a site was listed”.

The following should be considered when developing and assessing LAC:

- LAC are a tool by which ecological change can be measured. However, LAC do not constitute a management regime for the Ramsar site.
- Exceeding or not meeting LAC does not necessarily indicate that there has been a change in ecological character within the meaning of the Ramsar Convention. However, exceeding or not meeting LAC may require investigation to determine whether there has been a change in ecological character.
- While the best available information was used to prepare the ECD and define LAC for the site, a comprehensive understanding of site character may not be possible as in many cases only limited information and data are available for these purposes. The LAC may not accurately represent the variability of the critical components, processes, benefits or services under the management regime and natural conditions that prevailed at the time the site was listed as a Ramsar wetland.
- LAC can be updated as new information becomes available to ensure they more accurately reflect the natural variability (or normal range for artificial sites) of critical components, processes, benefits or services of the Ramsar wetland. The formal process for this is via the Ramsar Rolling Review, which is a three yearly assessment of ecological character at each Ramsar site (Butcher et al. 2011).

The LAC for the Edithvale-Seaford Wetlands Ramsar Site were developed in the recent ECD addendum (Hale 2016). The Ramsar Rolling Review (DELWP 2016e) provides the most recent assessment of current conditions against LAC, the 2016 assessment is provided in (Table 5). This assessment indicates that the site is currently within the LAC.

Table 5 Summary of assessment against Limits of Acceptable Change for the Edithvale-Seaford Wetlands Ramsar Site.

Key: CPS Critical ecosystem components, processes and services/benefits

Critical CPS	Limit of Acceptable Change	2016 Assessment
Waterbird diversity and abundance	<p>Abundance of waterbirds will not decline below the following (calculated as a rolling five year average of maximum annual count):</p> <ul style="list-style-type: none"> • Total waterbirds – 2500 • Migratory waders – 900 • Australasian waders – 125 • Ducks – 650 • Fishers – 200 • Large wading birds – 100 • Herbivores – 450 	<p>Abundance of waterbirds (2011 – 2015) from Edithvale-Seaford Wetlands was as follows (BirdLife Australia unpublished data):</p> <ul style="list-style-type: none"> • Total waterbirds – 7250 • Migratory waders – 2700 • Australasian waders – 300 • Ducks – 1960 • Fishers – 280 • Large wading birds – 275 • Herbivores – 1300 <p>The site is currently within the LAC.</p>
	<p>Abundance of Sharp-tailed Sandpiper will not decline below 1% of the population as stated in the most recent Wetlands International Population estimate (based on a five year rolling average of annual maximum counts).</p>	<p>Abundance of Sharp-tailed Sandpipers (2011 – 2015) from Edithvale-Seaford Wetlands was 3300 (1.8% of the population) (BirdLife Australia, unpublished data).</p> <p>The site is currently within the LAC.</p>
Waterbird breeding	<p>Breeding of the following species at least once every five years:</p> <p>Black Swan Blue-billed Duck Chestnut Teal Dusky Moorhen Purple Swamphen Swamp Harrier</p>	<p>All species were recorded breeding more than once in the past five years (2011-2015) (Birds Australia 2006, 2007, 2008, 2010, 2012; BirdLife Australia 2013a, 2013b, 2015 and 2016).</p> <p>The site is currently within the LAC.</p>
Physical habitat for waterbirds	<p>Habitat that comprises open water, emergent native vegetation (sedges, rushes and reeds) and exposed mudflats, with no habitat comprising more than 70 percent of the total wetland area for more than five successive years.</p>	<p>The extent of tall marsh dominated by Common Reed <i>Phragmites australis</i> has increased at the Ramsar site over the past two decades from 34 hectares in 1994 to 57 hectares in 2013 (Melbourne Water unpublished data). However, this is not greater than 70% of the wetland area and a mosaic of open water (deep and shallow) exposed mudflats and emergent vegetation is maintained.</p> <p>The site is currently within the LAC.</p>
Threatened species: birds	<p>Presence of Australasian Bittern and Curlew Sandpiper in at least three out of every five years.</p>	<p>Data from BirdLife Australia (2011–2015) indicate:</p> <ul style="list-style-type: none"> • Australasian Bittern = four out of the past five years. • Curlew Sandpiper = four out of the past five years. <p>The site is currently within the LAC.</p>

6 Site Condition

This section addresses the current condition of the vegetation, water quality and hydrology of Edithvale-Seaford Wetlands.

6.1 Vegetation

Edithvale-Seaford Wetlands have undergone much disturbance since pre-European settlement, formerly being part of the much larger Currum Currum Swamp that has now largely been drained and turned into residential land. The existing wetland remnants that comprise the deeper areas of the former Currum Currum Swamp are highly modified from their previous form with changed hydrology, water quality and salinity. Despite these modifications, the existing wetlands appear to be in quite good condition due to the large diversity of flora and fauna species and vegetation communities present that represent many aspects of the former swamp on a smaller scale.

The quality of the vegetation was assessed and mapped by Australian Ecosystems in 2010 and 2015 (Australian Ecosystems 2011a, 2015) following either the Vegetation Quality Assessment (VQA) or Index of Wetland Condition (IWC) assessment methodology. These assessments indicate that the vegetation quality at both sites is generally quite good, with the majority of vegetation containing moderate diversity and moderate to low levels of weed invasion. In general the higher dryland and ephemeral areas of vegetation contain more weeds than the lower permanently wet or longer inundated areas.

To compare the change in vegetation condition over time, Melbourne Water has commenced condition assessment surveys every five years. However, the 2015 survey primarily used the Index of Wetland Condition assessment methodology rather than the Vegetation Quality Assessment (Habitat Hectare) method. This makes comparison of the vegetation quality over time very difficult as the two assessment methods used are not readily comparable. The conclusions drawn in Australian Ecosystems (2015) regarding the overall change in Habitat Hectares present at the site between 2010 and 2015 is not valid as both assessments assigned all zones assessed using the IWC method a default VQA scores of 0.45. This is more pertinent in the 2015 assessment where the majority of the vegetation was assessed using the IWC method.

6.2 Water Quality

Water quality in the Edithvale-Seaford Wetlands is likely to be significantly influenced by saline water intrusion from groundwater, exposure of coastal acid sulphate soils, backflows of saline water in drains during high tides¹ and the quality of stormwater run-off from adjoining urban areas. The presence of sediment, nutrients and toxicants potentially affects the diversity and productivity of the wetlands (Ecology Australia 2001; GHD 2006; KBR 2009; SKM 2011; DSE 2012). Rising salinity is an issue and is likely to result in shifts from freshwater flora and fauna communities to those tolerant of more saline conditions. Furthermore, good water quality is important for sustaining aquatic flora and fauna (SKM 2011; Jacobs 2016).

¹ There are flood flaps to prevent backflows, however, rubbish can jam them open allowing water to backflow.

6.2.1 Water Quality Sampling 2009/2010

Occasional surface water quality sampling was undertaken at Edithvale Wetlands in 2009/2010 by the Waterwatch group (SKM 2011; Jacobs 2016). Water quality varied between wetland cells, but was overall poor with many indicators exceeding SEPP (Waters of Victoria or WoV) objectives due to the source of the water. Cells with wetlands that received direct inputs of stormwater exhibited high turbidity and nutrients, but lower salinity (EN3a and ES1), than other cells. Cell EN2 had particularly high salinity, and receives groundwater as a result of excavation into the groundwater layer in the 1980's; this cell was also suspected to have low pH from potential acid sulfate soils, but no evidence was obtained in water quality samples (SKM 2011; Jacobs 2016).

At Seaford Wetlands, occasional water quality sampling was undertaken by the Waterwatch group from 2003. Overall, water quality was also poor with many indicators exceeding SEPP objectives due to the sources of water (SKM 2011; Jacobs 2016). Low pH (<6), due to exposure of acid sulphate soils, was recorded in most cells. Nutrient concentrations were highest in the northern cells where inflows from Wadleys Road Drain occur (e.g. SN1 and SCW1), and declined as water flows through the system (from north to south). Seaford Wetlands were brackish and salinity increased through the system with the most brackish cells being SSW1, SSW3 and SSE4 in the south. Elevated salinity is due to saline groundwater intrusion, and tidal intrusion from Eel Race Creek and Kananook Creek (GHD 2006; KBR 2009; SKM 2011; Jacobs 2016).

6.2.2 Water Quality Monitoring 2009 Onwards

From September 2009–2015, more continuous records of water quality parameters were collected.

Salinity

Water quality monitoring from September 2009–2015 demonstrated that salinity levels in Seaford Wetlands are highly variable over time and between cells. Despite this spatial and temporal variation, salinity levels within a cell typically peak in mid-to-late summer and decrease through the year (Jacobs 2016).

As a coastal wetland, potentially subject to seawater incursions and with likely interactions with saline groundwater, SEPP (WoV) objectives were unlikely to be met (Table 6). The dissolved solids associated with catchment run-off are also a potential contributing factor to the increased conductivity seen in both sites on Wadleys Drain, which supplies Seaford Wetland northern pool (Cell SN1).

Conductivity measurements recorded may reveal excessive fluctuations over short periods of time, and unexpectedly freshwater recorded at times. As the pools at Seaford evaporate (i.e. there is no outflow), the salt becomes highly concentrated in the remaining water and then leaves a thick salt crust when it has dried out completely. Wadleys Drain receives backflows occasionally from the Patterson Lakes pumps, which pump sea water from Patterson Lakes into Eel Race Creek (Paul Rees, Melbourne Water, pers. comm.).

Freshwater typically has a conductivity of less than 1000 $\mu\text{S}/\text{cm}$, while seawater measurements are typically greater than 50,000 $\mu\text{S}/\text{cm}$. The conductivity in the Wadleys Drain sites ranged from 340 (freshwater) to 20,000 $\mu\text{S}/\text{cm}$. Similarly, Edithvale Wetlands sites ranged from 450 to 20,000 $\mu\text{S}/\text{cm}$. The observations for Seaford Wetlands ranged from 7.1 (levels associated with mountain streams) to 49,500 $\mu\text{S}/\text{cm}$; very nearly seawater (Table 6).

pH

The pH objectives for this region (SEPP WoV) require that the 25th and 75th percentile range should fall between 6.4 and 7.7. Between one and 27 snapshot surveys were conducted from 2009–2015. Limited numbers of surveys at some sites mean that an accurate calculation of percentiles is compromised.

All sites fall within or close to the SEPP (WoV) objectives for pH levels, with median levels for all wetland cells within a suitable neutral range of 7.0-7.5. There are occasional low readings of pH of below 6.0, in cells SSE4, SCW1, SSW3 and SCE2 and in Centre Drain (Jacobs 2016). This is potentially an indicator of Acid Sulfate Soils (ASS), which are present in areas that are influenced by sulphate-rich saline groundwater intrusion (Jacobs 2016). Low pH occurs when ASS is exposed to air and is then rewet. ASS has been identified in cells SSE4 and SSW1; low pH observations are typically recorded between August and November during the wetting cycle of Seaford wetlands (SKM 2011; Jacobs 2016).


The wetlands fall within an area that has been mapped as potentially impacted by ASS (DSE 2010; Figure 10). Measurements of pH ranged from 2.92 to 9.2. These are extreme fluctuations, and may be a result of waxing and waning inputs from catchment run-off (increasing the alkalinity) and coastal acid sulfate soils (increasing the acidity).

Typically, the Wadsleys Road Drain and Edithvale Wetland sites exhibited pH levels that ranged from acceptable to increased alkalinity, while the Seaford Wetland sites exhibited massive fluctuations, with one site ranging between 2.92 and 9.12 over the course of 17 snapshot surveys.

Table 6 Water quality monitoring data summary for salinity, 2009–2015, Edithvale-Seaford Wetlands.

	Wetland Cells	Monitoring Site	Count of EC (µS/cm)	Min of EC (µS/cm)	Average of EC (µS/cm)	Max of EC (µS/cm)
Wadsleys Drain	SN1	DER100	1	4100	4100	4100
		DSE500	2	7400	12700	18000
		DWA100	5	570	10134	20000
		DWA200	7	1270	7081	19600
		DWA300	2	340	435	530
Edithvale Wetlands	EN3	EV1	22	1655	3029	4600
	EN3a	EV2	9	450	2213	4110
	ES1	EV3	8	965	1839	3700
	EN2	EV4	2	20000	20000	20000
Seaford Wetlands	SN1	SF1	15	463	6389	37300
	SSW1	SF10	4	6250	15280	28000
	SCW1	SF2	28	1010	14048	42900
	SCE2	SF3	20	1490	16643	49500
	SSE2	SF4	9	108	1802	4380
	SSE4	SF5	17	7.1	10202	41300
	SCW1	SF6	13	19.7	7797	22500
	*SN1	SF7	14	23.9	13666	39000
	SSW3	SF8	9	41.3	7641	32800
	SCW1	SF9	9	425	8418	32600

*=Centre Drain



Map 3 Central Coast

Prospective Land: land that has the potential to contain Coastal Acid Sulfate Soils

© State of Victoria, Department of Environment and Water 2016
 This publication is copyright. No part may be reproduced by any means without the permission of the Copyright Act 1969.
 Disclaimer
 The publication is made a statement to the State of Victoria and its employees and contractors that the publication is not a liability for any particular person and therefore does not constitute any liability for any person, firm or corporation which may arise from any reliance on any information in this publication.

Figure 10 Map of land that has the potential to contain coastal acid sulfate soils from DSE (2010).

Dissolved Oxygen

Dissolved Oxygen (DO) levels recorded at Edithvale and Seaford Wetlands have been highly variable (Jacobs 2016). Stormwater inputs support high biochemical oxygen demand and can rapidly deplete DO levels. The wetlands are rich in organic plant biomass and decomposition during the die-back period would draw DO levels down. DO levels above 10 mg/L may indicate algal blooms or macrophyte growth in wetland cells (Jacobs 2016).

Turbidity

Median turbidity levels have generally not exceeded 50 NTU, with occasional recordings above 50 NTU in Seaford Wetland cells SCW1 and SCE2 (Jacobs 2016).

Reactive Phosphorous

The Waterwatch group collected Reactive Phosphorous data from April 2009 to November 2011 (Jacobs (2016). Phosphorous levels were found to be highest at the inlets to wetlands, such as in Wadleys Drain (SN1) and Centre Drain (SN1) at Seaford Wetlands and in stormwater receiving wetlands at Edithvale (EN3a and ES1). This reflects significant nutrient removal and stormwater treatment by the wetlands as water moves through the cells, and the important ecosystem service prior to water entering Port Phillip Bay (SKM 2011; Jacobs 2016).

Heavy metals in sediment

Eight heavy metals (arsenic, cadmium, chromium, copper, mercury, nickel, lead and zinc) were tested in sediment across eight sites (EV1–3, SF1–5) between 2012 and 2014. The only two metals to exceed the recommended levels for 95% protection of species (ANZECC and ARMCANZ 2000) were copper and zinc.

Edithvale Wetlands sites passed all copper assessments, while one quarter of assessments of Seaford Wetlands sites failed. Edithvale Wetlands sites failed two out of 10 zinc assessments, and Seaford Wetlands failed seven out of 20 assessments (Table 7).

A number of nutrients were also sampled, however, there are no default trigger values for wetlands to compare against. When comparing the results against the most conservative levels (freshwater lakes and reservoirs), none of the sites tested passed default trigger values for total phosphorus, while comparisons against the least conservative levels (lowland rivers) had a number of sites passing: EV1 (4 out of 4 surveys), EV2 (1 out of 3) and SF4 (3 out of 4). All sites failed tests for Total Nitrogen when compared against any default trigger values, ranging from marine to upland river sites, while all sites but one (SF2 on one occasion) passed the 95% level of protection of species for NH₃. While difficult to compare against relevant trigger values, and noting the small number of replicates for each site (three or four), these results suggest that there is a fairly high nutrient load across all sites.

Table 7 Water quality monitoring data summary for copper and zinc, 2009–2015, Edithvale-Seaford Wetlands.

Cell (Site)	Number of surveys	Number of times exceeding recommended levels	
		Copper	Zinc
EN3 (EV1)	4		
EN3a (EV2)	3		1
ES1 (EV3)	3		1
SN1 (SF1)	4	3	3
SCW1 (SF2)	4	2	1
SCE2 (SF3)	4		1
SSE2 (SF4)	4		
SSE4 (SF5)	4		2

6.3 Hydrology

The Hydrology of the Edithvale-Seaford Wetlands Ramsar Site is described in detail in GHD (2006) and summarised in KBR (2009), SKM (2011), DSE (2012) and Jacobs (2016a).

The Carrum Carrum Swamp is part of a geological trough bounded on the north side by a monoclinical fold near Beaumaris and on the south side by Selwyn's Fault, near Frankston (GHD 2006). Along with many of the coastal lagoons and wetlands in south-eastern Australia, the wetlands were formed as a result of the end of the last Ice Age and sea level rise c. 5,000 to 7,000 years ago (GHD 2006; SKM 2011; Heritage Insight 2016). A significant portion of the Carrum Carrum Swamp was drowned. Characteristics of the former coastal shoreline include former coastal dune systems behind which lay a series of inter-dunal lagoons. Drier and cooler conditions resulted in retreat of the sea establishing the current dune system and a more recent set of inter-dunal lagoons that filled with a mixture of eroded dunal sands and more recent alluvial deposits from the catchments of Dandenong Creek, Eumemmerring Creek, Boggy Creek and smaller creeks to the north (GHD 2006).

When the area was first surveyed in 1866, water from the Carrum Carrum Swamp drained either to the north into Mordialloc Creek or to the south to Kananook Creek. Five wetland-lagoonal complexes were evident, including (GHD 2006; SKM 2011; see Figure 11):

- Carrum Swamp – the main series of lagoons that existed and form the current suburb of Chelsea Heights, and Wannarkladdin and Carrum wetlands. Dandenong and Eumemmerring Creeks discharged into the Carrum Swamp. The Swamp formed the main seasonal pond and header pool that fed Kananook Creek that broke through the parallel dune systems of the semi-tidal creek. Construction of the Pattersons Cut in 1878–9 divided this swamp, resulting in a loss of connection to the major catchment flows;
- Seaford Swamp: a secondary swamp that extended from near the current alignment of Eel Race Drain to a sand dune ridge near the current alignment of Hummerstone Road. The swamp was an off-stream wetland and only filled seasonally with sufficient water flow from Boggy Creek. Interconnection with Carrum Swamp occurred under flood conditions. Outflow was via a small channel along the current alignment of Eel Race Drain;
- Edithvale Swamp: a backwater swamp from the Carrum Swamp with only seasonal connection to both the north and south cells of the wetland. A natural sill divided the cells from Carrum Swamp at the southern end of Edithvale Wetlands (near the current location of Thames Boulevard, Chelsea Heights);
- Mordialloc/Braeside Swamp: was an area of the former inter-coastal swamps where drainages from the north terminated that extended from higher elevations of Mordialloc through to the lowlands of Braeside and Keysborough; it was contained on the north side by higher ground. It was seasonally connected to all other swamps and drained to the old Mordialloc Creek which was a sand barrier estuary that was only breached in flood season. The Swamp was mostly drained by the Dandenong Roads Board in the 1880s with the construction of a drain along a similar alignment to that of the current Mordialloc Main Drain. Small vestiges of the Swamp still exist at Braeside Park and to the south of Woodlands.

- Boggy Creek Swamp: a smaller swamp that encompassed the lower floodplain of Boggy Creek (included Rossiters Road Swamp). The Boggy Creek alignment lay further to the east than its current alignment.

At the time of European Settlement, the Carrum Carrum Swamp was an extensive freshwater wetland that occupied c. 5,260 ha and extended along the eastern shoreline of Port Phillip Bay, from Mordialloc in the north, to Frankston in the south, and as far east as Bangholme. The water source of the Carrum Carrum Swamp came mostly from Dandenong Creek (headwaters in the Dandenong Ranges) and Eumemmering Creek (headwaters at Narre Warren), with smaller contributions from the catchments that currently drain to Mordialloc Creek and Boggy Creek catchment (GHD 2006; KBR 2009; SKM 2011).

Drainage of the Carrum Carrum Swamp commenced in 1868, when channels were constructed across the Swamp to carry water to Mordialloc Creek and Kananook Creek, in order to claim land for grazing and cultivation (GHD 2006).

In 1879, the Patterson Cut, now known as the “Patterson River”, was cut through the Swamp by the Dandenong Roads Board and coastal dunes to Port Phillip Bay to provide further drainage, with the focus on providing agricultural land. The “new river” commenced at the confluence of Dandenong and Eumemmering Creeks, near Bangholme (GHD 2006; SKM 2011).

Extensive flooding in 1889 resulted in the capacities of the small channels and Patterson Cut being exceeded. The Carrum Drainage Trust was formed with a major focus to increase the capacity of the system (GHD 2006).

By early twentieth century, most of the original wetlands area had been drained and land was being used for market gardens and dairy farming (Heritage Insight 2016a and 2016b).

Following extensive flooding in 1923 and 1924, drains were further increased in size by the State Rivers and Water Supply Commission (GHD 2006).

Major floods in 1934 resulted in works on the levee system to increase the capacity and further develop the drainage system (GHD 2006).

By the 1960’s, the former Swamp area had been largely urbanised (Heritage Insight 2016a and 2016b).

Aerial photograph history and contour mapping demonstrates that (Heritage Insight 2016a and 2016b):

- the areas encompassing Edithvale and Seaford Wetlands would have regularly been waterlogged and occasionally inundated (0–1 m above sea level);
- rapid residential development and subdivision during the 1940s for Edithvale Wetlands and 1950s for Seaford Wetlands; and
- significant modifications to drainage, including construction of pools, drains and weirs after 1980 at Edithvale Wetlands and since 1989 at Seaford Wetlands.

In late 1991, Melbourne Water was formed by the amalgamation of the Dandenong Valley Authority and Board of Works and took over management of drainage and floodplain systems.

As result of historic drainage, Edithvale and Seaford Wetlands have been largely isolated from their natural sources of water and most surface inflow is via drains, many of which are controlled (GHD 2006; KBR 2009; SKM 2011; DSE 2012; Jacobs 2016).

6.3.1 Edithvale South Wetlands

Edithvale South Wetlands comprises a natural depression on the south side of Edithvale Road with the lowest point being 0.3 m below sea level (GHD 2006). Despite being farmed for dairy cattle as late as 1969, the morphology of the basin and the peat layer on which it lies is intact, preserving the last and deepest remnant of the original Carrum Carrum Swamp (GHD 2006; KBR 2009; SKM 2011; DSE 2012).

Edithvale South Wetlands are predominantly fed by three drains from catchments to the east which initially entered sediment ponds at the edge of the macrophyte zone (Figure 12). The winter and spring standing water level generally sits at around 0.00 m Australia Height Datum (AHD), but rises up to 0.78 m AHD during rain events and discharges to Edithvale North Wetlands at Centre Swamp Drain; the invert level of the weir is -0.02m AHD, and therefore, any water above that level discharges to Edithvale North Wetlands (Appendix 7). Reduced rainfall during summer and autumn will result in evaporation and water levels drop below 0.00 m AHD, the drawdown creating the critical mudflat foraging habitat for waders (GHD 2006).

The system comprises one main shallow pool (ES1) and three smaller, deeper pools (ES1a, ES1b and ES1c). The invert of the main pool is -0.47 m AHD. It starts to overflow into Centre Swamp Drain above the invert level of the weir (-0.02 m AHD), and is generally c. 45 cm deep, but has been recorded at depths of up to 125 cm deep (i.e. 0.78 m AHD) (Paul Rees, Melbourne Water, pers. comm.).

The system dries-up over late-summer and autumn, facilitating a lush growth of Salt Club-rush (*Bolboschoenus caldwellii*) in late-summer and autumn, the species that sustains the peats and nutrient cycling. Natural drying of the wetland facilitates the control Common Reed *Phragmites australis* invasion.

A manually operated pump on Centre Swamp Drain can pump water when the drain holds water into the northern pool during dry periods (Appendix 7; GHD 2006; KBR 2009; DSE 2012).

6.3.2 Edithvale North Wetlands

Edithvale North Wetland predominantly comprises a series of constructed wetlands within the former floodplain on the north side of Edithvale Road (Appendix 7; Figure 12; GHD 2006; KBR 2009; DSE 2012). Construction commenced in 1987. Immediately north of Edithvale Road, EN1 was constructed as a shallow pool within the peat layers that established a similar morphology and depth to Edithvale South Wetlands and subsequently remains fresh-brackish (GHD 2006; KBR 2009; DSE 2012). EN1 is connected to Edithvale South via a pipe under Edithvale Road which supplies water at around 0.00m AHD.

A number of other pools further north (i.e. EN2, EN3 and EN3a) generally reach deep into the peat layer and sandy substrate and have varying inputs of groundwater. Water levels in EN3 and EN3a are generally responsive to the groundwater table, which varies in height, but can draw down to -1.95 m AHD during drought years. Salinity is therefore a problem in EN3 and EN3a e.g. 10,000 to 12,000 us/cm (GHD 2006). In wetter seasons, the standing water can reach 0.00 to 0.15 m AHD and salinity is lower e.g. 4,000 to 5,000 us/cm (GHD 2006).

The invert level of the northwest wetland (EN3) is -1.9 m AHD, while the invert level of the L-shaped lake (EN2) is -1.95 m AHD. Data for EN3 suggest that it ranges from 0.42 (-1.48 m AHD) to 2.00 (-0.1 m AHD) m deep. Data for EN2 suggest that it ranges from 0.00 m (i.e. empty) (-1.95 m AHD) recorded in Feb 2016, to more than 2.00 m (gauge under water) (>0.15 m AHD) (Paul Rees, Melbourne Water, pers. comm.).

In the west of the system, the water level in the Dog Pond (another constructed wetland), which is open to the public as a swimming hole for dogs, is also controlled by the groundwater table, but can receive water from cell EN2 via an overflow channel at -0.4 m AHD. Inadequate water retention and salinity are considered to be problems in this pond.

Stormwater drains enter the wetlands via sediment ponds EN4 and EN5. EN4 is a deep pool that generally supports freshwater and provides good foraging habitat for Blue-billed Duck *Oxyura australis*, Musk Duck *Biziura lobata* and Australasian Bittern (GHD 2006; KBR 2009; DSE 2012; Jacobs 2016).

A series of weirs control flow of water between EN1, EN2, EN3 and EN3a and limit total drawdown during prolonged dry or drought conditions. Stormwater supplies these cells from a series of drains and overland flow (GHD 2006; KBR 2009; SKM 2011; Jacobs 2016).

6.3.3 Seaford Wetlands

Seaford Wetlands was disconnected from its natural catchment (Boggy Creek) in the 1880's associated with the construction of the Patterson Cut. The wetlands were subsequently drained for agriculture and much of the land surrounding the core wetland area was still being farmed into the 1970's, with some excavation to facilitate drainage by the landowners (GHD 2006; KBR 2009; SKM 2011).

In addition to its disconnection from Boggy Creek, the system no longer receives water from the Carrum Swamp (now Patterson Lakes) and receives increased groundwater from the local drainage system (GHD 2006; SKM 2011).

Saline water can enter the system from three sources (Appendix 7; Figure 13; GHD 2006; KBR 2009; DSE 2012; Jacobs 2016):

1. In 1988, excavation occurred in the south of the swamp to create lakes (GHD 2006; KBR 2009; SKM 2011; DSE 2012). The excavation broke through peat layer resulting in saline water and acid sulphate groundwater entering this section of the wetland;
2. Ground seepage into stormwater drains to the west of the wetlands; and
3. Backflows from marine waters of Kananook Creek via connectivity to the Bardia Avenue drain and Weatherstone Road drainage system.

The hydrology is highly modified and regulated, and surface flows are almost all from stormwater drains (Appendix 7). A central drain, constructed after World War 1, receives pumped low flows from Wadsleys Road Drain into the northern cell (SN1). This central drain delivers water to varying degrees into cells further south. The levee crossing the central drain, and the central and southern flood gates on this drain were upgraded in 2013 (Paul Rees, Melbourne Water, pers. comm.). The wetlands also receive water during very large events (i.e. 1-in-100 year) from the Boggy Creek catchment i.e. overtops the levee (SKM 2011).

To control salinity at Seaford Wetlands, previous management interventions aimed to provide a freshwater supply during winter, spring and summer through a series of cascading pools which run from north to south through the system that constantly dilute and export the salts from the site. Other works included (GHD 2006; SKM 2011; DSE 2012):

- In 1989–1990, the inlet from Eel Race Drain was modified to prevent intrusion of saline water and permit freshwater to enter the wetlands;

- In 1991–1992, construction of a low embankment to enable ponding of freshwater over groundwater and isolate the northern cell (SN1) from saline areas and promote a fresh-brackish system (SKM 2011);
- Interventions to reduce saline intrusion from the drainage system and to intercept the local drains from the west, between Shirley Avenue and Mackenzie Street;
- There are pumps and flood flaps at Bardia Avenue and Weatherston Road to prevent water back-flowing from Kananook Creek; and
- In addition there is a pump system at James Street, which directs saline groundwater intercepted by the local drainage system and directs it around the wetland to the Austin Road Pit.

Seaford Wetlands support (GHD 2006; KBR 2009; SKM 2011; Jacobs 2016): semi-permanent brackish pools in the south which receive stormwater input; a deep, permanent and strongly saline pool in the southwest which has been excavated into the peat layer and exposed acid-sulphate soils; intermittent shallow brackish herb/sedgeland areas in the south; and small marshes fed by stormwater in the east and north.

Melbourne Water is currently investigating whether there is an opportunity to pump water from Boggy Creek into Seaford Wetlands (Paul Rees, Melbourne Water, pers. comm.).

SCE2 has an invert level of 0.19 m AHD, with a range of 0.00 m (0.19 m AHD) to 0.60 m (0.79m AHD) and generally dries out in summer. SCW1 has an invert level of 0.14 m AHD, with a range of 0.00 m (0.14 m AHD) to 0.70 m (0.84 m AHD). The SCE2 and SCW1 operate independently below 0.14 m AHD, but are connected above this point as essentially one large pool, as the levee on either side of the main drain has low points of 0.28 m AHD on the west and 0.31 m AHD on the east. Thus, in an average year, the cells will be disconnected initially in autumn, they will slowly fill until they reach 0.31 m AHD (0.12 to 0.14 m deep), at which point they will be connected over most of winter and spring, until in late-spring and early summer, they draw down below 0.31 m AHD and are no longer connected.

SSW1 has an invert level of 0.08 m AHD and ranges from 0.00 m (0.08 m AHD) to 0.68 m (0.76 m AHD). SSW3 has an invert level of 0.21 m AHD and ranges from 0.00 m (0.21 m AHD) to 0.66 m (0.87 m AHD).




Figure 11 Former Carrum Carrum Swamp, including remnant cells (from GHD 2006).




Figure 12 Location of hydrological cells, drains, weirs and sedimentation ponds and direction of flows for Edithvale Wetlands (reproduced from GHD 2006).




Figure 13 Location of hydrological cells, drains, pipes and weirs and direction of flows for Seaford Wetlands (reproduced from GHD 2006).

7 Risk Assessment

A risk assessment has been undertaken to evaluate the threats identified and their potential impacts to the Ramsar sites values. The risk assessment approach follows a structured and iterative process, with the following steps:

1. Establish the context – existing values and environmental conditions;
2. Identify risks – threats and associated potential impacts; and
3. Analyse risks – assign likelihoods and consequences to determine level of risk.

The methods and detailed results are presented in Appendix 8.

Table 8 summarises the results of the risk assessment for each threat.

These results feed into the Management Plan in Section 9.

Table 8 Summary of risk assessment for the Edithvale-Seaford Wetlands Ramsar Site.

Pressures	Stressors	Risk (highest risk for all potential impacts)
Pollution: urban stormwater	Increased nutrients and sediments	Low
Pollution: urban stormwater	Toxicants	Low
Water management: urban stormwater	Altered water regimes	Medium
Water management: urban stormwater	Increased salinity	High
Urban development and recreation	Litter	Medium
Disturbance of Acid Sulfate Soils (ASS)	Acidity and liberation of heavy metals	Medium
Invasive species	Invasive native plant species: Common Reed <i>Phragmites australis</i> and Cumbungi <i>Typha</i> spp.	High
Invasive species	Invasive native exotic species: Spiny rush <i>*Juncus acutus</i>	Medium
Invasive species	Predators (foxes, cats and rats)	Extreme
Invasive species	Native species: kangaroos	Medium
Invasive species	Grazing animals (pigs, rabbits)	Medium
Recreational activities	Human disturbance (noise)	Medium
Climate change	Increased frequency / intensity of fire	High
Climate change	Sea level rise	Extreme
Climate change	Increased frequency & intensity of storms	High

8 Threatening Processes

8.1 Water Management: Urban Stormwater – Altered Water Regimes

An important water-related threat to waterbirds and waterbird habitats is inadequate water levels in the wetlands cells to provide habitat at critical times to support life history requirements (e.g. ANCA 1996; Brett Lane and Associates 2001; Ecology Australia 2001; Birds Australia 2008; DSE 2012; BirdLife Australia 2016a). Adequate water levels are required for the duration of the migratory wader species visitation. If wetlands dry out too quickly over summer (e.g. by December), waders disperse from Edithvale-Seafood Wetlands. Adequate water levels are also important for the duration of the diving ducks breeding season to complete the breeding cycle (e.g. Blue-billed Duck, Hardhead).

The Edithvale South Wetland dries out over most summers. If this occurs early in summer, it predisposes young Black Swans, which are incapable of flying, to predation by foxes (e.g. BirdLife Australia 2016a). Maintaining adequate water levels is also an issue at Seafood Wetlands, with the wader pool (SCE2) drying out by November or December during drought conditions (SKM 2011).

An important water-related threat is the dominance of emergent vegetation (i.e. Common Reed and to a lesser extent Cumbungi) in response to shallow, stable water levels during the critical growing summer period, reducing the occurrence of other aquatic vegetation and habitat types (e.g. open mudflats for waders) (SKM 2011). Ideally, a natural pattern of wetting and drying that facilitates the maintenance of a mosaic of vegetation types (e.g. shallow open water with submerged or weakly emergent macrophytes to areas of dense emergent macrophytes, is desired (Ecology Australia 2001; SKM 2011).

The priority objective is to provide an inundation regime which inundates the wetlands over winter and early-spring and is then slowly drawn down from September to December to provide habitat for migratory waders, whilst maintaining the deeper pools/wetland cells as permanent wetlands for waterfowl (Ecology Australia 2001; KBR 2009; SKM 2011; DSE 2012).

Ecology Australia (2001), GHD (2006) and SKM (2011) also suggested that insufficient high level freshwater flows into Seafood Wetlands may lead to gradual accumulation of salt, which becomes concentrated especially during drought years. Maintaining the freshwater herbfield was considered to be dependent on seasonal freshwater flushing to avoid a further transition from freshwater to saline vegetation (also see DNRE 2002).

8.2 Water Management: Urban Stormwater – Increased Salinity

Saline water enters Edithvale North Wetland and Seafood Wetlands via excavations made in 1987 and 1988 which broke through the peat layer resulting in saline water and acid sulphate groundwater intrusions. In Seafood Wetlands, saline water also enters via groundwater seepage into stormwater drains to the west of the wetlands, occasional backflows from marine waters of Kananook Creek and saltwater inflow from Eel Race Drain during higher tides (GHD 2006 and 2015; KBR 2009; SKM 2011; DSE 2012). Both Edithvale and Seafood Wetlands will also become more saline with eustatic sea level rise exacerbating tidal intrusions. Increasing salinity will result in a further shift from flora and fauna communities that favour freshwater, to those that can tolerate brackish or saline water (Ecology

Australia 2001; GHD 2006; SKM 2011). Rising salinity is therefore a threat to waterbirds, frogs and vegetation diversity, resulting in an overall decline in habitat diversity (also see DNRE 2002).

Combined, there appears to be an inexorable trend to more saline conditions, with rising sea levels either directly or indirectly through groundwater intrusion, being the major driver (see also Section 6.2.1).

8.3 Pollution: Urban Stormwater – Increased Nutrients, Sediments and Toxicants

Stormwater inflows potentially carry pollutants, nutrients and sediments into the wetlands from urban areas. Sedimentation can result in smothering of aquatic biota and the creation of sites for weed establishment and favour the introduction of exotic or invasive native weed species (e.g. *Phragmites australis* and *Typha* spp.) and reduce the flood carrying capacity of the wetlands by making the wetlands shallower over time. Nutrients can favour conditions suitable for eutrophication and algal blooms of wetlands. Toxicants can be delivered into wetlands which may adversely impact fauna through bioaccumulation (Ecology Australia 2001; DNRE 2002; KBR 2009; SKM 2011; DSE 2012; DEPI 2014b).

8.4 Urban Development and Recreation – Litter

Litter not only harms the appearance of a wetland, but it can also be harmful to aquatic wildlife through entanglement or ingestion (Ecology Australia 2001; DNRE 2002; KBR 2009; SKM 2011; DSE 2012; DEPI 2014b). Litter can also affect functioning of drains and pipes (e.g. through blockages), ultimately affecting the water regime and drainage function of a wetland.

In order to assist with removal of rubbish from Council drains entering the wetlands, Melbourne Water would like to improve the treatment of some of the council stormwater pipes by installing 'daylight' pipes with treatment swales (Paul Rees, Melbourne Water, pers. comm.).

8.5 Disturbance of Potential Acid Sulfate Soils – Acidity and Liberation of Heavy Metals

Potential acid sulfate soils occur naturally in coastal and inland settings. These soils contain metal sulfide minerals, which, if drained, excavated or exposed to air, can react with oxygen and water to form sulfuric acid (H_2SO_4). This can result in the acidification of surrounding soils, groundwater and surface water, which can destroy soil structure and cause severe acidification of waterways. If exposed, actual acid sulfate soils can lead to the release of other contaminants, such as heavy metals, which, when combined with acid, can harm plants and animals, and contaminate drinking water (DSE 2010).

Actual acid sulfate soils can also have health, engineering, social and economic impacts as well as devastating effects on alkaline building foundations and infrastructure and the formation of acid can corrode concrete and steel (DSE 2010; Eco-tainable 2013).

Soil sulfide levels are elevated in unconsolidated saturated Holocene sediments present in the Seafood Wetlands (Eco-tainable 2013). Consequently, careful management of these sediments is required during

any earth works that need to be undertaken and an Acid Sulfate Soil Management Plan will be required for the management of excavated acid sulfate sediments (DSE 2010; Eco-tainable 2013).

8.6 Invasive Species – Exotic Plants

A number of previous ecological studies for the two wetlands have identified weed species occurring within each site. These include reports by Ecology Australia (2001), TBLD and Australian Ecosystems (2005), Australian Ecosystems (2011a, b, 2015). A large proportion of these are weeds of management concern for the protection of biodiversity values. They include 16 noxious weed species in Edithvale and 19 noxious weed species in Seafood, as listed under the *Catchment and Land Protection Act 1994* for the Port Phillip and Westernport Catchment Management Authority region. In addition, seven species in Edithvale and nine species in Seafood are Weeds of National Significance (WONS).

Many other weed species are present within the site as ubiquitous herbaceous species which are not prioritised for management as they are not particularly invasive or have reached the extent of their invasion potential.

The weeds of management concern listed in Section 9.4.6 (Tables 13 and 14) include highly invasive species as well as less invasive species requiring management for other reasons (e.g. species with easily eradicated populations, species with high visual impact). These weed species have been categorised into two management threat categories (high and medium), reflecting the level of invasiveness and impact posed by the species within the reserve. While it is imperative that the focus of weed management is on high threat species, it is important that populations of medium threat species are not allowed to expand in the short term. This is particularly pertinent in the groomed areas where weeds may be more likely to invade.

An important consideration for managing high threat weed species within the reserve is the ongoing seed input from source infestations within and adjoining the reserve. If source populations are identified, the land owners should be engaged to eradicate these infestations.

8.7 Invasive Species – Exotic Grazing Animals (European Rabbit and Hares)

European Rabbits *Oryctolagus cuniculus* occur at Seafood Wetlands, as evident from faeces observed across the site. Although rabbits can degrade biodiversity values by grazing on plants and disturbing the soil profile, their impacts were not observed to be excessive during site visits. Rabbit impacts can contribute to erosion and dispersal of seeds in faeces and on fur, and if unmanaged, can lead to the loss of plant populations, resulting in floristic changes, degraded fauna habitats and weed spread. Previous assessments (Australian Ecosystems 2011, 2015) noted grazing by introduced animals as a threat to Seafood Wetlands and recommended rabbit control programs be implemented at the site. European Hares *Lepus europaeus* have not been confirmed at Edithvale or Seafood Wetlands, however they should be managed alongside rabbits if found to occur at either site.

8.8 Invasive Species – Exotic Terrestrial Predators (foxes, cats and rats)

The impacts of other pest animals are less readily quantified. Domestic and feral cats *Felis catus* undoubtedly use Edithvale-Seafood Wetlands, as cats have been at Edithvale North and South and

Seafood Wetlands; cats will have some impacts to native fauna. Foxes *Vulpes vulpes* are also regularly seen at the wetlands, and fox control using soft-jaw trapping at Seafood Wetland resulted in the capture of 20 foxes over a two-week period in April/May of 2015 (Paul Rees, Melbourne Water, pers. comm.). The Red Fox is prevalent within Melbourne and surrounds and integrated control using baiting in residential areas is difficult at best and unfeasible in an area frequented by domestic dogs. Both cats and foxes can prey on birds, small mammals (e.g. if Swamp Rats are still extant) and frogs, which may impact detrimentally on local populations at the Wetlands.

8.9 Invasive Species – Exotic Predatory Fish

The Mosquitofish is a small freshwater fish introduced from USA and capable of tolerating a wide range of environmental conditions. Mosquitofish prey upon frog eggs and tadpoles and are considered a serious threat to frogs in the Melbourne region (reviewed in Ecology Australia 2014). It has been suggested that Mosquitofish may be associated with the decline of the threatened Growling Grass Frog *Litoria raniformis* in Victoria (Tyler 1997). The EPBC Act-listed Dwarf Galaxias *Galaxias pusilla* has not been recorded at either site, but if present, may also be threatened by predation and/or competition with Mosquitofish (DELWP 2015).

8.10 Out of Balance/Invasive Native Species

8.10.1 Common Reed and Cumbungi

Common Reed *Phragmites australis* Narrow-leaf Cumbungi *Typha domingensis* and Broad-leaf Cumbungi *Typha orientalis* are robust graminoids that grow to 3 m or more in height. These species now cover large areas of both Edithvale and Seafood Wetlands and are expanding in these sites. Common Reed is by far and away the most abundant of these species at Edithvale and Seafood Wetlands, but all are capable of shading out the smaller herbaceous species, reducing diversity and simplifying habitat structure as they expand in distribution. The thick reed beds provide limited fauna habitat, instead restricting the diversity of habitats, particularly open wetlands and mud-flats, available to wetland birds that might otherwise utilise the site.

A grooming regime commenced in 2013 which aims to reduce the density and distribution of Common Reed and recreate more habitat diversity. This is included in a study by Greet and Rees (2015) at Seafood Wetlands.

Consideration has been given to potential implications of the grooming regime. The potential promotion of weeds (now that the dominant vegetation structure has been temporarily removed and as a result of soil disturbance) are managed effectively now that the groomed areas are more accessible. In addition, the resultant mass of organic litter created is raked into piles as it can limit or inhibit natural recruitment of native species (and weeds), potentially deoxygenate the water as the litter decomposes and clog drains after large rainfall events.

8.10.2 Coast Tea-tree and Coast Wattle

Coast Tea-tree #*Leptospermum laevigatum* and Coast Wattle #*Acacia longifolia* subsp. *Sophorae* naturally occur on the dune systems and headlands along the coast (Curtis 1956, Pedley et al. 1983,

Thompson 1989, Morrison and Davies 1991, Whibley and Symon 1992, Entwisle et al. 1996, Court 2001, Bachmann 2009). Neither of these species are considered indigenous at Edithvale or Seafood Wetlands where they can form dense stands effectively outcompeting and limiting the recruitment of indigenous species.

As #Coast Tea-tree is fire sensitive, a reduction in fire frequency has been suggested by many authors as a precursor to invasion by this species (Burrell 1981, Ecology Australia 1987, Molnar et al. 1989). Fire is an essential management tool for eliminating this species as it does not resprout (instead releasing seed en-masse) and many indigenous species respond positively with enhanced post-fire recruitment. Fire has been used very effectively in many nearby regions where heathy and woodland vegetation communities are threatened by #Coast Tea-tree, e.g. by Kingston City Council and Bayside City Council at many of their inland reserves, by Parks Victoria at Wonthaggi Flora and Fauna Reserve and at 'The Pines' in Frankston. In other coastal heaths, e.g. in the Greater Otway National Park near Anglesea, fire is similarly employed to control #Coast Tea-tree and #Coast Wattle invasion (Katrina Lovett, Parks Victoria, pers. comm.). Follow up management post-fire is vital to manage the resultant crop of #Coast Tea-tree and #Coast Wattle seedlings.

8.10.3 Eastern Grey Kangaroo

The Eastern Grey Kangaroo population at Edithvale South Wetland has been slowly rising to 52 kangaroos estimated in March 2016. This is well above the density of 0.6 – 1.0 kangaroos per hectare recommended by DELWP, or 16 – 27 animals (see EcoPlan Australia 2015; VWSA 2016). Prior to management intervention in March 2016, numbers were considered to be at an unsustainable level.

Following the implementation of kangaroo management in March 2016, 20 Eastern Grey Kangaroos remained on-site (VWSA 2016), which is within the sustainable population limits of 0.6 – 1.0 kangaroos per hectare recommend by DELWP (see EcoPlan Australia 2015; VWSA 2016). Of the 20 kangaroos remaining, nine females were implanted with levonorgestrel implants in March 2016, while 11 animals were not darted, including three females implanted in 2009 (see VWSA 2016). For the 11 animals that were not darted, VWSA (2016) assumed a 1:1 sex ratio to conclude that five or six of these animals were females, including three reproductively inactive animals (i.e. implanted) and two or three females capable of breeding. The three inactive females will eventually return to reproductive activity during their life span. VWSA (2016) also noted that 13 females implanted in 2009 and darted in 2016 had not returned to reproductive condition in March 2016, providing confidence in the approach used in 2009 and 2016.

Based on the results outlined above, a life expectancy in the wild of about 15 years, and the fact that seven recently implanted females were over seven years of age and therefore unlikely to return to reproductive activity within their life span, VWSA (2016) concluded that the population should remain relatively stable for the next decade with natural mortality balancing recruitment.

8.11 Recreational Activities – Human Disturbance to Native Wildlife

Recreational activities pose a potential disturbance threat to waterbirds at Edithvale-Seafood Wetlands Ramsar site. Recreational activities undertaken on and adjoining the site include dog walking on- and off-leash, walking, jogging and trail-bike riding. Although Melbourne Water has a dogs-on-leash policy at the wetlands, this is not always obeyed by reserve users.

There is a substantial body of literature on the effects of noise and visual disturbance to waterbirds from human recreational activities and the potential implications of disturbance (e.g. Dowling and Weston 1999; Paton et al. 2000; Weston 2003; Weston and Elgar 2005 and 2007; Antos et al. 2007; Maguire 2008; DSE 2011; Glover et al. 2011, Weston et al. 2012, Taylor 2006; Weston et al. 2009 and 2012; McLeod et al. 2013; Martin 2015). Recreational disturbance has been identified as a significant risk and management issue at a number of coastal Ramsar sites, including at Cheetham Wetlands [part of the Port Phillip Bay (Western Shoreline) and Bellarine Peninsula Ramsar Site), Western Port Ramsar Site and Edithvale-Seafood Ramsar Site (DNRE 2002; Antos et al. 2007; KBR 2009; DSE 2011). The consequences include:

- increased energy expenditure when roosting or foraging birds are forced to fly for varying periods or distances until the disturbance has ceased. Increased movement and decreased foraging time is particularly important for migratory shorebirds which need to gain weight prior to long-distance return flights to their breeding grounds;
- Increased risks of mortality to young of breeding birds due to distraction of breeding adults from protecting nests, eggs and chicks from predators and weather (e.g. heat and cold). Decreased breeding success can result from decreased parental attention to young;
- Disturbance of roosting and foraging birds can result in reductions in breeding success and survival of adults, young and hatchlings having cumulative effects on the population;
- Avoidance of important foraging habitats resulting in a reduction in the area of available foraging habitat for waterbirds;
- Increased levels of competition between birds and mortality associated with a reduction in the area of available foraging habitat if there is a lack of other foraging habitat nearby;
- Sensitive species may be driven out of an area totally.

Of particular relevance is the study of Antos et al. (2007) who investigated the issue of disturbance to shorebirds in relation to urban development and the adjacent Cheetham Wetlands in the south-west of Melbourne. Their study focussed on unauthorised human use, including dogs-off-leash, and concluded that 'buffer creep' was a significant issue for these significant wetlands, with shorebirds effectively avoiding areas of the perimeter of wetlands where human access and associated disturbance occurred, resulting in a significant reduction in the area of available foraging habitat.

Melbourne Water is receiving an increasing number of applications for group active recreation, including running events and other activities, potentially resulting in an increased level of disturbance to waterbirds, including migratory shorebirds.

At Seafood Wetland, Frankston City Council restricts the scale and frequency of public events to minimise disturbance impacts on migratory shorebirds.

Clearly, any event application to Melbourne Water or Council would need to consider the timing of the event in relation to ecological values (e.g. in relation to the period of visitation of migratory shorebirds at Edithvale-Seafood Wetlands or in relation to the period of breeding waterbirds at the wetlands), the location of the event in relation to the ecological values, the number of people involved and level of disturbance, the duration of the event and disturbance factor, and means of reducing or minimising the event and intensity of disturbance.

At Edithvale-Seafood Wetlands, Melbourne Water has intentionally retained stands of Tall Marsh around wetlands, such as along both sides of Edithvale Road and between areas of open space and wetlands, which effectively offer some protection/buffers from busy roads and other recreational activities.

Installation of signage regarding dogs needing to be on-leash inside the Ramsar site and information programs via leaflets forwarded in the mail to residents adjoining the wetlands, should help enforce the dogs-on-leash policy.

There is scope for track, fence and revegetation works to direct users around, rather than through core conservation areas (see Sections 9.5.8 and 9.5.9).

8.12 Climate Change

Melbourne Water have recently requested Jacobs to provide site specific scenarios for Edithvale and Seafood as part of a state-wide assessment of climate change impacts to coastal wetlands (DELWP 2015).

These assessments are based on CSIRO / BOM (2015) modelling of moderate (RCP 4.5) and worse case (RCP 8.5) responses to climate change. Under both scenarios Edithvale and Seafood are highly exposed to the key components including:

- increased eustatic sea level;
- increased storm surge activity;
- higher temperatures;
- lower average rainfall;
- changes in seasonal rainfall with strong declines in winter and spring; and
- overall, more variable rainfall.

Sea level rise modelling shows increasing levels of inundation over the course of this century, with almost total inundation of both wetlands by 2100 (DELWP 2015). Melbourne Water's current thinking is that some mitigation of sea level rise should be possible at Edithvale, but the numerous connections via Kananook Creek and Eel Race Creek would make this very difficult at Seafood.

These assessments suggest that the impact of climate change is the pre-eminent issue for these wetlands. While this current plan has only a seven year timeframe, important decisions are required within this period as to how to manage the transition to increasingly saline conditions. The DELWP (2015) report confirms that the adaptive capacity of these systems is low, so key questions appear to relate to the cost-benefit of mitigation, opposed for example to facilitating or managing change.

8.13 Fire and Fuel Management

Inappropriate fire regimes are a potential threat to local flora and fauna, the underlying peat layer of Seafood Wetlands, cultural heritage sites, Melbourne Water infrastructure, local residents, visitors to the reserve and adjoining residential and commercial assets (KBR 2009; Terramatrix 2013a). Furthermore, significant areas of fuel occur in the reserve, including high grass cover and the extensive stands of

Common Reed *Phragmites australis* which form the dominant EVC at the site, Tall Marsh EVC (Terramatrix 2013a). The most significant fire that occurred in Seafood Wetlands was an unplanned fire in January 2004 that burnt c. 5 ha and ignited the underlying peat layer and could not be extinguished for six months (Terramatrix 2013a).

Current ground-based suppression methods used to extinguish peat fires are highly invasive and likely to penetrate the underlying peat layer and expose acid sulphate soils (Terramatrix 2013a).

Fire management and planning and prevention at Seafood Wetlands are the responsibility of both Melbourne Water and Frankston City Council and is directed by the *Country Fire Authority Act 1958* and a number of documents, including the Carrum Lowlands Fire Protection Plan (Melbourne Water 2007), the Frankston City Council Bushfire Management Strategy (Terramatrix 2013b) and Seafood Wetlands Fire Management Plan (Terramatrix 2013a).

Terramatrix (2013a) found that the existing extent of fuel management works around the perimeter of the reserve was generally of sufficient width to provide the necessary separation distance between housing and/or other assets and flammable vegetation within the reserve. They advised that additional fire protection measures should be undertaken at four key areas at Seafood Wetlands to protect assets such as dwellings. This includes managing these areas as 'Asset Protection Zones' that low fuel loads by ensuring:

- Grass will be no more than 50 mm in height;
- Trees must not over-hang or touch dwellings or assets;
- Shrubs under trees must be separated by at least 1.5 times their mature height;
- Tree canopy separation of at least 2 m; tree canopy cover of no more than 15%; and
- Tree branches pruned to a height of 2 m above ground level.

These recommendations do not pose a significant risk to the ecological values present on-site. However, the associated vegetation removal may require offsetting under Victoria's Native Vegetation Permitted Clearing Regulations. A number of fire protection exemptions may apply under the applicable Environmental Significance Overlay's and under Clause 52.48 of the *Planning and Environment Act 1987*. All fire suppression works are to be undertaken in accordance with *Victorian Best Practice Guidelines for Assessing and Managing Coastal Acid Sulfate Soils* (DSE 2010).

9 Management

This section provides a summary of the management undertaken at the wetlands since the production of the previous Management Plan and the management actions required to maintain or enhance biodiversity values within Edithvale – Seafood Wetlands.

The works undertaken by Melbourne Water at the Edithvale-Seafood Wetlands Ramsar site since the preparation of the last Management Plan (KBR 2009) are provided in Appendix 9. These are summarised as follows:

- The key difference to the last Management Plan with respect to works undertaken to enhancing/maintaining the Ramsar ecological values is the extensive grooming of Tall Marsh to provide more open areas of mudflat for waders, and associated works (e.g. Tall Marsh mapping and vegetation quadrat monitoring, e.g. Greet 2015; Greet and Rees 2015).
- Significant work has also been undertaken on the hydrology of the wetlands, including identification of the optimal flow regimes for the different wetland cells and capital works required to enhance/maintain habitat for waterbirds, and associated upgrades to infrastructure have been undertaken to deliver the water requirements (SKM 2011).
- Groundwater monitoring has been undertaken since 2014 (Figures 14 and 15; GHD 2014).
- Water level gauges have been monitored on a monthly basis.
- Water quality monitoring commenced at Edithvale Wetlands in 2009 (Figures 14 and 15).
- In relation to community education and recreation, the Edithvale Education Centre was constructed at Edithvale South Wetlands in 2011, and the Edithvale South Bird Hide which is now open (as at August 2016) (Paul Rees, Melbourne Water, pers. comm.).
- Monthly bird surveys have continued to be undertaken by BirdLife Australia.
- Weed control is an on-going monthly commitment.
- To assist with predator control at Edithvale North wetlands, a predator-proof fence that is buried 300 mm into the ground was installed.

With regard to this Management Plan, additional items that have been recommended include managing impacts from visitors and their dogs and revegetation. The following management recommendations build upon those detailed in previous reports for the reserve (e.g. TBLD and Australian Ecosystems 2005; KBR 2009; Australian Ecosystems 2011a, b, 2016, Terramatrix 2013, GHD 2006, 2015, SKM 2011, BirdLife Australia 2015 and 2016a) that were reviewed during field inspections.

These issues are to be addressed in a way that will avoid or minimise the risk of triggering sulfuric acid release from the potential acid sulfate soils.

Given that the site is a Ramsar-listed wetland of international significance, protection and enhancement of habitat for a suite of waterbirds is a primary focus of the management actions recommended. Potentially disruptive works (e.g. grooming of Common Reed) are generally undertaken between March and August when many of the migratory shorebird species are absent and the wetlands are driest (allowing easier access). Only maintenance and sampling (e.g. water quality) are undertaken outside of this.

9.1 Management Categories

Management actions are divided amongst Program Maintenance, Capital/Grants Programs and Monitoring and Assessment:

Program Maintenance

- Management actions that will be undertaken on a regular basis and will be implemented by Melbourne Water at Edithvale Wetlands and Melbourne Water and/or Frankston City Council at Seaford Wetland.

Capital/Grants Programs

- Management actions that will involve a large one-off investment of funds or grants.

Monitoring and Assessment

- Biodiversity monitoring and assessment works that are required in accordance with the Ramsar and SoBS Plan, and will be coordinated by a biodiversity specialist from Melbourne Water.

Investigation

- One-off investigations to determine the most appropriate management action(s).

Each management action recommended has been assigned to one of these categories.

9.2 Management Zones and Objectives

Edithvale Wetlands has been divided into three Management Zones, while Seaford Wetlands comprises five. The delineation of Management Zones reflects the differing values, threats and associated management prescriptions required across the reserves. They are mapped in Figure 14 and 15 and described in Tables 9 and 10 along with an outline of the management objectives.

Table 9 Management zones and objectives for Edithvale Wetlands.

Management Zone (MZ)	Management Objective
<p>MZ 1 (wetland vegetation where Common Reed <i>Phragmites australis</i> should not increase in cover)</p> <p>Management Zone 1 covers the wetland vegetation generally in the lower depressions and constructed wetlands. Major threats include overabundance of Common Reed reducing flora and fauna diversity, weeds and potential acid sulfate soils.</p>	<ul style="list-style-type: none"> • Grooming and maintaining the current hydrology regime to suppress Common Reed growth • Control high-priority weed species to maintain biodiversity values • Control foxes and other pest animals, including those that may be using nest boxes
<p>MZ 2 (higher drier Common Reed dominated Tall Marsh)</p> <p>Elevated in the slightly higher and drier areas of Common Reed dominated Tall Marsh vegetation is Management Zone 2. Common Reed dominates these areas with very few other indigenous plant species occurring, but provides crane, rail and bittern habitat. Threats include overabundance of Common Reed reducing flora and fauna diversity, weeds, pet dogs and other pest animals, potential acid sulfate soils and dumping of hard-rubbish.</p>	<ul style="list-style-type: none"> • Rotational grooming of Common Reed to promote regrowth that provides more suitable for cranes, rails and bitterns. • Control high-priority weed species to maintain biodiversity values • Control foxes and other pest animals, including those that may be using nest boxes
<p>MZ 3 (scrub, woodland and lawn areas)</p> <p>Management Zone 3 includes the Swamp Scrub, Damp Sands Herb-rich Woodland and exotic lawn that generally surround the perimeter of the wetlands. For the most part, the understorey of these areas comprises slashed exotic grasses, with a few pockets that comprise indigenous groundstorey species or have been revegetated. Some serious weed species occur in these areas either as naturalised or planted individuals (e.g. Swamp Oak <i>Casuarina cunninghamiana</i>, Giant Honey-myrtle <i>Melaleuca armillaris</i>; see Appendix 10 for mapping).</p>	<ul style="list-style-type: none"> • Control high-priority weed species to maintain biodiversity values • Revegetation and supplementary plantings to increase floristic and habitat diversity • Control foxes and other pest animals, including those that may be using nest boxes

Table 10 Management zones and objectives for Seaford Wetlands.

Management Zone (MZ)	Management Objective
<p>MZ 1 (wetland vegetation where Common Reed <i>Phragmites australis</i> should not increase in cover)</p> <p>The lower elevation wetlands that generally are not or previously were not dominated by Common Reed comprise Management Zone 1. Encroachment by Common Reed is the predominant threat to flora and fauna habitat values. Other threats include pest plants and animals (particularly foxes).</p>	<ul style="list-style-type: none"> • Grooming and maintaining the current hydrology regime to suppress Common Reed growth • Control high-priority weed species to maintain biodiversity values • Control foxes and other pest animals, including those that may be using nest boxes
<p>MZ 2 (Common Reed dominated Tall Marsh)</p> <p>Management Zone 2 includes the slightly more elevated and hence drier areas dominated by Common Reed. Common Reed dominates these areas with very few other indigenous plant species occurring and for the most part provides less suitable fauna habitat values. Threats include overabundance of Common Reed reducing flora and fauna diversity, weeds, pest animals, and potential acid sulfate soils.</p>	<ul style="list-style-type: none"> • Rotational grooming of Common Reed to promote regrowth that provides more suitable for crakes, rails and bitterns. • Control high-priority weed species to maintain biodiversity values • Control foxes and other pest animals, including those that may be using nest boxes
<p>MZ 3 (Woodland and scrub vegetation)</p> <p>Management Zone 3 comprises the remnant and revegetated Damp Sands Herb-rich Woodland and Swamp Scrub vegetation to the north and east of the site. Major threats include weed invasion and pest animals.</p>	<ul style="list-style-type: none"> • Control high-priority weed species to maintain biodiversity values • Control foxes and other pest animals, including those that may be using nest boxes • Undertaken revegetation to supplement diversity and rehabilitate disturbed areas after pest plant and animal control
<p>MZ 4 (Non-remnant areas to be slashed and/or revegetated)</p> <p>These areas primarily consist of non-remnant vegetation with scattered indigenous plantings that are recommended for revegetation to provide additional habitat structure and buffer.</p>	<ul style="list-style-type: none"> • Control high-priority weed species to maintain biodiversity values • Undertake revegetation to provide additional habitat structure and diversity
<p>MZ 5 (adjoining Council owned property outside the Ramsar Boundary)</p> <p>The Council owned Down's Estate (to the north east of the study area) primarily comprises wetland vegetation and pasture. It is recommended that this block be incorporated into the Ramsar Site.</p>	<ul style="list-style-type: none"> • Incorporate Management Zone 5 into the Seaford section of the Edithvale-Seafood Wetlands Ramsar site • If successfully incorporated, prepare a management plan to improve the quality of the vegetation and habitat values




Figure 14 The location of management zones and key monitoring features at Edithvale Wetlands. Refer to Figure 5 for detail of Ecological Vegetation Class (EVC) mapping.




Figure 15 The location of management zones and key monitoring features at Seaford Wetlands. Refer to Figure 6 for detail of Ecological Vegetation Class (EVC) mapping.

9.3 Management Priorities

Priority levels have been assigned to each management action, based on Melbourne Water's (2013b) *Waterways Sites of Biodiversity Significance Strategy – 2013*.

Priority 1 (legislative obligations)

- Actions that are a legal obligation under the *Environment Protection and Biodiversity Conservation Act 1999* (this includes Ramsar obligations), the *Flora and Fauna Guarantee Act 1988* or *Catchment and Land Protection Act 1994*;
- Actions to control threats to the Ecological Character of the Ramsar-listed wetlands and are therefore required actions to avoid breaches under the EPBC Act.
- It is expected that funding will be available to achieve all of these actions.

Priority 2 (best practice management)

- Actions that are not a legal obligation but would be considered best practice management, including:
 - the protection and enhancement of habitat for threatened species
 - actions that are relatively easy to achieve as part of ongoing maintenance
 - actions that provide direct benefit to biodiversity values
 - actions that are in line with community's expectation
 - ongoing monitoring of known biodiversity values
- Actions will be undertaken if funding and resources are available after completing all P1 actions.

Priority 3 (other management priorities)

- Actions that are desirable or recommended but with lower priority in the scheme of management actions to be implemented; these may indirectly benefit flora and fauna, such as signage and trail/track maintenance.
- Actions will be implemented if funding and resources are available after completing all P1 and P2 actions.

By prioritising management actions, Melbourne Water and Frankston City Council will be better informed regarding how to incorporate these management actions into Program Maintenance and Capital Works Programs depending on the level of available funding and resources. Consideration has been given to Melbourne Water and Frankston City Council's responsibilities under legislation as well as the site-specific type and level of threat to biodiversity values.

This management plan has been prepared with consideration of the hydrological and floodplain function of the wetland sites. Care has been taken to ensure that the management actions recommended in this plan do not conflict with the hydrological and/or flood capacity function of the site.

Table 11 lists the management actions to be undertaken to protect and enhance the ecological values of Edithvale-Seafood Wetlands. These actions relate to Ramsar and SoBS obligations. Table 12 summarises the estimated costs to undertake the recommended management actions.

Table 11 Management actions to protect and enhance the ecological values of Edithvale-Seaford Wetlands. CPS = critical components, processes and services are highlighted in bold type.**Key:**

MZ Management Zone EPBC *Environment Protection and Biodiversity Conservation Act 1999* MW Melbourne Water KCC Kingston City Council
 CaLP *Catchment and Land Protection Act 1994* FCC Frankston City Council

Action	Legislation applicable	Management Zone	Risk (based on risk assessment)	Values/CPS	Management Category	Target	Edithvale Wetlands Estimated cost	Seaford Wetlands Estimated cost	Management Responsibility (Seaford)	Timing
Priority 1 – Legislative Obligations										
Where possible, grooming of Common Reed to maintain Ecological Character	<i>EPBC Act 1999</i>	Edith: MZ 1, 2 Seaf: MZ 1, 2	High	Waterbird diversity and abundance; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Program maintenance	Edithvale: MZ 1 groomed annually (c. 33 ha); MZ 2 groomed in a rotational pattern with return frequency determined based on fauna habitat utilisation monitoring (c. 5 ha per year if on five year rotation) Seaford: MZ 1 groomed every two (to five) years depending on access due to seasonal conditions that may prevent slashing for one or more years (c. 61 ha); MZ 2 groomed in a rotational pattern with return frequency determined based on fauna habitat utilisation monitoring (c. 8 ha per year if on five year rotation)	\$114,000 per year (\$3,000 per hectare)	\$115,500 per year (\$3,000 per hectare)	MW	Autumn, after Common Reed has flowered, but before seed set.
Conduct a machinery trial for options to cut Common Reed at Seaford Wetland where soft sediments are restricting grooming machinery	<i>EPBC Act 1999</i>	Seaf: MZ 1	High	Waterbird diversity and abundance; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Investigation	Trial of machinery options completed by Year 3 to determine if alternative mechanically grooming options are viable. This is to occur in conjunction with the above grooming regime and is expected to involve trialing various machinery to cut the Common Reed in locations with soft sediments	NA	\$12,000 one-off cost (over three years) Includes additional investigation and hire costs beyond the grooming regime above.	MW	Autumn, after Common Reed has flowered, but before seed set.
Monitor Common Reed management program performance to maintain Ecological Character	<i>EPBC Act 1999</i>	Edith: MZ 1, 2 Seaf: MZ 1, 2	-	Waterbird diversity and abundance; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Monitoring and assessment	Flora: Quadrats established/assessed in Year 1, and reassessed annually Fauna: Waterbird abundance, diversity and visitation rates assessed across grooming regimes three times per year by year 3	Flora: \$9,000 per year Fauna: \$8,000 per year	Flora: \$9,000 per year Fauna: \$8,000 per year	MW	Flora: Mar/Apr Fauna: Sep–Dec
Where possible, maintain current hydrological regime in wetlands to maintain Ecological Character	<i>EPBC Act 1999</i>	Edith: MZ 1, 2 Seaf: MZ 1, 2	Medium	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Program maintenance	Where possible, wetlands inundated in winter and early spring, with gradual drawdown to the end of December (noting that this is largely driven by rainfall and evaporation with Melbourne Water able to input minimal influence). Ephemeral areas ideally dry between January and April while deeper pools remain inundated year round.	Running costs to be determined	Running costs to be determined	MW	All year

Action	Legislation applicable	Management Zone	Risk (based on risk assessment)	Values/CPS	Management Category	Target	Edithvale Wetlands Estimated cost	Seaford Wetlands Estimated cost	Management Responsibility (Seaford)	Timing
Undertake a cost-benefit analysis of management option associated with climate change to: <ul style="list-style-type: none"> - prolong the system as a freshwater to brackish environment; and - manage and facilitate the inevitable change of the system to a more saline environment. 	<i>EPBC Act 1999</i>	Edith: MZ 1, 2 Seaf: MZ 1, 2	Extreme	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Investigation	Build upon the Jacobs (2016b) impacts of climate change assessment to develop a cost-benefit analysis, completed by Year 6 with recommendations incorporated into the revised Ramsar management plan in Year 7.	\$10,000 for one-off assessment	\$10,000 for one-off assessment	MW	Any time
Monitor abiotic and biotic changes associated with climate change: <ul style="list-style-type: none"> - Salinity and water quality - Groundwater and surface water level - Bird surveys - Floristic surveys 	<i>EPBC Act 1999</i>	Edith: MZ 1, 2 Seaf: MZ 1, 2	-	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Monitoring and assessment	Monthly surveys undertaken to document surface water levels, water quality, salinity and bird populations, Quarterly surveys undertaken to record ground water levels, Annual surveys to monitor floristic changes	Abiotic: \$12,000 per year Floristic: \$6,000 per year (note monthly bird monitoring cost is addressed below)	Abiotic: \$12,000 per year Floristic: \$6,000 per year (note monthly bird monitoring cost is addressed below)	MW	All year
Manage CaLP listed "Regionally Controlled" weeds (Tables 13 and 14), particularly *Spiny Rush	<i>CaLP Act 1994</i>	All	High	Waterbird diversity and abundance; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities	Program maintenance	Targets (eliminate, control, contain) as specified in Tables 13 and 14	\$24,000 per year	\$24,000 per year	MW and FCC	All year, 12 visits/yr
Control foxes with soft-jaw traps, maintaining fencing and destroying dens and other harbour (if present)	<i>CaLP Act 1994</i>	All	Extreme	Waterbird diversity and abundance; and threatened waterbird species. Small mammals, frogs and reptiles.	Program maintenance	Soft-jaw traps checked daily for two weeks between April and May, and data on trap success rates collected and submitted to Melbourne Water Maintain predator-proof fence at Edithvale Wetlands No active warrens/harbour, new dens/harbour fumigated and destroyed within 3 months	\$4,000 per year \$2,000–\$4,000 per year \$0–\$3,000 per year	\$4,000 per year NA \$0–\$3,000 per year	MW and FCC	April-May All year
Control rabbits and hares by destroying warrens and other harbour (if present)	<i>CaLP Act 1994</i>	All	Medium	Flora and vegetation communities.	Program maintenance	No active warrens/harbour, new warrens/harbour fumigated and destroyed within 3 months	\$0–\$3,000 per year	\$5,000–\$10,000 per year (more for the first three years)	MW and FCC	Sep
Undertake mosquito monitoring and control (if necessary)	<i>Health (Infectious Diseases) Regulations 2001 and Public Health and Wellbeing Regulations 2009</i>	Seaf: MZ1	-	Human health.	Program maintenance	Monthly dip netting for larvae and CO ₂ light trapping undertaken in spring and summer (Sept-March/April) If monitoring data indicate that larvae are in high abundance, control with ProLink Briquettes will be undertaken	NA	\$8,000 per year	MW	Sept-Feb

Action	Legislation applicable	Management Zone	Risk (based on risk assessment)	Values/CPS	Management Category	Target	Edithvale Wetlands Estimated cost	Seaford Wetlands Estimated cost	Management Responsibility (Seaford)	Timing
Monthly monitoring of bird populations and water level to detect limits of acceptable change to maintain Ecological Character	<i>EPBC Act 1999</i>	All	-	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species.	Monitoring and assessment	Bird surveys and monitoring of water levels undertaken monthly and annual reporting to Melbourne Water	\$15,000 per year (this is an existing program)	\$15,000 per year (this is an existing program)	MW	All year
Potential Acid Sulfate Soil Management	<i>EPBC Act 1999, CaLP Act 1994, Environmental Protection Act 1970, Coastal Management Act 1995, Planning and Environment Act 1986 and Crown Land (Reserves) Act 1978</i>	All, especially Seaford	Medium	Waterbird diversity and abundance; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Program maintenance	Ensure an Acid Sulfate Management Plan is prepared prior to any excavation works	\$6,000 per one-off assessment	\$6,000 per one-off assessment	MW	Any time
Implement fire management recommendations provided by Terramatrix (2013)	<i>Country Fire Authority Act 1958</i>	Edith: MZ3 Seaf: MZ3, 4	High	Flora and vegetation communities.	Program maintenance	Vegetation within the recommended Asset Protection Zones and Bushfire Moderation Zone managed accordingly from Year 1.	NA	\$8,000 per year	MW and FCC	All year
Priority 2 – Best Practice Management										
Incorporate “Downs Estate” into the Ramsar boundary	-	Seaf: MZ 5	-	Physical habitat for waterbirds;	Capital/Grants Program	Down’s Estate incorporated into Ramsar site by Year 2	NA	\$5,000 per one-off investigation	MW and FCC	Any time
Manage high threat weeds (Tables 13 and 14)	-	All	High	Flora and vegetation communities.	Program maintenance	Targets (eliminate, control, contain) as specified in Tables 13 and 14	\$8,000 per year	\$12,000 per year	MW and FCC	All year, 4-6 visits/yr
Maintain current dog access restrictions (i.e. no dogs in wetlands, on-leash in buffers), install interpretative signage to educate the public of dog and cat impacts	-	All	-	Waterbird diversity and abundance; waterbird breeding; and threatened waterbird species.	Capital/Grants program	Signage installed by Year 2. These can be in the form of regulatory and educational interpretative signage.	\$5,000–\$10,000 for signage/letter drop	\$5,000–\$10,000 for one-off signage/letter drop	MW, KCC and FCC	Any time
Distribute letters to landholders in the local area regarding cat and dog impacts	-	All	Extreme	Waterbird diversity and abundance; waterbird breeding; and threatened waterbird species. Small mammals, frogs and reptiles.	Program maintenance	Letter distributed in Years 2, 5 and 8 to all registered cat and dog owners within 500 m of either wetland.	\$5,000 per year	\$6,000 per year	MW, KCC and FCC	All year
If incursions occurs in future, undertake pig trapping to remove released pigs (as had occurred at Seaford Wetland)	-	Seaford: All	Medium	Flora and vegetation communities	Program maintenance	Pig(s) removed within 6 months of confirming occurrence	NA	NA	MW and FCC	All year
Check nest boxes, document occupants and remove Common Myna nests and bee hives	-	All	-	Waterbird breeding.	Program maintenance	All nest boxes check and cleaned in May each year. Occupancy of nest boxes recorded and submitted to Melbourne Water.	\$1,600 per year	\$1,600 per year	MW and FCC	May

Action	Legislation applicable	Management Zone	Risk (based on risk assessment)	Values/CPS	Management Category	Target	Edithvale Wetlands Estimated cost	Seaford Wetlands Estimated cost	Management Responsibility (Seaford)	Timing
Revegetation of areas disturbed by management works (e.g. weed control, den destruction)	-	All	-	Flora and vegetation communities	Program maintenance	Bare ground revegetated within two months; all plant losses replaced with 90% survival to year 7; all guards removed once plants are established (c. 1-3 years depending on species and health of plant). To be undertaken in accordance with the species lists provided in Appendix 12 and the revegetation prescriptions prepared by TBLA and Australian Ecosystems (2005; Appendix 13)	Guarded: \$11/plant Unguarded: \$3/plant Maintenance: \$2/plant	Guarded: \$11/plant Unguarded: \$3/plant Maintenance: \$2/plant	MW and FCC	Ideally May-Oct
Consolidate Frankston City Council and Melbourne Water boundaries at Seaford Wetland (as per Section 2.5)	-	Seaf: All	-	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Capital/Grants program	Divide Seaford Wetland manager boundaries so that Melbourne Water manages all wetland areas and Frankston City Council manages all the exterior areas.	NA	\$5,000 for one-off investigation	MW and FCC	Anytime
At Seaford Wetland, investigate feasibility of extending the path through Downs Estate and along the northern boundary to create a complete loop track	-	Seaf: MZ 3, 5	-	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Investigation	Investigation undertaken by year 1, and findings implemented by year 2	NA	\$4,000 for one off assessment	MW and FCC	Anytime
Maintain fencing around the wetlands	-	Edith: MZ3 Seaf: MZ2, 3, 4, 5	-	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Program maintenance	Undertake fence checks twice per year and repair breaches.	\$1,000 per year (plus repairs)	\$1,000 per year (plus repairs)	MW and FCC	Anytime
Consider the potential need for fencing or seasonal closure of the internal path at Seaford Wetland, and fencing at Down's Estate if included in the Ramsar site (see proposed fencing on Figure 4).	-	Seaf: MZ 1, 2, 5	-	Waterbird diversity and abundance; waterbird breeding; physical habitat for waterbirds; and threatened waterbird species. Flora and vegetation communities.	Investigation	Investigation undertaken by year 2, and findings completed by year 4.	NA	\$3,000 for one-off investigation	MW and FCC	Anytime
Photo-point monitoring	-	All	-	Flora and vegetation communities	Program maintenance	Monitoring of photo-points undertaken every year	\$1,000 per year	\$1,000 per year	MW	In line with previous monitoring (October and March)
Native vegetation condition monitoring	-	All	-	Flora and vegetation communities	Monitoring and assessment	Habitat Hectare Assessment or other suitable monitoring approach collected at Year 5	\$9,000 for one-off assessment	\$9,000 for one-off assessment	MW	Sep-Dec
Monitor pest plant control performance	-	All	-	Flora and vegetation communities	Program maintenance	Distribution and cover of weeds of management concern assessed each year	\$2,000 per year	\$2,000 per year	MW and FCC	Sep-Nov

Action	Legislation applicable	Management Zone	Risk (based on risk assessment)	Values/CPS	Management Category	Target	Edithvale Wetlands Estimated cost	Seaford Wetlands Estimated cost	Management Responsibility (Seaford)	Timing
Monitor pest animal (rabbit/hare/fox) control performance	-	All	-	Waterbird diversity and abundance; and threatened waterbird species. Flora and vegetation communities	Program maintenance	Inspection for warrens / dens at least once per year (if active warrens found undertake more frequent inspections) BirdLife Australia to continue to record foxes and cats observed during monthly bird surveys	\$1,000 per year (more if active warrens found)	\$1,000 per year (more if active warrens found)	MW and FCC	Sep–Oct
Monitor Eastern Grey Kangaroo population	<i>Wildlife Act 1975 and Wildlife Regulations 2002</i>	Edith (south): MZ 1–3	Low-medium	Flora and vegetation communities	Program maintenance	Undertake annual monitoring of the Eastern Grey Kangaroo population using the sweep count procedure to ensure population is maintained between 16-27 kangaroos (see EcoPlan Australia 2015)	\$5,000 per year	NA	MW	March-April
Priority 3 – Other Management Priorities										
Manage medium threat weeds (Tables 13 and 14)	-	All	High-Medium	Flora and vegetation communities	Program maintenance	Targets (eliminate, control, contain) as specified in Tables 13 and 14	\$8,000 per year	\$8,000 per year	MW and FCC	All year, 4 visits/yr
Investigate potential for using fire as a management tool at Seaford Wetlands for #Coast Tea-tree control in MZ3.	-	Seaf: MZ 1–3	High	Flora and vegetation communities	Program maintenance	Feasibility of using fire as a management tool at Seaford for #Coast Tea-tree investigated by Year 3 and implemented accordingly.	NA	\$5,000 for one-off investigation	MW and FCC	Any time
Revegetation and supplementary plantings in Damp Sands Herb-rich Woodland and Swamp Scrub vegetation	-	Edith: MZ3 Seaf: MZ 3, 4	-	Flora and vegetation communities	Capital/Grants program	- Clustered plantings of shrubby and robust ground layer species in the Damp Sands Herb-rich Woodland vegetation with primarily mown lawn understorey - Supplementary and expansion plantings of Swamp Scrub - Revegetation of non-remnant areas opposite Seaford North Primary School - All plant losses replaced with 90% survival to year 7; all guards removed once plants are established (c. 1-3 years depending on species and health of plant). - To be undertaken in accordance with the species lists provided in Appendix 12 and the revegetation prescriptions prepared by TBLA and Australian Ecosystems (2005; Appendix 13)	Guarded: \$11/plant Unguarded: \$3/plant Maintenance: \$2/plant	Guarded: \$11/plant Unguarded: \$3/plant Maintenance: \$2/plant	FCC	May-Oct
User related issues: rehabilitate unauthorised tracks and disturbances	-	All	-	Flora and vegetation communities	Program maintenance	Negligible unauthorised tracks or disturbances by year 2, new disturbances rehabilitated within 2 months of identification	\$1,000–\$5,000 per incursion (depends on extent of damage)	\$1,000–\$5,000 per incursion (depends on extent of damage)	MW and FCC	All year
User related issues: undertake general litter collection and clear litter traps	-	All	Medium	Physical habitat for waterbirds. Flora and vegetation communities	Program maintenance	Negligible litter observed onsite and in litter traps	\$6,000–\$8000 per year	\$6,000–\$8000 per year	MW and FCC	All year
Mapping and monitoring of significant flora species (Appendix 2)	-	All	-	Flora and vegetation communities	Monitoring and assessment	Distribution and population size mapped and compared in Year 5	\$7,000 for one-off assessment	\$7,000 for one-off assessment	MW and FCC	Oct–Nov
Targeted surveys for reptiles (Swamp Skink)	-	Edith: MZ 1, 2 Seaf: MZ 1, 2	-	Reptiles.	Monitoring and assessment	One targeted survey completed by Year 3	\$8,000 per one-off assessment	\$8,000 per one-off assessment	MW and FCC	Nov-Mar
Fish survey to document fish species occurrence	-	Edith: MZ 1, 2 Seaf: MZ 1, 2	-	Fish.	Monitoring and assessment	Fish survey completed by Year 4	\$6,000 per one-off assessment	\$6,000 per one-off assessment	MW and FCC	Dec–May

Action	Legislation applicable	Management Zone	Risk (based on risk assessment)	Values/CPS	Management Category	Target	Edithvale Wetlands Estimated cost	Seaford Wetlands Estimated cost	Management Responsibility (Seaford)	Timing
Invertebrate survey, with the first priority on aquatic invertebrates and the lower priority on terrestrial invertebrates	-	All	-	Invertebrates	Monitoring and assessment	Invertebrate survey undertaken by Year 6	\$6,000 per one-off assessment	\$6,000 per one-off assessment	MW	Spring

Costs do not consider CPI.

Summary of costs:

Table 12 Summary of estimated costs to undertake priority management actions at each wetland, Edithvale-Seaford Wetlands.

Site	Priority	Estimated costs per year	Estimated costs for one-off assessments
Edithvale Wetland	Priority 1 management actions	\$194,000-\$202,000 ^	\$16,000
	Priority 2 management actions	\$29,600-\$34,600 ^	\$9,000
	Priority 3 management actions	\$15,000-\$21,000 ^	\$27,000
Seaford Wetland	Priority 1 management actions	\$214,500-\$222,500 ^	\$28,000
	Priority 2 management actions	\$29,600-\$34,600 ^	\$26,000
	Priority 3 management actions	\$15,000-\$21,000 ^	\$32,000

^ Does not include revegetation costs of \$11 / guarded plant or \$3 / unguarded plant, and \$2 / plant for maintenance.

9.4 Priority 1 Management Actions (Legislative Obligations)

9.4.1 Grooming of Common Reed (Program Maintenance)

One of the key criteria for assessing the Limits of Acceptable Change (LAC) for the Edithvale-Seafood Wetlands Ramsar Site is the extent of different habitat types; no habitat type should comprise more than 70% of the total wetland area (see Section 5; Table 5).

While Common Reed *Phragmites australis* plays an important role for birds in providing breeding sites and shelter for waterbirds and marshland passerines (e.g. Golden-headed Cisticola *Cisticola exilis*, Little Grassbird *Megalurus gramineus*, Australian Reed-warbler *Acrocephalus australis*), its tall and dense habit makes it largely impenetrable for wildlife to feed in, and being very invasive, the species can spread rapidly over wetlands at the expense of other aquatic plants and open mudflats.

To enhance the habitat diversity for a range of fauna species (particularly significant waterbird species that may visit the internationally significant wetlands) grooming of Common Reed is recommended. Although Common Reed is more abundant, this grooming should also address Cumbungi *Typha* spp. Overabundance which is comparatively a lesser issue. Opening these dense reed-beds to create a low open wetland and/or mudflats should diversify habitats making them more suitable to a larger variety of birds such as waders and other shorebirds.

An ongoing experiment has been undertaken at Seafood Wetlands by Greet and Rees (2015) to monitor the recovery of Common Reed after grooming. This study showed that the Common Reed can be effectively controlled provided the culms are subsequently flooded to approximately >20 cm for several months particularly during the spring-summer growth period. They also determined that grooming Common Reed culms after flowering but before seed set was important as it prevented recharging of below-ground energy reserves (i.e. rhizomes). This combination of timing and adequate inundation resulted in minimal Common Reed regrowth and an opening up of the wetland vegetation, with anecdotal evidence supporting a “significant increase in waterbirds using the wetland since it filled, most notably in and around the area where the *Phragmites* [Common Reed] has been cut” (Andrew Silcocks, Birdlife Australia, pers. comm.). In addition, the autumn slashing is best time to access the wetlands as they are driest and should minimise impacts on breeding wildlife.

In areas where flooding was inadequate for Common Reed suppression, the recovery was rapid (Greet and Rees 2015). However, it is thought that utilisation of the ‘regrowth’ Common Reed by crakes, rails and bitterns will increase significantly when compared to the ‘old growth’ Common Reed which has very low utilisation rates as it is too dense for effective foraging (Paul Rees, Melbourne Water, pers. comm.).

Melbourne Water has been undertaking regular grooming at Edithvale-Seafood Wetlands since autumn 2012. At Edithvale this included the majority of Edithvale Wetlands South excluding a narrow strip around the permitter, and much of the Common Reed at Edithvale Wetlands North (particularly the southern half toward Edithvale Road). At Seafood the grooming was not as extensive, as very soft sediments (even when the wetland was dry) inhibited access for the rubber tracked grooming bobcat.

Further Common Reed grooming is recommended as part of this management plan. The method to undertake Common Reed grooming at Seafood will need to be investigated to overcome boggy conditions – see below. The grooming regime will comprise two scenarios:

1. The wetland areas that are most likely to receive adequate inundation (Management Zone 1) should be groomed on an annual basis at Edithvale and every two (to five) years at Seaford (depending on sediment firmness and machinery access) until Common Reed is adequately suppressed, then groomed on an as needs basis thereafter; and
2. The slightly higher and drier areas (Management Zone 2) should be groomed in 1-ha patches' rotating over a number of years to ensure that habitat diversity is maintained across the site.

The optimum return frequency for the grooming regime needs to be determined based on monitoring of fauna utilisation and vegetation recovery, as discussed in Section 9.4.2 below. This will involve monitoring the diversity and abundance of indigenous plant species as well as weeds (ideally ensuring serious weeds do not increase in cover), and monitoring the utilisation of the various regrowth stages by fauna (returning the grooming to areas once the fauna utilisations rates begin to drop). Given that grooming is a costly management action that may also result in an increase in serious weeds, unnecessary grooming should be minimised where possible if the benefits to fauna habitat utilisation and/or indigenous floral diversity are not being demonstrated.

In an attempt to shield avifauna from some human disturbances a 3 m wide buffer of Common Reed should be retained near walking paths where possible. This may not be suitable in areas close to residential dwellings if fire fuel loads pose a risk. In these situations, the breaks should be strategically positioned to line up with other buffers such as trees on the opposite side of the fence.

Trial of Common Reed Grooming Alternatives

Due to soft sediments at Seaford Wetland restricting access for the rubber tracked grooming bobcat, an investigation into and trial of alternative machinery and/or methods is recommended. Soil firmness should be assessed prior to grooming to determine which areas will be accessible to machinery. Other alternatives include:

- Using smaller tracked or legged (e.g. Menzi Muck) machinery;
- Using machinery with a side arm groomer or basket mower for extra reach;
- Manually slashing the Common Reed either on foot or from a boat, or
- Potentially the use of herbicide to control Common Reed.

9.4.2 Monitoring of Common Reed Management Program Performance (Monitoring and Assessment)

The success of the Common Reed grooming regime should be monitored to evaluate the suppression of this species and Cumbungi, and document the change in diversity and abundance of weeds and indigenous flora species. This monitoring should follow the methods used to monitoring the Common Reed grooming program at Seaford Wetlands (Greet and Rees 2015), preferably (though not essentially) with the addition of more sites to extending the survey into Edithvale Wetland.

In summary, this will involve establishing 5 m x 5 m quadrats at random location in areas of mature Common Reed reed-beds. In line with Greet and Rees (2015), these will mostly be located in relatively low-lying areas along the wetter fringes with similar flooding exposure (i.e. Management Zone 1). A second treatment will also be assessed, with quadrats located in the higher drier Common Reed reed-beds (Management Zone 2). Some control quadrats should be left un-groomed for each treatment. The

quadrats should be surveyed in March/April each year with cover scores being recorded to the nearest 5% for the vegetation/habitat types present (EVC, open water, mudflat, etc.) and each species recorded.

The objective to improve habitat diversity, particularly for waders and other birds, should be assessed by monitoring the response of birds (abundance and diversity as well as frequency of visits) to the altered vegetation structures. Monitoring should occur in a) areas with differing densities and age of Common Reed regrowth, including 'old-growth' Common Reed, b) areas groomed annually as well as intermittently, and c) areas transitioning to more open habitats such as mudflats. There are excellent opportunities to collaborate with external organisations or community groups to implement this action (e.g. Birdlife Australia, Friends of Edithvale-Seafood Wetlands, University of Melbourne).

9.4.3 Maintain Current Hydrology Regime (Program Maintenance)

Although the precise relationship between bird abundances and water levels are not understood (e.g. BirdLife Australia 2016a), general agreement exists that the spread of Common Reed is detrimental to waterbird values through loss of open water habitats, especially mudflats that support large numbers of internationally migratory waders. Furthermore, shallow stable water levels during summer facilitate the growth of Common Reed and the objective is to provide as far as possible the appropriate regime which re-introduces a more natural wetting and drying pattern that facilitates the maintenance of a self-sustaining mosaic of vegetation types and controls the spread of emergent vegetation/Tall Marsh. Therefore, the ideal watering regime would be to fill wetlands to capacity during winter and early-spring, and the slowly drawdown the wetlands into summer to continually expose mudflat habitat for foraging waders (Figure 16; SKM 2011). In autumn, the wetlands would be sufficiently dry enabling grooming of Common Reed. Deeper pools would permanently hold water for waterfowl that require deeper, permanent water during the summer months (e.g. ducks). BirdLife Australia should continue to record water levels at wetland cells during the monthly bird monitoring surveys.

The Wadsleys Pump and regulator gates in the centre and at the bottom of the wetlands are the primary options for Melbourne Water to influence water levels. However, these provide a limited capacity as rainfall and evaporation factors are the biggest drivers (Paul Rees, Melbourne Water, pers. comm.).




Figure 16 Seasonal water level fluctuations for Edithvale-Seaford Wetlands (source: SKM 2011)

9.4.4 Cost-benefit Analysis of Management Actions Associated with Climate Change (Investigation)

The predicted sea level rise resulting from climate change indicates that saline incursions as a result of storm surges will be occurring at both by 2030, increasing in extent in subsequent decades (DELWP 2015a, b). This will inevitably change the ecological character of Edithvale-Seaford Wetlands, and needs to be recognised as an ongoing and irreversible threat.

A cost-benefit analysis should be undertaken for various management options to:

- prolong the system as a freshwater to brackish environment, noting this will only be a relatively short term benefit as the pressure of sea level rise increases over time; and
- manage and facilitate the inevitable change of the system to a more saline environment.

This should build on the findings of the Jacobs (2016b) climate change assessment.

9.4.5 Monitoring of Abiotic and Biotic Changes with Climate Change (Monitoring and Assessment)

To monitor the progressive transition of the sites to a more saline environment as a result of sea level rise, a number of physical and biotic aspects that will readily reflect the changing conditions should be targeted. These include:

- Recording salinity and water quality levels on a quarterly basis;
- Measuring groundwater levels on a quarterly basis;
- Continuing the monthly bird monitoring surveys (and record surface water levels) (Birdlife Australia) and allowing for an assessment of trends in species occurrences and abundances over time; and

- Developing a vegetation monitoring program that will document the changing abundance and diversity of flora tolerant of differing salinity levels. The design should be statistically robust, based on quantitative data (presumably transect based), cost-effective and repeatable. This should be resurveyed on a yearly basis.

A monitoring program should be designed and established to document and analyse changes over time and to allow management to be adapted accordingly.

9.4.6 Manage CaLP Listed 'Regionally Controlled' Weeds (Program Maintenance)

At Edithvale Wetlands, there are 12 noxious weed species listed as 'Regionally Controlled' under the *Catchment and Land Protection (CALP) Act 1994* that require management (eradicate, control, contain), while at Seafood Wetland there are 15 (see Tables 13 and 14). Land owners must take all reasonable steps to prevent the growth and spread of 'Regionally Controlled' species on their land. There is no legal obligation to control weeds listed as 'Restricted' under the CaLP Act, though their spread, either as plants, seeds or contaminants in other material is prohibited. Of the Regionally Controlled weeds identified, Spiny Rush **Juncus acutus* is of highest importance as it poses the biggest threat to wetland values. A large population of this species is present in the adjoining Frankston City Council owned land (Down's Estate) that is proposed for incorporation in to the Ramsar Site (Management Zones 5, see Section 9.5.1 for details). This population should be addressed in conjunction with Frankston City Council as it is an upstream source population that poses a threat to the ecological values of the Ramsar site.

These species were identified during the brief field visits and in the previous ecological reports for the site (TBLA and Australian Ecosystems 2005, Australia Ecosystems 2011a, 2016). Appendix 10 provides the most recent weed mapping. They represent a small proportion of the weed flora, are required to be managed under the *Catchment and Land Protection Act 1994*, and include high and medium threat weed species. At Edithvale, an additional 23 high threat weed and 21 medium threat weed species are listed in Table 13 and addressed as Priority 2 (high threat) and Priority 3 (medium threat) issues. Table 14 lists the additional 21 high threat weed and 26 medium threat weed species identified at Seafood Wetlands, which are addressed as Priority 2 and Priority 3 issues.

Numerous other species may require management in certain circumstances (e.g. to allow for revegetation or to protect specific values), but full-scale management would be untenable. It must also be noted that the weed flora of the site is not static; over time new weed species will colonise, dispersed there by a range of natural agents (e.g. wind and animals). Few recent detailed flora surveys have been undertaken for the entirety of either site, and those that have were not extensive and did not document all species observed. For these reasons, the weeds listed for control here should not be seen as exhaustive. Ongoing monitoring will allow for the identification of new weed species and their incorporation into the management program as appropriate.

Additional notes regarding weed control and herbicide use are provided in Appendix 11.

Friends of Edithvale-Seafood Wetlands regularly undertake weed control activities and should be continue to be supported to do so.

Table 13 Environmental weeds requiring management within Edithvale Wetlands.**Key:**

- * Denotes exotic species
- # Denotes Victorian native species that are introduced to the study area
- § Unclear if this species was recorded at Edithvale or Seaford Wetlands (Ecology Australia 2001)

Status

- R Restricted under the *Catchment and Land Protection (CaLP) Act 1994*
- C Regionally Controlled under the CaLP Act
- W Weed of National Significance

Threat (to biodiversity values, based on expert opinion)

- High Highly invasive and/or transformer species (in given situation), effecting severe more or less permanent deleterious changes in vegetation and fauna habitat, floristic composition and structure, preventing vegetation recruitment, and impacting ecosystem function
- Medium Moderately invasive and/or moderately high threat to vegetation and fauna habitat, floristic composition, structure, and recruitment (may be serious as part of multi-species weed invasions)

Strategy

- Eradicate Aim for elimination of populations within the study area, as well as source populations
- Control Control populations to ensure abundance and distribution does not increase beyond current levels
- Contain Manage all new populations to ensure populations do not increase in distribution or abundance (management should focus in areas of remnant vegetation)

Control Method

- H Herbicide: Applied to foliage with spray, wick applicator, etc.; annuals must be sprayed well before seed ripening.
- CD Cut and Dab: Cut down and concentrated herbicide immediately applied to stump, stems, or leaves.
- DF Drill and Fill: Stem drilled and injected with concentrated herbicide or bark "frilled" and herbicide applied.
- PR Physical removal: plants can be physically removed by hand-weeding or with tools when small and/or isolated but soil disturbance is kept to a minimum.
- Fire Fire used to kill mature plants that will not resprout (intensive follow up will be required to remove seedlings as they germinate).

	Scientific Name	Common Name	Status	Threat	Strategy	Treatment
	CaLP listed 'regionally controlled' weed species					
*	<i>Cirsium vulgare</i>	Spear Thistle	C	Medium	Contain	H
*	<i>Cynara cardunculus</i>	Artichoke Thistle	C	Medium	Eradicate	H
§ *	<i>Cytisus palmensis</i>	Tree Lucerne	C, W	High	Eradicate	H, CP, DF, PR
*	<i>Genista linifolia</i>	Flax-leaf Broom	C, W	High	Eradicate	H, CP, PR
*	<i>Genista monspessulana</i>	Montpellier Broom	C, W	High	Eradicate	H, CP, PR
*	<i>Juncus acutus</i> subsp. <i>Acutus</i>	Spiny Rush	C	High	Eradicate	H
*	<i>Lycium ferocissimum</i>	African Box-thorn	C, W	High	Eradicate	H, CP, DF
*	<i>Moraea flaccida</i>	One-leaf Cape-tulip	C	Medium	Control	H
*	<i>Nassella trichotoma</i>	Serrated Tussock	C, W	High	Control	H
*	<i>Rubus fruticosus</i> spp. Agg.	Blackberry	C, W	High	Control	H, CP, PR
§ *	<i>Senecio jacobaea</i>	Ragwort	C	Medium	Control	H
§ *	<i>Silybum marianum</i>	Variegated Thistle	C	Medium	Control	H
	High threat weeds					
#	<i>Acacia longifolia</i> subsp. <i>Longifolia</i>	Sallow Wattle		High	Eradicate	CP, PR
#	<i>Acacia longifolia</i> subsp. <i>Sophorae</i>	Coast Wattle		High	Eradicate	CP, PR
*	<i>Acacia saligna</i>	Golden Wreath Wattle		High	Eradicate	CP, PR
*	<i>Allium triquetrum</i>	Angled Onion	R	High	Control	H, PR
*	<i>Casuarina cunninghamiana</i> subsp. <i>Cunninghamiana</i>	River Oak		High	Eradicate	CP, DF
*	<i>Coprosma repens</i>	Mirror Bush		High	Eradicate	CP, DF
*	<i>Disa bracteata</i>	South African Orchid		High	Control	H, PR
*	<i>Foeniculum vulgare</i>	Fennel	R	High	Eradicate	H
*	<i>Fraxinus angustifolia</i>	Desert Ash		High	Eradicate	CP, DF, PR
*	<i>Hakea sericea</i>	Silky Hakea		High	Eradicate	CP, DF, PR
*	<i>Hedera helix</i>	English Ivy		High	Eradicate	H, CP, PR
#	<i>Leptospermum laevigatum</i>	Coast Tea-tree		High	Eradicate	CP, DF, PR, Fire
*	<i>Lophopyrum ponticum</i>	Tall Wheat-grass		High	Contain	H
#	<i>Melaleuca armillaris</i> subsp. <i>Armillaris</i>	Giant Honey-myrtle		High	Eradicate	CP, DF
	<i>Melaleuca parvistaminea</i>	Rough-barked Honey-myrtle		High	Eradicate	CP, DF
*	<i>Oxalis pes-caprae</i>	Soursob	R	High	Contain	H
*	<i>Pennisetum clandestinum</i>	Kikuyu		High	Contain	H
*	<i>Phalaris aquatica</i>	Toowoomba Canary-grass		High	Contain	H
#	<i>Pittosporum undulatum</i>	Sweet Pittosporum		High	Eradicate	CP, DF, PR
*	<i>Pinus radiata</i>	Radiata Pine		High	Eradicate	DF, PR
*	<i>Prunus cerasifera</i>	Cherry Plum		High	Eradicate	CP, PR
§ *	<i>Salix cinerea</i>	Grey Sallow	R, W	High	Eradicate	CP, DF
§ *	<i>Typha latifolia</i>	Great Reedmace		High	Control	H, CP, PR
	Medium threat weeds					
*	<i>Acacia floribunda</i>	White Sallow-wattle		Medium	Eradicate	CP, DF
*	<i>Anthoxanthum odoratum</i>	Sweet Vernal-grass		Medium	Contain	H
*	<i>Araujia sericifera</i>	White Bladder-flower		Medium	Eradicate	H, CP
*	<i>Callistemon hybrid</i>	Bottlebrush		Medium	Eradicate	CP
*	<i>Cotula coronopifolia</i>	Water Buttons		Medium	Contain	PR
*	<i>Cynodon dactylon</i> var. <i>dactylon</i>	Couch		Medium	Contain	H
*	<i>Epilobium hirsutum</i>	Great Willow-herb		Medium	Control	H
#	<i>Eucalyptus botryoides</i>	Southern Mahogany		Medium	Eradicate	CP, DF
*	<i>Eucalyptus leucoxylon</i> cv. 'rosea'	Yellow Gum		Medium	Contain	CP, DF
*	<i>Eucalyptus robusta</i>	Swamp Mahogany		Medium	Contain	CP, DF
#	<i>Eucalyptus sideroxylon</i>	Mugga		Medium	Contain	CP, DF
*	<i>Eucalyptus spathulata</i>	Swamp Mallet		Medium	Contain	CP, DF
*	<i>Festuca arundinacea</i>	Tall Fescue		Medium	Contain	CP, DF

	Scientific Name	Common Name	Status	Threat	Strategy	Treatment
*	<i>Juncus articulatus</i>	Jointed Rush		Medium	Contain	H
*	<i>Lagunaria patersonia</i> subsp. <i>patersonia</i>	Pyramid Tree		Medium	Eradicate	CP, DF
#	<i>Melaleuca lanceolata</i>	Moonah		Medium	Contain	CP, DF
*	<i>Paspalum dilatatum</i>	Paspalum		Medium	Contain	H
*	<i>Verbena bonariensis</i>	Purple-top Verbena		Medium	Control	H
§ *	<i>Paraserianthes lophantha</i>	Cape Wattle		Medium	Eradicate	CP, DF
§ *	<i>Salix babylonica</i>	Weeping Willow		Medium	Eradicate	CP, DF
§ *	<i>Senecio vulgaris</i>	Common Groundsel		Medium	Control	H

Table 14 Environmental weeds requiring management within Seaford Wetlands.

Key:

- * Denotes exotic species
- # Denotes Victorian native species that are introduced to the study area
- § Unclear if this species was recorded at Edithvale or Seaford Wetlands (Ecology Australia 2001)

Status

- R Restricted under the *Catchment and Land Protection (CaLP) Act 1994*
- C Regionally Controlled under the CaLP Act
- W Weed of National Significance

Threat (to biodiversity values)

- High Highly invasive and/or transformer species (in given situation), effecting severe more or less permanent deleterious changes in vegetation and fauna habitat, floristic composition and structure, preventing vegetation recruitment, and impacting ecosystem function
- Medium Moderately invasive and/or moderately high threat to vegetation and fauna habitat, floristic composition, structure, and recruitment (may be serious as part of multi-species weed invasions)

Strategy

- Eradicate Aim for elimination of populations within the study area, as well as source populations
- Control Control populations to ensure abundance and distribution does not increase beyond current levels
- Contain Manage all new populations to ensure populations do not increase in distribution or abundance (management should focus in areas of remnant vegetation)

Control Method

- H Herbicide: Applied to foliage with spray, wick applicator, etc.; annuals must be sprayed well before seed ripening.
- CD Cut and Dab: Cut down and concentrated herbicide immediately applied to stump, stems, or leaves.
- DF Drill and Fill: Stem drilled and injected with concentrated herbicide or bark "frilled" and herbicide applied.
- PR Physical removal: plants can be physically removed by hand-weeding or with tools when small and/or isolated but soil disturbance is kept to a minimum.
- Fire Fire used to kill mature plants that will not resprout (intensive follow up will be required to remove seedlings as they germinate).

	Scientific Name	Common Name	Status	Threat	Strategy	Treatment
	CaLP listed 'regionally controlled' weed species					
*	<i>Chrysanthemoides monilifera</i>	Boneseed	C, W	High	Eradicate	CP, PR
*	<i>Cirsium vulgare</i>	Spear Thistle	C	Medium	Contain	H
§ *	<i>Cytisus palmensis</i>	Tree Lucerne	C, W	High	Eradicate	H, CP, DF, PR
*	<i>Eragrostis curvula</i>	African Love-grass	C	High	Eradicate	H
*	<i>Genista linifolia</i>	Flax-leaf Broom	C, W	High	Eradicate	H, CP, PR
*	<i>Genista monspessulana</i>	Montpellier Broom	C, W	High	Eradicate	H, CP, PR
*	<i>Juncus acutus</i> subsp. <i>Acutus</i>	Spiny Rush	C	High	Eradicate	H
*	<i>Lycium ferocissimum</i>	African Box-thorn	C, W	High	Eradicate	CP, DF
*	<i>Moraea flaccida</i>	One-leaf Cape-tulip	C	Medium	Control	H
*	<i>Rosa rubiginosa</i>	Sweet Briar	C	High	Eradicate	CP, H
*	<i>Rubus fruticosus</i> spp. Agg.	Blackberry	C, W	High	Control	H, PR
*	<i>Salpichroa origanifolia</i>	Pampas Lily-of-the-Valley	C	High	Control	H
§ *	<i>Senecio jacobaea</i>	Ragwort	C	Medium	Control	H
§ *	<i>Silybum marianum</i>	Variiegated Thistle	C	Medium	Control	H
*	<i>Ulex europaeus</i>	Gorse	C, W	High	Eradicate	H, CP, PR
	High threat weeds					
#	<i>Acacia longifolia</i> subsp. <i>Longifolia</i>	Sallow Wattle		High	Eradicate	CP, DF
#	<i>Acacia longifolia</i> subsp. <i>Sophorae</i>	Coast Wattle		High	Eradicate	CP, DF
*	<i>Acacia saligna</i>	Golden Wreath Wattle		High	Eradicate	CP, DF
*	<i>Allium triquetrum</i>	Angled Onion	R	High	Control	H
*	<i>Asparagus asparagoides</i>	Bridal Creeper	R, W	High	Control	H, PR
*	<i>Casuarina cunninghamiana</i> subsp. <i>Cunninghamiana</i>	River Oak		High	Eradicate	CP, DF
*	<i>Coprosma repens</i>	Mirror Bush		High	Eradicate	CP, DF
*	<i>Cortaderia selloana</i>	Pampas Grass		High	Eradicate	CP
*	<i>Delairea odorata</i>	Cape Ivy		High	Eradicate	CP, DF
*	<i>Ehrharta calycina</i>	Perennial Veldt-grass		High	Eradicate	H
*	<i>Fraxinus angustifolia</i>	Desert Ash		High	Eradicate	CP, DF
#	<i>Leptospermum laevigatum</i>	Coast Tea-tree		High	Eradicate	CP, DF
*	<i>Lophopyrum ponticum</i>	Tall Wheat-grass		High	Contain	H
#	<i>Melaleuca armillaris</i> subsp. <i>Armillaris</i>	Giant Honey-myrtle		High	Eradicate	CP, DF
	<i>Melaleuca parvistaminea</i>	Rough-barked Honey-myrtle		High	Eradicate	CP, DF
*	<i>Pinus radiata</i>	Radiata Pine		High	Contain	CP, DF
#	<i>Pittosporum undulatum</i>	Sweet Pittosporum		High	Eradicate	CP, DF, PR
*	<i>Prunus cerasifera</i>	Cherry Plum		High	Eradicate	CP, DF, PR
§ *	<i>Salix cinerea</i>	Grey Sallow	R, W	High	Eradicate	CP, DF
*	<i>Tradescantia fluminensis</i>	Wandering Jew		High	Eradicate	H
*	<i>Zantedeschia aethiopica</i>	White Arum-lily		High	Eradicate	H

	Scientific Name	Common Name	Status	Threat	Strategy	Treatment
	Medium threat weeds					
*	<i>Acacia elata</i>	Cedar Wattle		Medium	Eradicate	CP, DF
*	<i>Anthoxanthum odoratum</i>	Sweet Vernal-grass		Medium	Contain	H
*	<i>Cordyline australis</i>	New Zealand Cabbage-tree		Medium	Eradicate	CP, DF
*	<i>Cotula coronopifolia</i>	Water Buttons		Medium	Contain	PR
*	<i>Crassula natans</i> var. <i>minus</i>	Water Crassula		Medium	Contain	PR
*	<i>Cupressus macrocarpa</i>	Monterey Cypress		Medium	Eradicate	DF
*	<i>Cynodon dactylon</i> var. <i>dactylon</i>	Couch		Medium	Contain	H
*	<i>Ehrharta erecta</i> var. <i>erecta</i>	Panic Veldt-grass		Medium	Contain	H
#	<i>Eucalyptus botryoides</i>	Southern Mahogany		Medium	Eradicate	DF
*	<i>Festuca arundinacea</i>	Tall Fescue		Medium	Contain	H
*	<i>Gladiolus undulatus</i>	Wild Gladiolus		Medium	Contain	H
*	<i>Hyacinthoides hispanica</i>	Spanish Bluebell		Medium	Eradicate	H
*	<i>Juncus articulatus</i>	Jointed Rush		Medium	Contain	H
*	<i>Lampranthus tegens</i>	Little Noon-flower		Medium	Control	H
*	<i>Mesembryanthemum crystallinum</i>	Common Ice-plant		Medium	Control	H
*	<i>Oxalis pes-caprae</i>	Soursob	R	Medium	Contain	H
§ *	<i>Paraserianthes lophantha</i>	Cape Wattle		Medium	Eradicate	CP, DF
*	<i>Paspalum dilatatum</i>	Paspalum		Medium	Contain	H
*	<i>Paspalum distichum</i>	Water Couch		Medium	Contain	H
*	<i>Pennisetum clandestinum</i>	Kikuyu		Medium	Contain	H
*	<i>Phalaris aquatica</i>	Toowoomba Canary-grass		Medium	Contain	H
*	<i>Phoenix canariensis</i>	Canary Island Date-palm		Medium	Eradicate	CP, DF
*	<i>Phytolacca octandra</i>	Red-ink Weed		Medium	Control	CP
§ *	<i>Salix babylonica</i>	Weeping Willow		Medium	Eradicate	CP, DF
§ *	<i>Senecio vulgaris</i>	Common Groundsel		Medium	Control	H
§ *	<i>Typha latifolia</i>	Great Reedmace		Medium	Control	H, CP, PR

9.4.7 Control CaLP Listed Pest Animals (Program Maintenance)

Fox control is a high priority action that should be implemented in April/May each year, as has been undertaken during the period of the previous Management Plan. In reserves adjoining residential areas, it is undertaken using soft-jaw trapping as 1080 and PAPP (para-aminopropiophenone) baiting cannot occur within 150 m of a dwelling or 20 m of a waterway or waterbody, effectively eliminating the entire sites from baiting (DEDJTR 2016). The contractor should report on the results, including the number of traps set and number of foxes taken each day, in order to calculate a capture success, and therefore, gauge the success and identify a decline in fox captures over the period of each annual control program.

Maintenance checks of fences (e.g. at Edithvale South Wetland) should be undertaken on a monthly basis with rapid repair where needed to minimise entry of foxes into the site due to non-functioning fences.

European Rabbits *Oryctolagus cuniculus*, European Hares *Lepus europaeus* and Red Foxes *Vulpes vulpes* are listed as established pest animals under the *Catchment and Land Protection (CaLP) Act 1994*. The area of most concern appears to be Seaford Wetlands where foxes and rabbits are known to occur (hare may also occur, though this has not been confirmed). Their harbour (rabbit warrens, fox dens, piles of logs/rubbish) within the property, should be destroyed, ensuring indigenous vegetation is not unduly disturbed in the process. If warrens or dens are identified, they should be fumigated and then destroyed/collapsed. Some harbour can be important habitat for native fauna, hence prior to harbour removal, any impacts on native fauna should be considered. Given the impracticality of undertaking pest animal management on a single property (beyond destroying harbour), baiting/trapping actions would only be recommended if they are part of an ongoing landscape strategy being undertaken by Melbourne Water in collaboration with other management agencies.

All pest animal control should be undertaken by a qualified pest control contractor and follow best practice management principals to minimise suffering to target species. Codes of practice and standard operating procedures for managing vertebrate pests which address animal welfare are available at <http://www.pestsmart.org.au/>.

9.4.8 Mosquito Monitoring and Control (Program Maintenance)

Mosquitoes require management at Seaford Wetland under the *Health (Infectious Diseases) Regulations 2001* and the *Public Health and Wellbeing Regulations 2009* as some species can transmit disease causing pathogens (GHD 2015; Melbourne Water 2016; see Section 4.1.6). Currently, Seaford Wetlands is the only high risk Melbourne Water asset for mosquito outbreaks (DSE 2004; ALS 2010; GHD 2015; Melbourne Water undated^b).

Annual mosquito and larvae monitoring and control has occurred at Seaford Wetland since 2007 after concerns about the high abundance of mosquitoes had been raised by local residents and schools (Melbourne Water 2016). Melbourne Water proactively monitors Seaford Wetlands each year, throughout spring and summer, when mosquito populations are normally at their highest, commencing in September and continuing until the wetlands have dried up, usually in March/April (Paul Rees, Melbourne Water, pers. comm.). If monitoring data indicate that larvae are in high abundance, control strategies are implemented. During a warm year, Melbourne Water may receive complaints about mosquitoes as early as August, in which case Melbourne Water will undertake a field assessment of the wetlands. The same protocols should continue to be followed.

Monthly monitoring should involve assessing the wetland for suitable breeding habitat, sampling water quality and mosquito trapping:

- Dip sampling (10 dips at each location) to detect mosquito larvae using a standard size dipper (1.5 cups/350 ml) and compared to an abundance rating (ALS 2010; GHD 2015; Melbourne Water undated^b); and
- CO₂ light traps powered by a 6 volt battery are installed around the wetland and set overnight to capture adult mosquitoes. The number of adult mosquitoes captured in each light trap is compared with an abundance rating and specimens are sent to the Arbovirology Laboratory to be screened for arboviruses (ALS 2010; GHD 2015; Melbourne Water undated^b).

Mosquito control should involve spreading ProLink Briquettes (see DSE 2004; Garrards 2016; Melbourne Water 2016; Pacific Biologics 2016; Melbourne Water undated^a and undated^b) — a c. 5 cm² brick of 18 g/kg S-methoprene — into water pools. The active ingredient prevents larvae and pupae development into adult mosquitoes, is considered to have an insignificant effect on other aquatic species and does not bio-accumulate. ProLink Briquettes are effective over an area of c. 10-20 m² and should last 150 days (though have found to be effective at Seaford Wetland for three months).

9.4.9 Monitoring of Bird Populations (Biodiversity Monitoring and Assessment)

The Limits of Acceptable Change (LAC) for the Edithvale-Seaford Wetlands Ramsar Site were developed in the recent Ecological Character Description (ECD) addendum (Hale 2016). The Ramsar Rolling Review (DELWP 2016e) provides the most recent assessment of current conditions against LAC, and the 2016 assessment is provided in Section 5 (see Table 5). The LAC are the mechanism against which change in ecological character of a Ramsar site is assessed.

LAC for Edithvale–Seaford Wetlands Ramsar relate to the (Table 5):

- abundance of bird in particular waterbird guilds;
- frequency and abundance of EPBC Act-listed threatened species, the Australasian Bittern and Curlew Sandpiper;
- abundance of EPBC Act-listed migratory Sharp-tailed Sandpiper;
- annual occurrence of breeding waterbirds; and
- extent of different habitat types (i.e. with no habitat comprising more than 70% of the total wetland area).

Therefore, a high priority action is that monthly bird monitoring and water level monitoring undertaken by BirdLife Australia continue to ensure the ecological character of the Ramsar site is maintained within the limits of acceptable change. The surveys undertaken by BirdLife Australia since 1989 have played a key role in developing the LAC, undertaking the 2016 assessment and determining that the LAC were met.

9.4.10 Potential Acid Sulfate Soil Management

Potential acid sulfate soils are well known from the wetlands and broader area (see Sections 3.3 and 8.5; Figure 10). For any works involving excavation into the soil, an Acid Sulfate Soil Management Plan needs to be prepared in accordance with DSE (2010). This is a legal requirement under *Environment Protection and Biodiversity Conservation (EPBC) Act 1999*, *Catchment and Land Protection (CaLP) Act 1994*, *Environmental Protection Act 1970* and the *Coastal Management Act 1995* (through the *Victorian Coastal Acid Sulfate Soils Strategy* (DSE 2009c)).

9.4.11 Fire Management (Program Maintenance)

The Fire Management Plan by Terramatrix (2013) provides recommended works to minimise the risk of fire at Seaford Wetlands. They provide recommendations for:

- Signage to raise awareness of fire risk, suitable egress, and no entry during Code Red fire danger days;
- Continue managing the perimeter vegetation in line with the requirements of an Asset Protection Zone with low fuel loads:
 - Grass will be no more than 50 mm in height;
 - Trees must not over-hang or touch dwellings or assets;
 - Shrubs under trees must be separated by at least 1.5 times their mature height;
 - Tree canopy separation of at least 2 m;
 - tree canopy cover of no more than 15%; and
 - Tree branches pruned to a height of 2 m above ground level.

Four areas of residential land immediately adjoining the reserve were identified as requiring additional fire protection measures to bring them in line with the Asset Protection Zone (Mersey Crescent/Coolibar Avenue, Wilson Grove, Armstrongs Road (east)/Bethune Court, and Bennett Court/Rossiter Court/Greaves Court):

- Classifying and maintaining two areas as Bushfire Management Zones with moderate fuel loads;
- Considering the use of prescribed fire in the Common Reed dominated wetlands (noting that this may not be effective in reducing biomass in the long-term) and the Damp Sands Herb-rich Woodland;
- Undertaking mechanical removal of fuel in Asset Protection Zones or Bushfire Moderation Zones; and
- Improving site access for tankers.

These recommendations should be followed where possible. If determined to be a feasible option, the use of fire to manage #Coast Tea-tree in the Damp Sands Herb-rich Woodland at Seaford Wetlands could be considered.

Using fire to manage Common Reed is not considered viable due to the proximity to houses and the likely outcome of a peat-fire burning for multiple weeks after the above ground fire was extinguished.

9.5 Priority 2 Management Actions (Best Practice Management)

9.5.1 Incorporate 'Downs Estate' into Ramsar Site (Capital/Grants Program)

Frankston City Council's Planning and Environment Department and Melbourne Water are interested in including Downs Estate within the Ramsar boundary (see Sections 2.5 and 4.1.12). Incorporation of Downs Estate into the Ramsar boundary would need to be passed at a Council meeting.

Planning zones and overlays over the property include a Green Wedge Zone, an Urban Floodway Zone, an Environmental Significance Overlay, a Land Subject to Inundation Overlay and a Wildfire Management Overlay. EVCs represented on the property include Aquatic Herbland, Brackish Aquatic Herbland and Tall Marsh (Aspect Studios 2012).

Currently, the water source for Downs Estate comes via flow through the natural drainage line which is supplied from local run-off from rainfall. Melbourne Water is considering pumping water from Wadsleys Drain into the 'old watercourse' natural drainage line which runs southwest across Downs Estate and into the downstream Seaford Wetlands. As the pump delivers c. 15L/min (and there are 1 million cubic metres of storage in the Seaford wetland), the main benefit would be through increasing the habitat in Downs Estate while providing only slightly more water in SN1 (Paul Rees, Melbourne Water, pers. comm.).

Therefore, the periodically filled wetlands have potential to be managed for late-winter and early-spring habitat for waders (e.g. Latham's Snipe). The land immediately adjoining the Seaford Wetlands (to the west of the shared trail) is undoubtedly the most important part of the 'Downs Estate', as it adjoins, is low-lying and subject to flooding and therefore supports complementary values to the Ramsar site. The trail represents a logical boundary between the wetter low-lying area adjoining Seaford and the dryland area to the east of the trail, if the property was to be divided into multiple use zones. For example, the area to the west of the trail could be incorporated into the Ramsar site and the area to the east could be used for other purposes, such as a community centre. A recent Council resolution supported some community activities east of the shared trail (see Section 4.1.12).

The Victorian Waterway Management Strategy provides guidance on extending the boundary of an existing Ramsar site (DEPI 2013b). Under Policy 12.6 of the Strategy, the Victorian Government can recommend extending the boundaries of an existing site to the Australian Government where:

- there is agreement by the owner or land manager and key stakeholders involved in management of the wetland and the actions they propose to meet relevant Ramsar obligations;
- there is compelling evidence that listing will provide clear benefits in:
 - protecting highly significant wetland values relating to the Ramsar criteria for listing;
 - raising the wetland profile; and
 - increasing the level of support for conservation and wise use measures that cannot be achieved through other mechanisms.

Regarding these criteria and incorporation of at least the adjoining parts of 'Downs Estate' into the Ramsar boundary:

- there is agreement between Melbourne Water, Frankston City Council and community groups, such as Bird Life Australia, that ‘Downs Estate’ should be incorporated into the Ramsar site and managed to protect and enhance Ramsar values (e.g. enhance bird values);
- the land would be used to clearly benefit the existing Ramsar site by:
 - protecting and enhancing the ecosystem services that contributed to the Seaford Wetlands Ramsar listing;
 - buffer and protect the Seaford Wetlands, including the values which contributed to Ramsar listing, from future potential incompatible uses of land to the east; and
 - provide additional/complimentary wetland habitat (at Downs Estate) to the habitat within the Ramsar site (i.e. Seaford Wetlands).
- satisfy the principle of ‘wise use’ of a Ramsar wetland (i.e. through provision of additional habitat and habitat to buffer the existing Ramsar values).

Melbourne Water and Frankston City Council should continue to hold discussions and investigate the opportunity for incorporation of Downs Estate into the Ramsar boundary. There is considerable scope for involvement of community groups (e.g. Friends of Edithvale-Seaford Wetlands, Down’s Estate Community Working Group, BirdLife Australia) in the future use and management of this parcel. If the parcel is incorporated into the Ramsar site, a management plan will need to be developed for the site addressing fencing, weed management and revegetation, and controlling activities to the east of the shared trail under a licence issued by Council to minimise any impacts on migratory shorebirds and other birds utilising Downs Estate if water is returned.

9.5.2 Manage High Threat Weeds (Program Maintenance)

Twenty three and twenty one high threat weeds, not including those listed under the *Catchment and Land Protection (CALP) Act 1994*, have been identified as requiring management (eradicate, control, contain) within Edithvale and Seaford Wetlands respectively (Tables 13 and 14). They represent a small proportion of the weed flora, but pose a serious risk to the biodiversity values onsite due to their highly invasive and deleterious impact on the native flora.

At Edithvale and Seaford Wetlands an additional 33 and 41 weeds of management concern have been identified respectively, during the field survey and/or in previous reports for the site (Tables 13 and 14). They are addressed as Priority 1 (Regionally Controlled under the CaLP Act) or Priority 3 (medium threat) issues.

The weeds listed for control here should not be seen as exhaustive as additional weeds of management may be identified onsite in certain situations or as they colonise. An adaptive management process should be undertaken with ongoing monitoring to identify new weeds as they arise.

Additional weed control notes regarding and herbicide use are provided in Appendix 11.

9.5.3 Maintain Dog Restrictions and Increase Signage (capital works)

To minimise impacts of dogs on the wildlife utilising Edithvale-Seaford-Wetland, the current dog restrictions should remain in place. That is, not dogs allowed in the wetland areas, and dogs on-leash only in the buffers. Signage explaining the dog restrictions and educating the public of the impacts of dogs on native wildlife should be installed.

A letter drop to surrounding neighbourhoods within 500 m of either site informing dog owners of domestic dog management issues and potential impacts of pets on biodiversity values within the wetlands should occur in conjunction with the information distributed to cat owners in order to encourage responsible pet ownership (see Section 9.5.4).

9.5.4 Control of Other Pest Animals (Program Maintenance)

It is difficult to control feral and domestic cats from entering Edithvale or Seaford Wetlands due to the largely residential landscape context of the wetlands. Trapping is unlikely to be effective at reducing cat incursions, especially if not undertaken in conjunction with adjoining landowners. Notices to registered cat owners within 500 m of either site regarding cat management issues should be distributed each year as this may contribute towards more responsible cat ownership and a subsequent reduction cat visitation to the wetlands. Friends of Edithvale-Seaford Wetlands could also be supported to hold information evenings regarding responsible pet ownership.

If feral animal issues arise they should be managed accordingly in consultation with a pest control expert.

Control of Mosquitofish at Edithvale-Seaford Wetlands is considered to be unfeasible, primarily because there is likely to be a continuous source for recolonization with inflows from drains feeding the wetlands (Hamer and Parris 2011). Increasing levels of aquatic vegetation may provide shelter for frogs from Mosquitofish, resulting in higher frog survivorship (e.g. Webb and Joss 1997). Options for reducing Mosquitofish abundance may need to be revisited if surveys ever located threatened fish species, such as Dwarf Galaxias at the site.

9.5.5 Undertake Nest Box Monitoring (Program Maintenance)

Nest boxes should be checked and cleaned in May each year. Occupancy of each nest box should be recorded and any pest animals removed (e.g. Common Myna nests, bee hives). Occupancy data should be submitted to Melbourne Water.

9.5.6 Revegetation after Disturbance (Program Maintenance)

Revegetation is proposed for three main purposes:

- To rehabilitate areas after disturbances from activities such as weed removal (e.g. *Spiny Rush), warren/den destruction or grooming;
- To enhance the structural and floristic diversity of Damp Sands Herb-rich Woodland and Swamp Scrub patches; and
- To revegetate an area of currently non-remnant vegetation.

Of these, the rehabilitation after disturbance is considered a Priority 2 management action, while the other two purposes are considered Priority 3 (addressed in Section 9.6.2).

Lists of species suitable for revegetation in various Ecological Vegetation Classes are provided in Appendix 12. For wetland communities, the applicable zone has also been provided based on their tolerance to inundation whereby (Figure 17 illustrates this zonation):

Terrestrial – Dryland vegetation that is primarily outside the natural flooding zone.

Zone 1 – Seasonally wet margins to permanently moist; shallow seasonal inundation in lower part of zone.

Zone 2 – Shallow inundation; upper minimum depth of inundation c. 10 cm; amphibious and emergent aquatic herbs, some straddling Zones 1 and 2.

Zone 3 – Permanent water; submergent and emergent aquatic-herbs, some straddling Zones 2 and 3.

There is considerable scope for involvement of the Friends of Edithvale-Seaford Wetlands in revegetation following disturbance works.

Additional information regarding revegetation is provided in Appendices 12 and 13. The *Edithvale-Seaford Wetlands Revegetation Prescriptions* (TBLA and Australia Ecosystems 2005) document provides more detailed revegetation prescriptions, of which the maps illustrating revegetation locations are provided in Appendix 13.




Figure 17 Conceptual cross section of the embankment for revegetating wetland environments, showing terrestrial vegetation and wetland vegetation zones 1, 2 and 3.

9.5.7 Consolidate Melbourne Water and Frankston City Council Management Boundaries at Seaford Wetland (Capital/Grants Program)

For ease of maintenance and to maintain equal standards, the management boundaries at Seaford Wetland should be consolidated giving ownership of the wetland interior to Melbourne Water and the dryland surrounds to Frankston City Council (see Section 2.5 for additional information).

9.5.8 Investigate Path Extension at Seaford Wetland (Investigation)

There is increasing pressure at Seaford Wetland to use the site for running events and other activities. Due to potential disturbances to wildlife, these should be carefully considered before being permitted. To minimise disturbances along the internal track and to give users an alternative route when this track is closed (e.g. during fox trapping), the shared trail could be extended through Down's Estate to, and along the northern boundary (i.e. on the east side of the proposed fencing shown on Figure 4) to link

with the track on the west side of the wetlands. This would create a complete loop/circuit track that does not require site users to traverse through the wetland, reducing disturbance related impacts.

Disturbance impacts could be further reduced through revegetation designed to screen the wetlands from the users traversing around the wetlands (i.e. on the inside of the shared trail/walking track).

The installation of a loop track with screening revegetation (and fencing) would offset the closure of the internal track to the public and allow visitors to get from one side of the wetland to the other without traversing through the core conservation areas (see Section 9.5.9).

9.5.9 Fencing (Program Maintenance)

Both Edithvale and Seafood Wetlands have extensive fencing to restrict access to the wetlands (Figures 3, 4, 14 and 15). These will need to be maintained should breaches occur.

At Seafood Wetland, extension of fencing should be considered through Downs Estate as indicated in Figure 4. The proposed fence extension would ensure the core conservation areas (wetlands and adjoining areas) are fully enclosed within fences. The core conservation areas could be permanently or seasonally closed to the public, restricting public viewing to designated areas along the boundary to minimise noise and visual disturbance impacts on birdlife. A fully enclosed fence around the conservation areas of the wetland would also facilitate enforcement of dog walkers to areas away from the wetlands. This could be undertaken whether or not Downs Estate gets incorporated into the Ramsar boundary.

Closure of the internal track to the public should be considered to prevent users causing disturbance to waterbirds at Seafood Wetland.

9.5.10 Monitoring (Program Maintenance)

Photo-point monitoring

Photo-points previously established across the site in 2011 (Australian Ecosystems 2011b) should be photographed annually and images stored by Melbourne Water (see Figures 14 and 15). Photographs must be taken at each point in the direction of the documented magnetic bearing. The field of view from baseline photos should be replicated in each photograph. Baseline photographs are provided Australian Ecosystems (2011b).

Monitor pest plant control performance

For the most part, weeds of management concern are generally in low abundance. Targets for pest plant control are identified in Tables 13 and 14 (eliminate, control, contain). A standard monitoring form should be developed to record pest plant control performance for each zone.

Monitor pest animal control performance

The site should be inspected in early spring each year to determine whether there are any active rabbit warrens or fox dens. This should be undertaken more regularly if warrens or dens are treated to control pest animals. BirdLife Australia should continue to record the locations of foxes and cats whenever seen during monthly bird surveys.

Monitor Eastern Grey Kangaroo population

The kangaroo management undertaken in March 2016 by VWSA (2016) has reduced the kangaroo population to the recommended sustainable level. Annual monitoring in March/April (when the wetlands are dry) should be undertaken (see EcoPlan Australia 2015) to ensure that the population remains within this stated sustainable range of 16-27 kangaroos. This should use the drive count method as per previous years, to ensure that the counts are as repeatable as possible.

9.5.11 Monitoring (Biodiversity Monitoring and Assessment)

Native vegetation condition monitoring

As part of the *Waterways Sites of Biodiversity Significance Plan 2013*, Melbourne Water intends to monitor the condition of remnant vegetation across each SoBS every five years using the Vegetation Quality Assessment (Habitat Hectares) method. Baseline data were collected in 2010 and then updated in 2015 (Australian Ecosystems 2011a, 2016). The 2010 and 2015 assessments variously used the Vegetation Quality Assessment (VQA) and Index of Wetland Condition (IWC) assessment method making it difficult to compare the condition of vegetation over the five year period (see Section 6.1). Though not designed as a monitoring assessment method, future surveys should follow the VQA methodology as it provides a more detailed assessment of vegetation quality than the IWC assessment method. As the VQA method was not designed to be used as a monitoring tool, the details required to score each component (observed percent cover, diversity and cover of each understorey lifeform, etc.) should also be documented for each assessment component.

The Vegetation Quality Assessments should include an inventory of the entire flora for each Habitat Zone or remnant vegetation within the reserve. Resulting data should be added to the Victorian Biodiversity Atlas.

Where possible, the boundaries of each zone should remain consistent with previous years. However, it is expected that some variations will occur as the Common Reed management program proceeds. This has in part been the reason for discrepancies in the EVC and Habitat Zone mapping in 2010 and 2015 (Australian Ecosystems 2011, 2015).

9.6 Priority 3 Management Actions (Other Management Priorities)

9.6.1 Manage Medium Threat Weeds (Program Maintenance)

Tables 13 and 14 list 21 and 26 weed species listed as 'medium threat' at Edithvale and Seaford Wetlands, respectively. These species require management within the sites, though precedence must be given to managing CaLP Act listed 'regionally controlled' noxious weeds and 'high threat' weeds. No medium threat weed species listed in Tables 13 and 14 should be allowed to increase in cover, particularly in areas of groomed Common Reed.

9.6.2 Revegetation and Supplementary Plantings in Woodland, Scrub and Non-Remnant Vegetation (Capital/Grants Program)

To increase the structural and floristic diversity of the Damp Sands Herb-rich Woodland vegetation that primarily comprises canopy trees over mown lawn, clustered plantings of understorey shrubs and robust

ground layer species should be undertaken. Supplementary plantings could also occur in better quality Damp Sands Herb-rich Woodland remnants, particularly after the removal of woody weeds such as #Coast Tea-tree at Seaford Wetlands.

Similarly areas of Swamp Scrub could be expanded or the floristic diversity increased with supplementary plantings. At Seaford Wetlands, a small area of non-remnant vegetation still comprising scattered indigenous wetland and shrub species located opposite Seaford North Primary School should also be revegetated with Swamp Scrub vegetation.

A list of species suitable for revegetation in various Ecological Vegetation Classes is provided in Appendix 12. Additional information regarding revegetation is provided in Appendix 13. The *Edithvale-Seaford Wetlands Revegetation Prescriptions* (TBLA and Australia Ecosystems 2005) document provides more detailed revegetation prescriptions, of which the maps illustrating revegetation locations are provided in Appendix 13.

9.6.3 Manage User-related Issues: New Tracks and Disturbances (Program Maintenance)

Inspections for new tracks or disturbances such as bike tracks and cubby huts should be undertaken during regular maintenance visit. If new tracks or disturbances are located they should be dismantled and the disturbed area rehabilitated as appropriate (e.g. weed control, revegetation). Any resultant weed infestation should be addressed. For example, a bike track with jumps was observed in the eastern most Damp Sands Herb-rich Woodland at Seaford Wetlands, with a resultant African Love-grass *Eragrostis curvula* infestation that requires management.

9.6.4 Manage Lower Priority User Related Issues (Program Maintenance)

General litter is primarily in low abundance around the site. Given the sites close proximity to the sea, litter collection should be undertaken as part of the regular ongoing maintenance of the site. Litter traps at stormwater inlets and outlets should also be regularly cleared, particularly after moderate to large rain events.

9.6.5 Monitoring (Biodiversity Monitoring and Assessment)

Significant Flora Monitoring

No extensive floristic lists are available for the site (TBLA and Australian Ecosystems 2005; Australian Ecosystems 2011a, 2016; DELWP 2016a) or distribution mapping of rare or threatened flora. For this reason, an extensive flora survey is recommended to document the occurrence of rare or threatened (and ideally also regionally significant species) within the site. This should be completed in year one of the management plan, and revisited every five years. This will allow managers to better understand the ecological values of the reserve and manage them accordingly.

The survey should be undertaken in spring by systematically traversing the site, focusing on areas of suitable habitat for rare or threatened flora species identified as potentially utilising the site (Tables 15 and 16 in Appendix 2). The resultant species list should be entered into the Victorian Biodiversity Atlas (DELWP 2016a) and a report prepared including management recommendations to enhance the habitat for the significant flora species found.

Targeted Surveys for Significant Fauna Species

Very little information is available on the reptile and fish species of the Edithvale-Seaford Wetlands (Section 4.1.6).

The wetlands provide potential habitat for the Swamp Skink. Melbourne Water may wish to determine the status of the Swamp Skink at the wetlands as well as the value of the wetlands in general for reptile species by undertaking a targeted survey for this group.

Dwarf Galaxias occurs in slow-flowing shallow ephemeral or permanent freshwater wetlands, such as swamps and drains (DELWP 2015), and has been recorded in other waterways that flow into the Patterson River and in catchments to the south of the site (Coleman et al. 2015). Although it seems unlikely that the Dwarf Galaxias occurs at Edithvale-Seaford Wetlands due to the isolation of the wetlands from their original catchments, a fish survey would provide useful information for the site, as both freshwater and saline water flows enter the wetlands from drains, potentially delivering both freshwater and estuarine inhabiting fish species to the wetlands.

Invertebrate Monitoring

As very little is known about the terrestrial and aquatic invertebrates at the wetlands, it is desirable to undertake an invertebrate inventory survey, particularly for aquatic species. An invertebrate survey would provide useful information for the site, including information or indices relating to the condition of the wetlands.

9.7 Implications for Cultural and Heritage Values

As per the findings of the desktop cultural heritage assessments by Heritage Insight (2016a, b), a Cultural Heritage Management Plan (CHMP) is required for an activity if it:

- occurs within an area of cultural heritage sensitivity (such as a declared Ramsar wetland); and
- is considered a high impact activity (see Heritage Insight 2016a, b for a list of high impact activities).

Although a CHMP is not required for any of the works recommended in this management plan, soils disturbance should be avoided wherever possible. If significant soil disturbance is proposed, the potential impacts and need for a CHMP should be evaluated on a case-by-case basis. A field survey of the study area would be able to provide a more detailed archaeological assessment and identify areas of specific archaeological potential along with any surface Aboriginal sites within the study area. This information could then be used to determine whether mandatory or voluntary CHMP preparation is required for specific proposed works. This would be particularly useful for any activities that do not trigger a mandatory CHMP and could prevent accidental harm to unrecorded Aboriginal cultural heritage material.

10 References

- ALS (2010) Melbourne Water Mosquito Monitoring Risk Assessment. Report No. 054-2010-066 prepared by ALS Water Resources Group for Melbourne Water. (ALS Water Resources Group)
- ANCA (1996) A Directory of Important Wetlands in Australia (second edition) (Australian Nature Conservation Agency, Canberra)
- Antos MJ, Ehmke GC, Tzaros CL and Weston M (2007) Unauthorised human use of an urban coastal wetlands sanctuary: Current and future patterns. *Landscape and Urban Planning* **80** 173–183.
- ANZECC and ARMCANZ (2000) Australian and New Zealand guidelines for fresh and marine water quality. Volume 1, The guidelines. Australian and New Zealand Environment and Conservation Council, Agriculture and Resource Management Council of Australia and New Zealand.
- Aspect Studios (2012) Downs Estate Masterplan. Unpublished report prepared in association with Practical Ecology for Frankston City Council (Aspect Studios, Melbourne)
- Aurecon (2011) Stage 1 Interim Report Weatherstone Road and Bardia Avenue Drainage Pumping Stations Melbourne Water. Unpublished report prepared for Melbourne Water (Aurecon Australia Pty Ltd, South Melbourne)
- Australian Ecosystems (2011a) Melbourne Water Sites of Biodiversity Significance Vegetation Assessments Autumn 2010. Unpublished report prepared for Melbourne Water. (Australian Ecosystems: Port Melbourne)
- Australian Ecosystems (2011b) Edithvale-Seaford Wetlands: Mapping the extent and distribution of over-abundant native and exotic species. Unpublished report prepared for Melbourne Water. (Australian Ecosystems: Port Melbourne)
- Australian Ecosystems (2016) Melbourne Water Sites of Biodiversity Significance Vegetation Assessments 2014-2015. Unpublished report prepared for Melbourne Water. (Australian Ecosystems Pty Ltd: Port Melbourne)
- Australian Government (2000) Australian Ramsar Management Principles, available for download at <http://www.environment.gov.au/water/wetlands/managing/australian-ramsar-management-principles>
- Bachmann, M (2009) *Acacia longifolia* in the South East region – Unravelling the myths and working towards the adoption of sensible “Guidelines for Management”. Report to Accompany Field Trip on Friday 27th March [2009]. Government of South Australia, Department of Environment and Heritage
- Birds Australia (2004) Bird Habitat Values at the Edithvale and Seaford Wetlands. Unpublished report prepared by C Tzaros and A Silcocks for Melbourne Water Corporation (Birds Australia, East Hawthorn)

- Birds Australia (2006) Edithvale and Seaford Wetlands Bird Survey Project 2003-06: Final Report 2003-2006, Results for 2005-2006, Management Recommendations, and Trend Analysis. Unpublished report prepared by A Silcocks, G Ehmke, C Tzaros and M Weston for Melbourne Water (Birds Australia, Carlton)
- Birds Australia (2007) Edithvale and Seaford Wetlands Bird Survey Project 2006-09: Report No. 1 May 2006 to April 2007. Unpublished report prepared by A Silcocks and J O'Connor for Melbourne Water (Birds Australia, Carlton)
- Birds Australia (2008) Edithvale and Seaford Wetlands Bird Survey Project 2006-09: Report No. 1 May 2007 to April 2008. Unpublished report prepared by A Silcocks and J O'Connor for Melbourne Water (Birds Australia, Carlton)
- Birds Australia (2010) Edithvale and Seaford Wetlands Bird Survey Project 2009-10: May 2009 to June 2010. Unpublished report prepared by A Silcocks and J O'Connor for Melbourne Water (Birds Australia, Carlton)
- Birds Australia (2011) Edithvale and Seaford Wetlands Bird Survey Project 2010-11: July 2010 to June 2011. Unpublished report prepared by A Silcocks and J O'Connor for Melbourne Water (Birds Australia, Carlton)
- BirdLife Australia (2013a) Edithvale and Seaford Wetlands Bird Survey Project 2011-12: July 2011 to June 2012. Unpublished report prepared by A Silcocks for Melbourne Water (BirdLife Australia, Carlton)
- BirdLife Australia (2013b) Edithvale and Seaford Wetlands Bird Survey Project 2012-13: July 2012 to June 2013. Unpublished report prepared by A Silcocks for Melbourne Water (BirdLife Australia, Carlton)
- BirdLife Australia (2015a) Edithvale and Seaford Wetlands Bird Survey Project 2013-14: July 2013 to June 2014. Unpublished report prepared by A Silcocks for Melbourne Water (BirdLife Australia, Carlton)
- BirdLife Australia (2015b) Edithvale Banding Project. Unpublished report prepared by A Silcocks for Melbourne Water (BirdLife Australia, Carlton)
- BirdLife Australia (2016a) Edithvale and Seaford Wetlands Bird Survey Project 2014-15: July 2014 to June 2015. Unpublished report prepared by A Silcocks for Melbourne Water (BirdLife Australia, Carlton)
- BirdLife Australia (2016b) Australian IBAs. Web page available at:
<http://birdlife.org.au/projects/important-bird-areas/iba-maps>
- Brett Lane and Associates (2001) Information Sheet on Ramsar Wetlands (for Edithvale-Seaford Wetlands). (Brett Lane and Associates, Burnley) available at:
<http://www.environment.gov.au/water/topics/wetlands/database/pubs/57-ris.pdf>
- Burrell JP (1981) Invasion of Coastal Heaths of Victoria by *Leptospermum laevigatum*. *Australian Journal of Botany*, 29: 747-764.
- Carter M (1975) Conservation and Management Proposals for Edithvale – Carrum – Seaford Wetlands Areas. Unpublished report prepared by the Wetlands Investigation Committee, Frankston.

- Chatterjee A, Phillips B and Stroud D (2008) Wetland Management Planning: A Guide for Site Managers (WWF, Wetlands International, IUCN and Ramsar Bureau)
- Coleman RA, Hoffmann AA and Raadik TA (2015) A review of *Galaxiella pusilla* (Mack) (Teleostei: Galaxiidae) in south-eastern Australia with a description of a new species *Zootaxa* **4021**(2), 243–281.
- Convention on Wetlands (1999) Wetland Risk Assessment Framework, Resolution 7.10 of the Seventh Meeting of the Contracting Parties to the Convention on Wetlands, Costa Rica 1999. (Secretariat of the Ramsar Convention: Gland, Switzerland)
- Court A.B (2001) *Acacia longifolia*. Flora of Australia Vol 11B Mimosaceae, *Acacia* part 2. CSIRO Publishing.
- Curtis WM (1956) The Student's Flora of Tasmania Part 1. Gymnospermae, Angiospermae. LG Shea, Government Printer, Tasmania
- DEDJTR (2016) Directions for the Use of 1080 and PAPP Pest Animal Bait Products in Victoria. (Department of Economic Development, Jobs, Transport and Resources, Melbourne)
Available at: http://agriculture.vic.gov.au/_data/assets/pdf_file/0003/263541/Directions-for-the-use-of-1080-and-PAPP-pest-animal-bait-products-in-Victoria.pdf
- DELWP (2015) Dwarf Galaxias *Galaxiella pusilla* Action Statement No. 258 *Flora and Fauna Guarantee Act 1988*. (Department of Environment, Land, Water and Planning: East Melbourne)
- DELWP (2016a) Victorian Biodiversity Atlas Version 3.0.6 database. Available at <http://www.depi.vic.gov.au/environment-and-wildlife/biodiversity/victorian-biodiversity-atlas> [Accessed 7 March 2016].
- DELWP (2016b) Biodiversity Interactive Map 3.2. Available at: <http://mapshare2.dse.vic.gov.au/MapShare2EXT/imf.jsp?site=bim> [Accessed 11 April 2016]
- DELWP (2016c) Planning Schemes Online. Available at <http://planningschemes.dpcd.vic.gov.au/> [Accessed May 2016]
- DELWP (2016d) Planning Schemes Online. Available at: <http://services.land.vic.gov.au/maps/pmo.jsp>
- DELWP (2016e) Victorian Ramsar Rolling Review 2015-16. Summary report prepared by J Hale for the Department of Environment, Land, Water and Planning (J Hale, Wandong)
- DEPI (2004) Vegetation Quality Assessment Manual – Guidelines for Applying the Habitat Hectares Scoring Method, Version 1.3. (Department of Sustainability and Environment: East Melbourne)
- DEPI (2013a) Edithvale-Seaford Wetlands Ramsar Site Boundary Description Technical report (Department of Environment and Primary Industries, East Melbourne)
http://www.depi.vic.gov.au/_data/assets/pdf_file/0004/242761/FINAL_Edithvale_Seaford_Wetlands_Boundary_Description_Dec2013.pdf

- DEPI (2013b) Improving Our Waterways Victorian Waterways Management Strategy (Department of Environment and Primary Industries, East Melbourne)(Section 12) available at:
http://www.depi.vic.gov.au/_data/assets/pdf_file/0017/200375/VWMS_Part3.pdf
- DEPI (2014) Advisory list of rare or threatened plants in Victoria – 2014. (Department of Sustainability and Environment: East Melbourne)
- DoE (2016) EPBC Act Protected Matters Search Tool. Available at:
<http://www.environment.gov.au/epbc/protected-matters-search-tool> [Accessed 5 April 2016]
- DEWHA (2008) National Framework and Guidance For Describing the Ecological Character of Australian Ramsar Wetlands Module 2 of the National Guidelines for Ramsar Wetlands — Implementing the Ramsar Convention in Australia Department of the Environment, Water, Heritage and the Arts , Canberra)
- DEWHA (2009) EPBC Act Policy Statement 3.21 Draft Significant Impact Assessment Guidelines for 36 Migratory Shorebird species (Department of Environment, Heritage, Water and the Arts, Canberra)
- DNRE (2002) Management of Victoria’s Ramsar Wetlands Strategic Directions Statement. (Department of Natural Resources and Environment: East Melbourne)
- Dowling B. and Weston M.A. (1999). Managing a breeding population of the Hooded Plover *Thinornis rubricollis* in a high-use recreational environment. *Bird Conservation International* 9, 255–270.
- DSE (2004) Sites of Zoological Significance of South East Melbourne and the Mornington Peninsula – a compendium of information collected between 1987 and 1991. Arthur Rylah Institute for Environmental Research Technical Report Series No. 92. Unpublished report prepared by R Brereton, M Schulz, I Mansergh, K Sandiford and S Bennett (Department of Sustainability and Environment, Arthur Rylah Institute for Environmental Research, Heidelberg)
- DSE (2004) Framework for Mosquito Management in Victoria (Department of Sustainability and Environment, East Melbourne). Available at:
<https://www2.health.vic.gov.au/about/publications/researchandreports/Framework%20for%20Mosquito%20Management%20in%20Victoria%20-%20DSE%202004> Accessed 29 August 2016.
- DSE (2006) Ministerial Guidelines for Assessment of Environmental Effects under the Environment Effects Act 1978. Seventh edition (Department of Sustainability and Environment, Melbourne) available at:
http://www.dtpli.vic.gov.au/_data/assets/pdf_file/0010/230599/DSE097_EES_FA.pdf
- DSE (2009a) Advisory list of threatened invertebrate fauna in Victoria – 2009. (Department of Sustainability and Environment: East Melbourne)
- DSE (2009b) A field guide to Victorian Wetland Ecological Vegetation Classes for the Index of Wetland Condition. (Department of Sustainability and Environment: East Melbourne).

- DSE (2009c) Victorian Coastal Acid Sulfate Soils Strategy (Department of Sustainability and Environment, East Melbourne)
- DSE (2010) Victorian Best Practice Guidelines for Assessing and Managing Coastal Acid Sulfate Soils (Department of Sustainability and Environment, Melbourne)
- DSE (2011) Western Port Welcomes Waterbirds: Waterbird Usage of Western Port. Technical Report Series No. 222. Report prepared by B Hansen, P Menkhorst and R Loyn (Arthur Rylah Institute for Environmental Research, Department of Sustainability and Environment, Heidelberg)
- DSE (2012) Description of the Ecological Character Edithvale-Seaford Wetlands Ramsar Site (Department of Sustainability and Environment, Melbourne)
- DSE (2013) Advisory list of threatened vertebrate fauna in Victoria – 2013 (Department of Sustainability and Environment, Melbourne)
- Ecology Australia (1987) The vegetation on public land in the Balcombe Creek catchment, Mornington Peninsula. Unpublished report by ARG McMahon, GW Carr, SE Bedggood for the Shire of Mornington.
- Ecology Australia (2000) Edithvale–Seaford Wetlands Ramsar Management Plan. Unpublished report prepared by BA Lane, MR Bezuijen, CK Orscheg, JA Todd and GW Carr for Melbourne Water (Ecology Australia Pty Ltd, Fairfield)
- Ecology Australia (2005c) Unimin Australia Limited, Lang Lang – Flora and Fauna Assessment. Unpublished report prepared for Unimin Australia by ED Moysey, C Campbell, N Roberts, C Wilson, D Quin and A McMahon (Ecology Australia, Fairfield).\
- Ecology Australia (2013) Sites of Biodiversity Significance Management Strategy – Background Document 2013. Unpublished report prepared by C Renowden, ARG McMahon and DG Quin for Melbourne Water Corporation (Ecology Australia Pty Ltd, Fairfield)
- Ecology Partners (2007) Distribution and Conservation Significance of Frog Species on the Mornington Peninsula, Victoria. Unpublished report prepared by A Aboltins and A Organ for Melbourne Water (Ecology Partners, Brunswick)
- Ecoplan Australia (2015) Eastern Grey Kangaroo Management Plan for Edithvale South Wetland. Unpublished report prepared by E Hynes for Melbourne Water (Ecoplan Australia Pty Ltd, Hurstbridge)
- Eco-tainable (2013) Acid Sulfate Soil Assessment Seaford Wetland, Seaford, Victoria. Unpublished report prepared by P O’Neill for Melbourne Water (Eco-tainable, Moorabbin)
- Entwisle TJ, Maslin BR, Cowam RS and Court AB (1996) ‘Mimosaceae’ In Walsh, N.G and Entwisle TJ eds., ‘Flora of Victoria’ Vol 3, pp. 585-656. Inkata Press: Melbourne.
- Garrards (2016) ProLink™ XR Briquets Mosquito Growth Regulator. Available at: http://www.garrards.com.au/images/stories/zone_files/labels/l_prolink_briquets.pdf
- GHD (2015) Melbourne Water Mosquito Monitoring Seaford 2014-2015. Unpublished report prepared by L Bateson for Melbourne Water (GHD Pty Ltd Melbourne)

- GHD and Aquatic Systems Management Pty Ltd (2006) Edithvale-Seaford Wetlands Review of Wetland Operations. Unpublished report prepared by G Corda for Melbourne Water (GHD Pty Ltd Melbourne)
- Glover HK, Weston MA, Maguire GS, Miller KK and Christie BA (2011) Towards ecologically meaningful and socially acceptable buffers: Response distances of shorebirds in Victoria, Australia, to human disturbance. *Landscape and Urban Planning* **103**, 326–334.
- Greet J (2015) Seaford Wetlands Vegetation Monitoring Program: report on *Phragmites* expansion and control. Technical 15.4. Melbourne Waterway Research Practice Partnership.
- Greet J and Rees P (2015) Slashing may have potential for controlling *Phragmites australis* in long-inundated parts of a Ramsar-listed wetland. *Ecological Management Restoration* **16** (3), 233–236.
- Hale J (2016) Ecological Character Description for the Edithvale-Seaford Wetlands (Addendum) and Ramsar Information Sheet Review. Unpublished report prepared by J Hale for the Department of Environment, Land, Water and Planning (J Hale, Wandong)
- Hamer AJ, Parris KM (2011) Local and landscape determinants of amphibian communities in urban ponds. *Ecological Applications* **21**(2), 378–390.
- Heritage Insight (2016a) Edithvale-Seaford Ramsar Wetlands Edithvale Section Desktop Cultural Heritage Assessment. Unpublished report prepared by R McAlister and D Di Fazio for Melbourne Water (Heritage Insight, Kensington)
- Heritage Insight (2016b) Edithvale-Seaford Ramsar Wetlands Seaford Section Desktop Cultural Heritage Assessment. Unpublished report prepared by R McAlister and D Di Fazio for Melbourne Water (Heritage Insight, Kensington)
- Higgins PJ and Davies SJJF (Eds) (1996) Handbook of Australian, New Zealand and Antarctic Birds. Volume 3 Snipe to Pigeons (Oxford University Press, Melbourne)
- Hua N, Tan K, Chen Y, Ma Z (2015) Key research issues concerning the conservation of migratory shorebirds in the Yellow Sea region. *Bird Conservation International* 25:38–52.
- IUCN (2008). International Union for Conservation of Nature and Natural Resources Red List of Threatened Species. <http://www.iucnredlist.org>
- Jacobs (2016a) Seaford Wetlands Water Quality Analysis. Unpublished report prepared by B Bonneville for Melbourne Water [Jacobs Group (Australia) Pty Limited, Melbourne]
- Jacobs (2016b) Coastal Wetlands and Climate Change Decision Support Framework. Unpublished report prepared for Melbourne Water [Jacobs Group (Australia) Pty Limited, Melbourne]
- KBR (2009) Edithvale-Seaford Ramsar Site Management Plan. Unpublished report prepared by S Heron for Melbourne Water (Kellogg, Brown and Root, Melbourne)
- Kennedy S, Wheeler J and Foley E (2012) 44 First Avenue, Chelsea Heights. (Australian Cultural Heritage Management (ACHM): Brunswick).
- Kingsford RT, Porter JL and Halse SA (2012) National Waterbird Assessment. Unpublished report prepared for the National Water Commission (Australian Wetlands, Rivers, and Landscape Centre, University of New South Wales)

- Komak S and Crossland MR (2000) An assessment of the introduced mosquitofish (*Gambusia affinis holbrooki*) as a predator of eggs, hatchlings and tadpoles of native and non-native anurans. *Wildlife Research* **27**, 185–189.
- MacKinnon J, Verkuil Y and Murray N (2012) IUCN situation analysis on East and Southeast Asian intertidal habitats, with particular reference to the Yellow Sea (including the Bohai Sea). Occasional Paper of the IUCN Species Survival Commission No. 47. IUCN, Gland, Switzerland and Cambridge, UK. li + 70pp. Also available at: www.iucn.org/asiancoastalwetlands
- McLeod EM, Guay PJ, Tayson AJ, Robinson RW and Weston MA (2013). Buses, cards, bicycles and walkers: the influence of the type of human transport on the flight responses of waterbirds. *PLoS One* **8(12)**:e82008. Doi:10.1371/journal.pone.0082008.
- Maguire GS (2008) A Practical Guide for Managing Beach-nesting Birds. Unpublished report prepared for the Australian Government (Birds Australia, Carlton)
- Marchant S and Higgins PJ (1990) Handbook of Australian, New Zealand and Antarctic Birds. Volume 1 Ratites to Ducks (Oxford University Press: Melbourne)
- Martín B, Delgado S, de la Cruz A, Tirado S and Ferrer M (2014) Effects of human presence on the long-term trends of migrant and resident shorebirds: Evidence of local population declines. *Animal Conservation* **18**, 73-81 (Print ISSN 1367-9430)
- Melbourne Water (undated^a) Melbourne Water Mosquito Management Decision Support Tool. Report prepared by Melbourne Water.
- Melbourne Water (undated^b) Melbourne Water Mosquito Management Sampling and Abundance Guidance. Report prepared by Melbourne Water.
- Melbourne Water (2007) Carrum Lowlands Fire Protection Plan. Prepared by Melbourne Water Infrastructure Group.
- Melbourne Water (2013a) Healthy Waterways Strategy, November 2013. (Melbourne Water Corporation: Docklands)
- Melbourne Water (2013b) Waterways Sites of Biodiversity Significance Plan 2013 – October 2013. (Melbourne Water Corporation: Docklands)
- Melbourne Water (2013c) Melbourne Water Provenance Policy (Melbourne Water Corporation: Docklands)
- Melbourne Water (2014a) Melbourne Water Standard for Plant Installation. (Melbourne Water Corporation: Docklands)
- Melbourne Water (2014b) Melbourne Water Standard for Plant Supply (Melbourne Water Corporation: Docklands)
- Melbourne Water (2016) Mosquito monitoring and control at Seaford Wetlands. Available at: <http://www.melbournewater.com.au/whatwedo/projectsaroundmelbourne/pages/mosquito-monitoring-and-control-at-seaford-wetlands.aspx> Accessed 29 August 2016.
- Menkhorst P (2012) The food and foraging rate of an Australasian Bittern. *Australian Field Ornithology* **29**, 133–142.

- Molnar, C. D., Fletcher, D. and Parsons, R. F. (1989) Relationships between heath and *Leptospermum laevigatum* scrub at Sandringham, Victoria. *Proceedings of the Royal Society of Victoria* 101: 77-87.
- Morgan LA, Buttemer MA (1996) Predation by the non-native fish *Gambusia holbrooki* on small *Litoria aurea* and *L. dentata* tadpoles. *Australian Zoologist* **30** (2), 143–149.
- Morrison DA and Davies SJ (1991) *Acacia*. In: GJ Harden ed. 'Flora of New South Wales' Vol. 2 New South Wales University Press, Sydney
- Murray NJ, Ma Z, Fuller RA (2015) Tidal flats of the Yellow Sea: a review of ecosystem status and anthropogenic threats. *Austral Ecology* 40 (4), 272–281. Doi: 10.1111/aec.12211
- NWPASS (2000) National Strategy for the Management of Coastal Acid Sulfate Soils. Report prepared by National Working Party on Acid Sulfate Soils for NSW Agriculture Wollongbar Agricultural Institute Wollongbar)
- <https://www.environment.gov.au/system/files/resources/f682fc6e-4b46-4f4f-abab-f42857489136/files/natass.pdf>
- Pacific Biologics (2016) ProLink Mosquito Growth Regulator. Available at: <http://www.pacificbiologics.com/index.php/larvicides/prolink> Accessed 6 September 2016
- Parsons Brinckerhoff (2009) Preliminary Bat Surveys Eastern Treatment Plant, Edithvale-Seaford Wetlands, Wannarkladdin Wetlands, Boundary Road Wetland and PARCS Wetland. Unpublished report prepared by Rob Gratton and Chad Browning for Melbourne Water (Parsons Brinckerhoff, Melbourne)
- Parsons Brinckerhoff (2010) Level 2 Targeted Microbat Survey Eastern Treatment Plant and Edithvale-Seaford Wetlands Unpublished report prepared by Rob Gratton and Chad Browning for Melbourne Water (Parsons Brinckerhoff, Melbourne)
- Paton D, Ziembicki M, Owen P and Heddle C (2000) Distances for Waterbirds and the Management of Human Recreation with Specific Reference to the Coorong Region of South Australia (University of Adelaide, South Australia)
- Pedley L, Henderson RJF and Reynolds ST (1983) *Acacia* in In: TD Stanley and EM Ross eds: 'Flora of south-eastern Queensland' Vol 1, pp 336-383 Queensland Department of Primary Industries Miscellaneous Publication 81020
- Phillips B (2006) Critique of the Framework for Describing the Ecological Character of Ramsar Wetlands (Department of Sustainability and Environment, Victoria, 2005) based on its application at three Ramsar sites: Ashmore Reed National Nature Reserve, the Coral Sea Reserves (Coringa-Herald and Lihou Reeds and Cays), and Elizabeth and Middleton Reefs Marine National Nature Reserve (Mainstream Environmental Consulting, Waramanga ACT)
- Pyke GH (2005) A review of the biology of *Gambusia affinis* and *G. holbrooki*. *Reviews in Fish Biology and Fisheries* **15**, 339–365.
- Pyke GH, White AW (1996) Habitat requirements for the Green and Golden Bell Frog *Litoria aurea* (Anura: Hylidae). *Australian Zoologist* **30**, 224-232.

- Pyke GH, White AW (2000) Factors influencing predation on eggs and tadpoles of the endangered Green and Golden Bell Frog *Litoria aurea* by the introduced Plague Minnow *Gambusia holbrooki*. *Australian Zoologist* **31**(3), 496–505.
- Ramsar Contracting Parties (2002) New Guidelines for Management Planning for Ramsar Sites and Other Wetlands. Adopted by Resolution VIII.14 (2002) of the Ramsar Convention (Valencia, Spain, available for download at –
<http://www.ramsar.org/sites/default/files/documents/library/new-mgt-guide.pdf>;
- Ramsar Convention (2005) Resolution IX.1 Annex A. A Conceptual Framework for the wise use of wetlands and the maintenance of their ecological character. Web page available at:
http://www.ramsar.org/sites/default/files/documents/pdf/res/key_res_ix_01_annexa_e.pdf
- Ramsar Convention (2016) The Ramsar Convention and its Mission. Web page available at:
<http://www.ramsar.org/about/the-ramsar-convention-and-its-mission> (accessed 30 May 2016)
- RAOU (2001) Edithvale Wetlands Bird Census Project September 1989 – August 1999. Unpublished report prepared by I Endersby and M Carter for Melbourne Water (Royal Australasian Ornithologists Union, Hawthorn East)
- SKM (2011) Environmental water requirements and associated capital works for the Edithvale-Seaford Wetlands. Unpublished report prepared by B Bonneville, P Boon, D Quin, S Treadwell, R Middleton and R Morden for Melbourne Water (Sinclair Knight Merz, Melbourne)
- Taylor (2006) Managing visitor disturbance of waterbirds on Australian inland wetlands. Pp. 150–157. In: ‘Wetlands of the Murrumbidgee River Catchment: Practical Management in an Altered Environment’. Eds. I. Taylor, P. Murray and S. Taylor. Fivebough and Tuckerbil Wetlands Trust (Active Print, Wagga, Wagga, New South Wales)
- TBLD and Australian Ecosystems (2005) Edithvale-Seaford Wetlands Revegetation Prescriptions Unpublished report prepared for Melbourne Water and Frankston City Council. (Thompson Berrill Landscape Design Pty Ltd: Prahran)
- Terramatrix (2013a) Seaford Wetlands Fire Management Plan. Unpublished report prepared by M Hansby, A Tibbits and J Boura for Frankston City Council (Terramatrix Pty Ltd, Collingwood)
- Terramatrix (2013b) Frankston City Council Bushfire Management Strategy. Unpublished report prepared for Frankston City Council (Terramatrix Pty Ltd, Collingwood)
- Thomas L and Middleton J (2003) Guidelines for Management Planning of Protected Areas (IUCN Gland, Switzerland and Cambridge, UK) available for download at
http://cmsdata.iucn.org/downloads/pag_010.pdf
- Thompson, J (1989) A revision of the genus *Leptospermum* (Myrtaceae) *Telopea* 3(3): 301 – 449
- USEPA (1998) Guidelines for Ecological Risk Assessment (US Environment Protection Authority)

- VWSA (2016) Edithvale Kangaroos Management April 18th-22nd 2016 Veterinary Report. Unpublished report prepared by K Watter for Melbourne Water (Veterinary Wildlife Services Australia Pty Ltd, Maleny, Queensland)
- VicFlora (2015) Flora of Victoria, Royal Botanic Gardens Victoria. Available at: <http://data.rbg.vic.gov.au/vicflora>. Accessed April 2016.
- Victorian Coastal Council (2014) Victorian Coastal Strategy 2014. Fourth iteration (Victorian Coastal Council, East Melbourne)
- Watkins D (1993) A National Plan for Shorebird Conservation in Australia. Australasian Wader Studies Group (Royal Australasian Ornithologists Union, Moonee Ponds)
- Webb C and Joss J (1997) Does predation by the fish *Gambusia holbrooki* (Atheriniformes: Poeciliidae) contribute to declining frog populations? *Australian Zoologist* **30**, 316–323.
- Weston MA, Antos MJ and Glover HK (2009) Birds, buffers and bicycles: a review and case study of wetland buffers. *The Victorian Naturalist* **126** (3), 79–86.
- Weston MA, McLeod EM, Blumstein DT and Guay P-J (2012) A review of flight-initiation distances and their application to managing disturbance to Australian birds. *Emu* **112**, 269–286.
- Weston MA (2003) Managing the Hooded Plover in Victoria – a review of existing information. Parks Victoria Technical Series No. 4 (Parks Victoria: Melbourne)
- Weston MA and Elgar MA (2005) Disturbance to brood-rearing Hooded Plovers *Thinornis rubricollis*: responses and consequences. *Bird Conservation International* **15**, 193209.
- Weston MA and Elgar MA (2007) Responses of incubating Hooded Plovers (*Thinornis rubricollis*) to disturbance. *Journal of Coastal Research* **23**, 569–576.
- Whibley DJE and Symon DE (1992) 'Acacias of South Australia'. The Flora and Fauna of South Australia Handbooks Committee, Adelaide.
- Wildlife Profiles (1999) Sunshine Reserve and Fairbairn Park Reptile and Amphibian Study. Report of a Preliminary Field Evaluation of the Habitat for Reptiles and Amphibians at Sunshine Reserve and Fairbairn Park. Unpublished report prepared by P Robertson for the Mornington Peninsula Shire Council (Wildlife Profiles, Heidelberg)

11 Glossary

Acronym	Synonym
AMIS	Asset Management Information System
CaLP Act	Victorian <i>Catchment and Land Protection Act 1994</i>
CWP	Capital Works Program
EPBC Act	Commonwealth <i>Environment Protection and Biodiversity Conservation Act 1999</i>
EVC	Ecological Vegetation Class
FFG Act	Victorian <i>Flora and Fauna Guarantee Act 1988</i>
PM	Program Maintenance
SoBS	Sites of Biodiversity Significance
VROTS	Victorian Rare or Threatened Species
WONS	Weeds of National Significance

Term	Definition
Biodiversity	The variety of all life-forms, the different plants, animals and micro-organisms, the genes they contain, and the ecosystems of which they form a part.
Bioregion	A landscape based approach to classifying the land surface using a range of environmental attributes such as climate, geomorphology, lithology and vegetation.
Ecological Vegetation Class	A type of native vegetation classification that is described through a combination of its floristics, life form and ecological characteristics, and through an inferred fidelity to particular environment attributes. Each EVC includes a collection of floristic communities that occurs across a biogeographic range, and although differing in species, have similar habitat and ecological processes operating.
Exotic	Plants, animals, fungi and other organisms that have been introduced (deliberately or accidentally) to Australia or a given area after European settlement
Habitat Hectares	A measure of the quality and extent of native vegetation, incorporating attributes including presence of large trees, tree canopy health, understorey structure and diversity, weed cover and landscape context
High Threat Weed	Introduced species (including non-indigenous 'natives') with the ability to out-compete and substantially reduce one or more indigenous life forms in the longer term assuming on-going current site characteristics and disturbance regime.
Indigenous	Plant and animal species found naturally in pre-European Australia
Introduced	Deliberately or accidentally brought to Australia or part of Australia, usually by human agency
Native Vegetation	Indigenous vegetation includes vegetation that is native to Australia as well as being native to a specific geographic region.

Appendix 1 Edithvale-Seafood Wetlands Ramsar site land parcels, land status and land managers (from KBR 2009, DSE 2012; updated by DEPI 2013a and DELWP 2016d).

<http://services.land.vic.gov.au/maps/pmo.jsp>

Wetland	Land parcels (Allotments) (Parish of Lyndhurst)	Land status	Owner/formally assigned land manager	Land Manager (on-ground)
Edithvale North	Lot 1\TP131999, Lot 1\TP225777, Lot 2\TP225777, Lot 1\TP82835, Lot 1\TP414444, Lot 1\TP83139, Lot 1\TP820840, Lot 1\TP820843, Lot 2\TP820843 and Lot 1\TP385644.	Freehold	Melbourne Water	Melbourne Water
Edithvale South	Lot 1\TP95924, Lot 1\TP370109, Lot 1\TP366503, Lot 1\TP132070 and Lot 1\TP138507.	Freehold	Melbourne Water	Melbourne Water
Seafood Wetlands	<u>Northwest</u> : Lot 2\LP138935 and Lot 1\TP117202	Freehold	Frankston City Council	Frankston City Council
	<u>Northwest</u> : Lot 1\TP758882, Lot 1\TP659206, Lot 2\TP659206, Lot 3\TP659206, Lot 4\TP659206, Lot 5\TP659206, Lot 6\225759, Lot 5\225759, Lot 4\225759, Lot 3\225759, Lot 2\225759, Lot 1\225759 and Lot 1\TP824347	Freehold	Melbourne Water	Melbourne Water
	<u>Central</u> : Lot 2070\PP3025.	Freehold	Melbourne Water	Melbourne Water
	<u>Northeast</u> : Lot 1\TP382307.	Freehold	Melbourne Water	Melbourne Water
	<u>Northeast</u> : Lot 93\PP3025.	Freehold	Frankston City Council	Frankston City Council
	<u>Northeast</u> : Lot 86B\PP3025.	Crown Land reserved for conservation of area of natural interest	Melbourne Water as appointed Committee of Management under the <i>Crown Land (Reserves) Act 1978</i> , Victoria	Melbourne Water
<u>Southwest</u> : Lot 25\LP11828, Lot 26\LP11828, Lot 27\LP11828, Lot 28\LP11828, Lot 29\LP11828, Lot 30\LP11828, Lot 52\LP11828, Lot 53\LP11828, Lot 54\LP11828, Lot 55\LP11828, Lot 56\LP11828, Lot 57\LP11828, Lot 1\TP820912, Lot 1\TP821028, Lot 2\TP821028, Lot 2\TP824349, Lot 20\LP11828, Lot 21\LP11828, Lot 22\LP11828, Lot 23\LP11828, Lot 24\LP11828, Lot 21\LP13210, Lot 22\LP13210, Lot 23\LP13210, Lot 24\LP13210, Lot 25\LP13210, Lot 26\LP13210, Lot 97\LP13210, Lot 98\LP13210, Lot 99\LP13210, Lot 100\LP13210, Lot 101\LP13210, Lot 145\LP13210, Lot 146\LP13210, Lot 147\LP13210, Lot 148\LP13210, Lot 149\LP13210, Lot 150\LP13210, Lot 217\LP13210, Lot 218\LP13210, Lot 219\LP13210, Lot 220\LP13210, Lot 221\LP13210, Lot 222\LP13210, Lot 1\TP188903, Lot 1\TP376615, Lot 3\TP82102, Lot	Freehold	Melbourne Water	Melbourne Water	

Wetland	Land parcels (Allotments) (Parish of Lyndhurst)	Land status	Owner/formally assigned land manager	Land Manager (on-ground)
	4\850290, Lot 4\850290, Lot 2\TP850292, Lot 1\TP820882, Lot 2\TP820882, Lot 4\TP850292, Lot 1\TP820912, Lot 2\TP850299, Lot 4\TP850299, Lot 3\TP820882, Lot 1\TP820912, Lot 6 TP850299, Lot 12\LP6611, Lot 13\LP6611, Lot 14\LP6611, Lot 15\LP6611, Lot 16\LP6611, Lot 17\LP6611, Lot 18\LP6611, Lot 19\LP6611, Lot 20\LP6611, Lot 1\TP881753, Lot 22\LP6611, Lot 23\LP6611, Lot 24\LP6611, Lot 25\LP6611, Lot 2\TP850301, Lot 27\LP6611, Lot 28\LP6611, Lot 29\LP6611, Lot 30\LP6611, Lot 31\LP6611, Lot 32\LP6611, Lot 33\LP6611, Lot 34\LP6611, Lot 35\LP6611, Lot 36\LP6611, Lot 37\LP6611 and Lot 38 LP6611.			
	<u>Southwest:</u> Parcel with no Lot numbers, Lot 9\TP146701, Lot 1\TP850292, Lot 10\TP146701, Lot 3\TP850292, Lot 29\LP11717, Lot 30\LP11717, Lot 31\LP11717, Lot 32\LP11717, Lot 33\LP11717, Lot 34\LP11717, Lot 35\LP11717, Lot 36\LP11717, Lot 37\LP11717, Lot 38\LP11717, Lot 39\LP11717, Lot 54\LP11717, Lot 55\LP11717, Lot 56\LP11717, Lot 57\LP11717, Lot 58\LP11717, Lot 59\LP11717, Lot 60\LP11717, Lot 61\LP11717, Lot 62\LP11717, Lot 63\LP11717, Lot 64\LP11717, Lot 88\LP11717, Lot 89\LP11717, Lot 90\LP11717, Lot 91\LP11717, Lot 92\LP11717, Lot 93\LP11717, Lot 94\LP11717, Lot 95\LP11717, Lot 96\LP11717, Lot 97\LP11717, Lot 98\LP11717, Lot 28\LP12131, Lot 29\LP12131, Lot 30\LP12131, Lot 33\LP12131, Lot 34\LP12131, Lot 35\LP12131, Lot 36\LP12131, Lot 1\TP201413, Lot 1\TP912428, Lot 1\TP850299, Lot 3\TP850299, Lot 1\TP146701, Lot 2\TP146701, Lot 3\TP146701, Lot 4\TP146701, Lot 5\TP146701, Lot 6\TP146701, Lot 7\TP146701, Lot 8\TP146701, Lot 10\TP146701, Lot 11\TP146701, No parcel identification, Lot 118\LP11717, Lot 119\LP11717, Lot 143\LP11717, Lot 144\LP11717, Lot 168\LP11717, Lot 169\LP11717, Lot 10\LP6611, Lot 11\LP6611, Lot 5\TP850299, Lot 1,TP850301, Lot 39 LP6611, Lot 40 LP6611 and Lot 41 LP6611.	Freehold	Frankston City Council	Frankston City Council
	<u>Southeast:</u> Lot 1\TP824348, Lot 24\LP13454, Lot 25\LP13454, Lot 26\LP13454, Lot 27\LP13454, Lot 28\LP13454, Lot 29\LP13454, Lot 30\LP13454, Lot 31\LP13454, Lot 32\LP13454, Lot 33\LP13454, Lot 34\LP13454, Lot 35\LP13454, Lot 36\LP13454, Lot 37\LP13454, Lot 38\LP13454, Lot 39\LP13454, Lot 40\LP13454, Lot 41\LP13454, Lot 42\LP13454, Lot 43\LP13454, Lot 44\LP13454, Lot 45\LP13454, Lot 46\LP13454, Lot 47\LP13454, Lot 48\LP13454, Lot 49\LP13454, Lot 50\LP13454, Lot 77\LP13454, Lot 78\LP13454, Lot 79\LP13454, Lot 80\LP13454, Lot 81\LP13454, Lot 82\LP13454, Lot 83\LP13454, Lot 84\LP13454, Lot 85\LP13454, Lot 86\LP13454, Lot	Freehold	Melbourne Water	Melbourne Water

Wetland	Land parcels (Allotments) (Parish of Lyndhurst)	Land status	Owner/formally assigned land manager	Land Manager (on-ground)
	87\LP13454, Lot 88\LP13454, Lot 89\LP13454, Lot 1\TP850291, Lot 3\TP850291, Lot 45\LP13210, Lot 46\LP13210, Lot 76\LP13210, Lot 77\LP13210, Lot 164\LP13210, Lot 169\LP13210, Lot 200\LP13210, Lot 1\TP188903, Lot 1\TP850298, Lot 3 TP850298, Lot 1\TP820919, Lot 1\TP850300, Lot 3\TP850300, Lot 1\LP10032, Lot 2\LP10032, Lot 3\LP10032, Lot 4\LP10032, Lot 5\LP10032, Lot 6\LP10032, Lot 7\LP10032, Lot 8\LP10032, Lot 9\LP10032, Lot 10\LP10032, Lot 11\LP10032, Lot 12\LP10032, Lot 13\LP10032, Lot 14\LP10032, Lot 15\LP10032, Lot 17\LP10032, Lot 18\LP10032, Lot 19\LP10032, Lot 20\LP10032, Lot 21\LP10032, Lot 22\LP10032, Lot 23\LP10032, Lot 24\LP10032, Lot 25\LP10032, Lot 32\LP10032, Lot 33\LP10032, Lot 34\LP10032, Lot 35\LP10032, Lot 36\LP10032, Lot 37\LP10032, Lot 38\LP10032, Lot 39\LP10032, Lot 40\LP10032, Lot 41\LP10032, Lot 42\LP10032, Lot 43\LP10032, Lot 44\LP10032, Lot 45\LP10032, Lot 46\LP10032, 47\LP10032, Lot 48\LP10032, Lot 49\LP10032, Lot 1\TP865059			
	<u>Southeast:</u> Lot PC369422, Lot 1\TP944917, Lot 2\TP944917 and Lot 3\TP944917.	Freehold	Frankston City Council	Frankston City Council

Appendix 2 Significant flora species potentially occurring within the Edithvale-Seaford Wetlands.

A data review on the Victorian Biodiversity Atlas (DELWP 2016a) and the EPBC Act Protected Matters Search Tool (DoE 2016) was conducted for the site to identify rare or threatened flora species that have been recorded or may occur within 5 km of the site. These databases identified 19 rare or threatened plant species that may occur at Edithvale Wetlands and 22 rare or threatened plant species that may occur at Seaford Wetlands. A likelihood of occurrence has been assigned to each of these species at each site based on Ecology Australia's expert opinion and consideration of the following factors:

- Date(s) and number of past records;
- General condition and land use history of the study area, i.e. level of modification;
- Comparisons of site factors (climate, soils, topography) between the study area and sites known to support populations of each threatened species; and
- Whether there is a reasonable expectation that the species would have been recorded during the field survey.

Tables 15 and 16 present the likelihood of occurrence of rare or threatened flora species that could potentially occur Edithvale and Seaford Wetlands, respectively (species with a negligible likelihood of occurrence are not included). Of these, seven species at Edithvale Wetlands and five species at Seaford Wetlands are considered to have a moderate likelihood of occurrence given the proximity of relatively recent records and the presence of potentially suitable habitat.

Table 15 Significant flora species recorded (or modelled to possibly occur) within 5 km of the Edithvale Wetlands (excludes species with negligible likelihood of occurrence).

Key:

EPBC Act *Environment Protection and Biodiversity Conservation Act 1999*

VU Vulnerable under the EPBC Act

FFG Act *Flora and Fauna Guarantee Act 1988*

L Listed under the FFG Act

en Endangered as classified by DELWP (DEPI 2014, DSE 2013)

vu Vulnerable as classified by DELWP (DEPI 2014, DSE 2013)

r Rare as classified by DELWP (DEPI 2014)

k Poorly known as classified by DELWP (DEPI 2014)

Scientific Name	Common Name	Status	Likelihood of occurrence
<i>Amphibromus fluitans</i>	River Swamp Wallaby-grass	VU X	Moderate likelihood of occurrence – suitable habitat present
<i>Austrostipa rudis</i> subsp. <i>Australis</i>	Veined Spear-grass	r	Low likelihood of occurrence – potentially suitable habitat
<i>Carex chlorantha</i>	Green-top Sedge	k	Moderate likelihood of occurrence – suitable habitat present
<i>Coronidium gunnianum</i>	Pale Swamp Everlasting	vu	Moderate likelihood of occurrence – potentially habitat present and recorded nearby at the Peninsula Link interchange
<i>Correa alba</i> var. <i>pannosa</i>	Velvet White Correa	r	Low likelihood of occurrence – potentially suitable habitat
<i>Lachnagrostis punicea</i> subsp. <i>Filifolia</i>	Purple Blown-grass	r L	Moderate likelihood of occurrence – potentially suitable habitat
<i>Philydrum lanuginosum</i>	Woolly Waterlily	vu	Low likelihood of occurrence – modified habitat less suitable
<i>Ranunculus amplus</i>	Lacey River Buttercup	r	Moderate likelihood of occurrence – suitable habitat present and recent records nearby
<i>Ranunculus papulentus</i>	Large River Buttercup	k	Moderate likelihood of occurrence – suitable habitat present and recent records nearby
<i>Xerochrysum palustre</i>	Swamp Everlasting	VU vu L	Moderate likelihood of occurrence – suitable habitat present and recent records nearby

Table 16 Significant flora species recorded (or modelled to possibly occur) within 5 km of the Seaford Wetlands (excludes species with negligible likelihood of occurrence).

Key:

EPBC Act	<i>Environment Protection and Biodiversity Conservation Act 1999</i>
VU	Vulnerable under the EPBC Act
FFG Act	<i>Flora and Fauna Guarantee Act 1988</i>
L	Listed under the FFG Act
en	Endangered as classified by DELWP (DEPI 2014, DSE 2013)
vu	Vulnerable as classified by DELWP (DEPI 2014, DSE 2013)
r	Rare as classified by DELWP (DEPI 2014)
k	Poorly known as classified by DELWP (DEPI 2014)
▲	Species not recorded within 5 km of the study area (DELWP 2016a); identified as potentially occurring or potential habitat occurring by the EPBC Act protected matters search tool (DoE 2016)

Scientific Name	Common Name	Status	Likelihood of occurrence
<i>Amphibromus fluitans</i>	River Swamp Wallaby-grass	VU	Moderate likelihood of occurrence – suitable habitat present
<i>Chorizandra australis</i>	Southern Bristle-sedge	k	Moderate likelihood of occurrence – suitable habitat present
<i>Coronidium gunnianum</i>	Pale Swamp Everlasting	vu	Low likelihood of occurrence – suboptimal habitat present
<i>Eleocharis macbarronii</i>	Grey Spike-sedge	k	Moderate likelihood of occurrence – suitable habitat present and recent records nearby
<i>Lachnagrostis punicea</i> subsp. <i>Filifolia</i>	Purple Blown-grass	r L	Low likelihood of occurrence – Potentially suitable habitat
<i>Poa labillardierei</i> var. <i>(Volcanic Plains)</i>	Basalt Tussock-grass	k	Low likelihood of occurrence – prefers basalt habitats
<i>Poa poiformis</i> var. <i>ramifer</i>	Dune Poa	r	Low likelihood of occurrence – calcareous sands or shallow siliceous sands overlying basalt
<i>Ranunculus amplus</i>	Lacey River Buttercup	r	Moderate likelihood of occurrence – suitable habitat present and recent records nearby
<i>Xerochrysum palustre</i> ▲	Swamp Everlasting	VU vu L	Moderate likelihood of occurrence – suitable habitat present and recent records nearby

Appendix 3 Lists of species recorded and frequency of records of bird species recorded at Edithvale Wetlands from 1989 to 2015 and at Seafood Wetlands from 1994 to 2015 during monthly surveys undertaken by Bird Life Australia (summary of surveys from BirdLife Australia Atlas database).

Key:

- * denotes introduced species
- X Recorded
- EPBC *Environment Protection and Biodiversity Conservation Act 1999*
 - CE Listed as Critically Endangered under the EPBC Act
 - EN Listed as Endangered under the EPBC Act
 - VU Listed as Vulnerable under the EPBC Act
- Mi Migratory
- M Marine overfly
- FFG *Flora and Fauna Guarantee Act 1988*
 - L Listed as threatened under the FFG Act
 - N Nominated for listing under the FFG Act
- DSE Status according to DSE (2013): Advisory List of Threatened Vertebrate Fauna in Victoria – 2013:
 - ce classified as critically endangered in DSE (2013)
 - en classified as endangered in DSE (2013)
 - vu classified as vulnerable in DSE (2013)
 - nt classified as near threatened in DSE (2013)
 - dd classified as data deficient in DSE (2013)

Edithvale North Wetlands

Common Name	Scientific Name	No of sightings from 312 surveys	Most recent record	EPBC	FFG	DSE
Stubble Quail	<i>Coturnix pectoralis</i>	18	21-Dec-12	M		
Brown Quail	<i>Coturnix ypsilophora</i>	2	30-Jun-12			
Magpie Goose	<i>Anseranas semipalmata</i>	53	25-Sep-15	M	L	nt
Musk Duck	<i>Biziura lobata</i>	232	25-Sep-15	M		vu
Freckled Duck	<i>Stictonetta naevosa</i>	11	22-Apr-15		L	en
Black Swan	<i>Cygnus atratus</i>	271	20-Jul-15	M		
Australian Shelduck	<i>Tadorna tadornoides</i>	8	13-Jan-01			
Australian Wood Duck	<i>Chenonetta jubata</i>	25	16-Mar-15			
Pink-eared Duck	<i>Malacorhynchus membranaceus</i>	1	27-Aug-14			
Australasian Shoveler	<i>Anas rhynchos</i>	90	16-Jan-15			vu
Grey Teal	<i>Anas gracilis</i>	117	25-Sep-15			
Chestnut Teal	<i>Anas castanea</i>	270	25-Sep-15			
Pacific Black Duck	<i>Anas superciliosa</i>	297	25-Sep-15			
Hardhead	<i>Aythya australis</i>	153	16-Jun-15			vu
Blue-billed Duck	<i>Oxyura australis</i>	170	22-Apr-15		L	en

Common Name	Scientific Name	No of sightings from 312 surveys	Most recent record	EPBC	FFG	DSE
Australasian Grebe	<i>Tachybaptus novaehollandiae</i>	240	20-Jul-15			
Hoary-headed Grebe	<i>Poliocephalus poliocephalus</i>	244	25-Sep-15			
Great Crested Grebe	<i>Podiceps cristatus</i>	18	30-Dec-14			
*Rock Dove	<i>Columba livia</i>	63	12-Sep-13			
*Spotted Dove	<i>Streptopelia chinensis</i>	290	25-Sep-15			
Common Bronzewing	<i>Phaps chalcoptera</i>	6	13-Dec-13			
Brush Bronzewing	<i>Phaps elegans</i>	2	14-Feb-07			
Crested Pigeon	<i>Ocyphaps lophotes</i>	124	16-Jun-15			
Tawny Frogmouth	<i>Podargus strigoides</i>	3	18-Mar-14			
Fork-tailed Swift	<i>Apus pacificus</i>	1	23-Mar-07	Mi,M		
Australasian Darter	<i>Anhinga novaehollandiae</i>	4	21-Dec-12			
Little Pied Cormorant	<i>Microcarbo melanoleucos</i>	248	20-Jul-15			
Great Cormorant	<i>Phalacrocorax carbo</i>	37	22-Nov-13			
Little Black Cormorant	<i>Phalacrocorax sulcirostris</i>	33	26-May-15			
Pied Cormorant	<i>Phalacrocorax varius</i>	3	12-May-05			nt
Australian Pelican	<i>Pelecanus conspicillatus</i>	82	27-Aug-14	M		
Australasian Bittern	<i>Botaurus poiciloptilus</i>	98	20-Jul-15	EN	L	en
Australian Little Bittern	<i>Ixobrychus dubius</i>	3	14-Nov-12		L	en
White-necked Heron	<i>Ardea pacifica</i>	16	21-Nov-14			
Eastern Great Egret	<i>Ardea modesta</i>	56	25-Sep-15		L	vu
Intermediate Egret	<i>Ardea intermedia</i>	3	21-Feb-06	M	L	en
Cattle Egret	<i>Ardea ibis</i>	6	21-Nov-14	Mi,M		
White-faced Heron	<i>Egretta novaehollandiae</i>	266	25-Sep-15			
Little Egret	<i>Egretta garzetta</i>	3	18-Mar-12	M	L	en
Nankeen Night-Heron	<i>Nycticorax caledonicus</i>	10	13-Feb-15	M		nt
Glossy Ibis	<i>Plegadis falcinellus</i>	4	19-Sep-12	Mi,M		nt
Australian White Ibis	<i>Threskiornis molucca</i>	109	16-Jun-15	M		
Straw-necked Ibis	<i>Threskiornis spinicollis</i>	128	16-Jun-15	M		

Common Name	Scientific Name	No of sightings from 312 surveys	Most recent record	EPBC	FFG	DSE
Royal Spoonbill	<i>Platalea regia</i>	72	18-Aug-15			nt
Yellow-billed Spoonbill	<i>Platalea flavipes</i>	56	26-May-15			
Black-shouldered Kite	<i>Elanus axillaris</i>	139	18-Aug-15			
Whistling Kite	<i>Haliastur sphenurus</i>	18	16-Apr-14	M		
Brown Goshawk	<i>Accipiter fasciatus</i>	48	26-May-15	M		
Collared Sparrowhawk	<i>Accipiter cirrocephalus</i>	9	22-Apr-15			
Swamp Harrier	<i>Circus approximans</i>	189	25-Sep-15	M		
Little Eagle	<i>Hieraetus morphnoides</i>	5	18-Aug-15			
Nankeen Kestrel	<i>Falco cenchroides</i>	53	14-Nov-12	Mi,M		
Brown Falcon	<i>Falco berigora</i>	9	27-Apr-12			
Australian Hobby	<i>Falco longipennis</i>	69	16-Mar-15			
Peregrine Falcon	<i>Falco peregrinus</i>	3	25-Sep-15			
Purple Swamphen	<i>Porphyrio porphyrio</i>	283	25-Sep-15			
Lewin's Rail	<i>Lewinia pectoralis</i>	2	18-Mar-14		L	vu
Buff-banded Rail	<i>Gallirallus philippensis</i>	16	30-Dec-14	M		
Baillon's Crake	<i>Porzana pusilla</i>	47	22-Jan-14	M	L	vu
Australian Spotted Crake	<i>Porzana fluminea</i>	41	15-Sep-14			
Spotless Crake	<i>Porzana tabuensis</i>	29	18-Aug-15	M		
Black-tailed Native-hen	<i>Tribonyx ventralis</i>	3	13-Feb-13			
Dusky Moorhen	<i>Gallinula tenebrosa</i>	177	26-May-15			
Eurasian Coot	<i>Fulica atra</i>	264	16-Jun-15			
Black-winged Stilt	<i>Himantopus himantopus</i>	98	16-Jun-15	M		
Red-necked Avocet	<i>Recurvirostra novaehollandiae</i>	1	20-Aug-94	M		
Red-capped Plover	<i>Charadrius ruficapillus</i>	9	13-Feb-95	M		
Double-banded Plover	<i>Charadrius bicinctus</i>	2	02-May-91	Mi,M		
Black-fronted Dotterel	<i>Euseyornis melanops</i>	183	25-Sep-15			
Red-kneed Dotterel	<i>Erythronyx cinctus</i>	21	22-Apr-15			
Masked Lapwing	<i>Vanellus miles</i>	224	25-Sep-15			
Latham's Snipe	<i>Gallinago hardwickii</i>	113	25-Sep-15	Mi,M	N	nt
Common Greenshank	<i>Tringa nebularia</i>	16	20-Dec-07	Mi,M		vu
Marsh Sandpiper	<i>Tringa stagnatilis</i>	9	06-Dec-06	Mi,M		vu

Common Name	Scientific Name	No of sightings from 312 surveys	Most recent record	EPBC	FFG	DSE
Wood Sandpiper	<i>Tringa glareola</i>	5	12-Mar-01	Mi,M		vu
Red Knot	<i>Calidris canutus</i>	1	17-Nov-91	Mi,M		en
Red-necked Stint	<i>Calidris ruficollis</i>	8	06-Dec-06	Mi,M		
Long-toed Stint	<i>Calidris subminuta</i>	1	15-Jan-00	Mi,M		nt
Pectoral Sandpiper	<i>Calidris melanotos</i>	6	13-Feb-15	Mi,M		nt
Sharp-tailed Sandpiper	<i>Calidris acuminata</i>	45	13-Feb-15	Mi,M		
Curlew Sandpiper	<i>Calidris ferruginea</i>	6	26-Sep-94	CE,Mi,M		en
Whiskered Tern	<i>Chlidonias hybrida</i>	59	16-Jan-15	M		nt
White-winged Black Tern	<i>Chlidonias leucopterus</i>	2	14-Dec-95	Mi,M		nt
Pacific Gull	<i>Larus pacificus</i>	23	18-Mar-14	M		
Silver Gull	<i>Chroicocephalus novaehollandiae</i>	266	25-Sep-15	M		
Yellow-tailed Black-Cockatoo	<i>Calyptorhynchus funereus</i>	1	14-Jul-09			
Galah	<i>Eolophus roseicapillus</i>	62	18-Aug-15			
Long-billed Corella	<i>Cacatua tenuirostris</i>	4	16-Jun-15			
Little Corella	<i>Cacatua sanguinea</i>	9	22-Apr-15			
Sulphur-crested Cockatoo	<i>Cacatua galerita</i>	60	25-Sep-15			
Cockatiel	<i>Nymphicus hollandicus</i>	3	13-Feb-13			
Rainbow Lorikeet	<i>Trichoglossus haematodus</i>	121	18-Aug-15			
Musk Lorikeet	<i>Glossopsitta concinna</i>	43	22-Apr-15			
Little Lorikeet	<i>Glossopsitta pusilla</i>	1	06-Dec-09			
Crimson Rosella	<i>Platycercus elegans</i>	3	18-Mar-14			
Eastern Rosella	<i>Platycercus eximius</i>	187	25-Sep-15			
Red-rumped Parrot	<i>Psephotus haematonotus</i>	36	18-Aug-15			
Blue-winged Parrot	<i>Neophema chrysostoma</i>	5	28-May-14	M		
Horsfield's Bronze-Cuckoo	<i>Chalcites basalis</i>	40	21-Nov-14	M		
Pallid Cuckoo	<i>Cacomantis pallidus</i>	11	02-Feb-09	M		
Fan-tailed Cuckoo	<i>Cacomantis flabelliformis</i>	5	27-Aug-14	M		
Eastern Barn Owl	<i>Tyto javanica</i>	1	27-Apr-98			

Common Name	Scientific Name	No of sightings from 312 surveys	Most recent record	EPBC	FFG	DSE
Laughing Kookaburra	<i>Dacelo novaeguineae</i>	2	24-Apr-13			
Sacred Kingfisher	<i>Todiramphus sanctus</i>	6	21-Nov-14	M		
Superb Fairy-wren	<i>Malurus cyaneus</i>	229	25-Sep-15			
White-browed Scrubwren	<i>Sericornis frontalis</i>	74	25-Sep-15			
Yellow Thornbill	<i>Acanthiza nana</i>	17	10-Jul-98			
Yellow-rumped Thornbill	<i>Acanthiza chrysorrhoa</i>	9	23-Jul-94			
Brown Thornbill	<i>Acanthiza pusilla</i>	139	25-Sep-15			
Spotted Pardalote	<i>Pardalotus punctatus</i>	36	16-Jun-15			
Striated Pardalote	<i>Pardalotus striatus</i>	5	26-May-15			
Eastern Spinebill	<i>Acanthorhynchus tenuirostris</i>	2	23-Mar-07			
White-plumed Honeyeater	<i>Lichenostomus penicillatus</i>	263	18-Aug-15			
Noisy Miner	<i>Manorina melanocephala</i>	222	25-Sep-15			
Spiny-cheeked Honeyeater	<i>Acanthagenys rufogularis</i>	10	22-Apr-15			
Little Wattlebird	<i>Anthochaera chrysoptera</i>	140	25-Sep-15			
Red Wattlebird	<i>Anthochaera carunculata</i>	253	25-Sep-15			
White-fronted Chat	<i>Epthianura albifrons</i>	44	10-Jul-03			
New Holland Honeyeater	<i>Phylidonyris novaehollandiae</i>	9	27-Apr-12			
Noisy Friarbird	<i>Philemon corniculatus</i>	1	18-Sep-08			
Black-faced Cuckoo-shrike	<i>Coracina novaehollandiae</i>	73	18-Aug-15	M		
White-winged Triller	<i>Lalage sueurii</i>	1	23-Oct-11			
Golden Whistler	<i>Pachycephala pectoralis</i>	29	20-Jul-15			
Grey Shrike-thrush	<i>Colluricincla harmonica</i>	21	24-Apr-13			
Olive-backed Oriole	<i>Oriolus sagittatus</i>	1	16-Nov-05			
Grey Butcherbird	<i>Cracticus torquatus</i>	132	25-Sep-15			
Australian Magpie	<i>Cracticus tibicen</i>	225	25-Sep-15			
Pied Currawong	<i>Strepera graculina</i>	2	11-Jul-07			
Rufous Fantail	<i>Rhipidura rufifrons</i>	1	08-Dec-98			

Common Name	Scientific Name	No of sightings from 312 surveys	Most recent record	EPBC	FFG	DSE
Grey Fantail	<i>Rhipidura albiscapa</i>	96	25-Sep-15			
Willie Wagtail	<i>Rhipidura leucophrys</i>	270	25-Sep-15			
Little Raven	<i>Corvus mellori</i>	251	25-Sep-15	M		
Magpie-lark	<i>Grallina cyanoleuca</i>	309	25-Sep-15			
Scarlet Robin	<i>Petroica boodang</i>	1	16-Jan-15			
Flame Robin	<i>Petroica phoenicea</i>	91	18-Aug-15	M		
Horsfield's Bushlark	<i>Mirafrja javanica</i>	1	24-Oct-10			
*Eurasian Skylark	<i>Alauda arvensis</i>	236	25-Sep-15			
Golden-headed Cisticola	<i>Cisticola exilis</i>	308	25-Sep-15			
Australian Reed-Warbler	<i>Acrocephalus australis</i>	181	25-Sep-15	Mi,M		
Little Grassbird	<i>Megalurus gramineus</i>	263	25-Sep-15			
Rufous Songlark	<i>Cincloramphus mathewsi</i>	1	20-Dec-07			
Brown Songlark	<i>Cincloramphus cruralis</i>	1	13-Dec-13			
Silvereeye	<i>Zosterops lateralis</i>	22	18-Aug-15	M		
Welcome Swallow	<i>Hirundo neoxena</i>	298	25-Sep-15			
Fairy Martin	<i>Petrochelidon ariel</i>	3	11-May-95			
Tree Martin	<i>Petrochelidon nigricans</i>	5	07-Feb-14	M		
*Common Blackbird	<i>Turdus merula</i>	276	25-Sep-15			
*Common Starling	<i>Sturnus vulgaris</i>	302	18-Aug-15			
*Common Myna	<i>Sturnus tristis</i>	285	25-Sep-15			
Mistletoebird	<i>Dicaeum hirundinaceum</i>	2	12-Apr-04			
Red-browed Finch	<i>Neochmia temporalis</i>	11	13-Feb-15			
*House Sparrow	<i>Passer domesticus</i>	126	13-Feb-15			
*Eurasian Tree Sparrow	<i>Passer montanus</i>	10	12-May-05			
Australasian Pipit	<i>Anthus novaeseelandiae</i>	21	18-Mar-12	M		
*European Goldfinch	<i>Carduelis carduelis</i>	213	18-Aug-15			
*Common Greenfinch	<i>Chloris chloris</i>	160	25-Sep-15			
*Black Duck-Mallard hybrid	<i>Anas superciliosa – Anas platyrhynchos</i>	1	10-Jul-03			

Edithvale South Wetlands

Common Name	Scientific Name	No of sightings from 312 surveys	Most recent record	EPBC	FFG	DSE
Stubble Quail	<i>Coturnix pectoralis</i>	1	03-Dec-90	M		
Brown Quail	<i>Coturnix ypsilophora</i>	2	12-May-01			
Magpie Goose	<i>Anseranas semipalmata</i>	35	20-Jul-15	M	L	nt
Musk Duck	<i>Biziura lobata</i>	14	14-Jun-11	M		vu
Freckled Duck	<i>Stictonetta naevosa</i>	5	18-Jan-12		L	en
Cape Barren Goose	<i>Cereopsis novaehollandiae</i>	2	24-Mar-07	M		
Black Swan	<i>Cygnus atratus</i>	253	25-Sep-15			
Australian Shelduck	<i>Tadorna tadornoides</i>	19	30-Dec-14			
Australian Wood Duck	<i>Chenonetta jubata</i>	29	21-Aug-13			
Pink-eared Duck	<i>Malacorhynchus membranaceus</i>	15	23-Dec-11			
Australasian Shoveler	<i>Anas rhynchotis</i>	189	25-Sep-15			vu
Grey Teal	<i>Anas gracilis</i>	212	25-Sep-15			
Chestnut Teal	<i>Anas castanea</i>	259	25-Sep-15			
*Northern Mallard	<i>Anas platyrhynchos</i>	2	26-Nov-96			
Pacific Black Duck	<i>Anas superciliosa</i>	258	25-Sep-15			
Hardhead	<i>Aythya australis</i>	105	30-Dec-14			vu
Blue-billed Duck	<i>Oxyura australis</i>	110	17-Oct-14		L	en
Australasian Grebe	<i>Tachybaptus novaehollandiae</i>	201	25-Sep-15			
Hoary-headed Grebe	<i>Poliocephalus poliocephalus</i>	187	25-Sep-15			
*Rock Dove	<i>Columba livia</i>	31	21-Aug-13			
*Spotted Dove	<i>Streptopelia chinensis</i>	306	25-Sep-15			
Common Bronzewing	<i>Phaps chalcoptera</i>	12	16-Mar-15			
Brush Bronzewing	<i>Phaps elegans</i>	1	14-Apr-95			
Crested Pigeon	<i>Ocyphaps lophotes</i>	107	25-Sep-15			

Common Name	Scientific Name	No of sightings from 312 surveys	Most recent record	EPBC	FFG	DSE
Tawny Frogmouth	<i>Podargus strigoides</i>	1	19-May-11			
Short-tailed Shearwater	<i>Ardenna tenuirostris</i>	1	29-Oct-00	Mi,M		
Australasian Darter	<i>Anhinga novaehollandiae</i>	6	25-Oct-13			
Little Pied Cormorant	<i>Microcarbo melanoleucos</i>	157	25-Sep-15			
Great Cormorant	<i>Phalacrocorax carbo</i>	66	18-Mar-14			
Little Black Cormorant	<i>Phalacrocorax sulcirostris</i>	42	18-Aug-15			
Pied Cormorant	<i>Phalacrocorax varius</i>	2	23-Oct-11			nt
Australian Pelican	<i>Pelecanus conspicillatus</i>	96	25-Sep-15	M		
Australasian Bittern	<i>Botaurus poiciloptilus</i>	66	20-Jul-15	EN	L	en
Australian Little Bittern	<i>Ixobrychus dubius</i>	5	26-Jun-13		L	en
White-necked Heron	<i>Ardea pacifica</i>	38	30-Dec-14			
Eastern Great Egret	<i>Ardea modesta</i>	51	25-Sep-15	Mi,M	L	vu
Intermediate Egret	<i>Ardea intermedia</i>	2	10-Mar-06	M	L	en
Cattle Egret	<i>Ardea ibis</i>	21	21-Nov-14	Mi,M		
White-faced Heron	<i>Egretta novaehollandiae</i>	238	25-Sep-15			
Little Egret	<i>Egretta garzetta</i>	4	14-Feb-12	M	L	en
Nankeen Night-Heron	<i>Nycticorax caledonicus</i>	13	02-Jan-13	M		nt
Glossy Ibis	<i>Plegadis falcinellus</i>	12	16-Jan-15	Mi,M		nt
Australian White Ibis	<i>Threskiornis molucca</i>	146	25-Sep-15	M		
Straw-necked Ibis	<i>Threskiornis spinicollis</i>	146	25-Sep-15	M		
Royal Spoonbill	<i>Platalea regia</i>	70	13-Feb-15			nt
Yellow-billed Spoonbill	<i>Platalea flavipes</i>	80	16-Jan-15			
Black-shouldered Kite	<i>Elanus axillaris</i>	149	26-May-15			
White-bellied Sea-Eagle	<i>Haliaeetus leucogaster</i>	1	16-Oct-95	Mi,M	L	vu
Whistling Kite	<i>Haliastur sphenurus</i>	17	25-Sep-15	M		
Brown Goshawk	<i>Accipiter fasciatus</i>	114	20-Jul-15	M		
Collared Sparrowhawk	<i>Accipiter cirrocephalus</i>	19	26-May-15			

Common Name	Scientific Name	No of sightings from 312 surveys	Most recent record	EPBC	FFG	DSE
Spotted Harrier	<i>Circus assimilis</i>	1	02-Jan-13			nt
Swamp Harrier	<i>Circus approximans</i>	218	25-Sep-15	M		
Little Eagle	<i>Hieraaetus morphnoides</i>	1	21-Feb-05			
Nankeen Kestrel	<i>Falco cenchroides</i>	30	28-May-14	Mi,M		
Brown Falcon	<i>Falco berigora</i>	16	21-May-13			
Australian Hobby	<i>Falco longipennis</i>	49	27-Aug-14			
Peregrine Falcon	<i>Falco peregrinus</i>	5	21-Jun-10			
Purple Swamphen	<i>Porphyrio porphyrio</i>	285	25-Sep-15			
Lewin's Rail	<i>Lewinia pectoralis</i>	4	10-May-07		L	vu
Buff-banded Rail	<i>Gallirallus philippensis</i>	37	22-Jan-14	M		
Baillon's Crake	<i>Porzana pusilla</i>	43	13-Feb-15	M	L	vu
Australian Spotted Crake	<i>Porzana fluminea</i>	78	15-Sep-14			
Spotless Crake	<i>Porzana tabuensis</i>	77	12-Sep-13	M		
Black-tailed Native-hen	<i>Tribonyx ventralis</i>	12	22-Jan-14			
Dusky Moorhen	<i>Gallinula tenebrosa</i>	225	16-Jan-15			
Eurasian Coot	<i>Fulica atra</i>	219	16-Jan-15			
Black-winged Stilt	<i>Himantopus himantopus</i>	150	25-Sep-15	M		
Red-necked Avocet	<i>Recurvirostra novaehollandiae</i>	6	02-Jan-13	M		
Red-capped Plover	<i>Charadrius ruficapillus</i>	2	16-Dec-97	M		
Black-fronted Dotterel	<i>Euseyonis melanops</i>	45	18-Aug-15			
Red-kneed Dotterel	<i>Erythronyx cinctus</i>	54	16-Jan-15			
Masked Lapwing	<i>Vanellus miles</i>	188	18-Aug-15			
Australian Painted Snipe	<i>Rostratula australis</i>	2	09-Dec-08	CE; Mi,M	L	ce
Latham's Snipe	<i>Gallinago hardwickii</i>	82	16-Jan-15	Mi,M	N	nt
Little Curlew	<i>Numenius minutus</i>	1	17-Aug-90	Mi,M		
Common Greenshank	<i>Tringa nebularia</i>	42	25-Oct-13	Mi,M		vu
Marsh Sandpiper	<i>Tringa stagnatilis</i>	38	16-Jan-15	Mi,M		vu
Wood Sandpiper	<i>Tringa glareola</i>	49	16-Jan-15	Mi,M		vu

Common Name	Scientific Name	No of sightings from 312 surveys	Most recent record	EPBC	FFG	DSE
Red-necked Stint	<i>Calidris ruficollis</i>	18	16-Jan-15	Mi,M		
Long-toed Stint	<i>Calidris subminuta</i>	1	07-Feb-02	Mi,M		nt
Pectoral Sandpiper	<i>Calidris melanotos</i>	13	16-Jan-15	Mi,M		nt
Sharp-tailed Sandpiper	<i>Calidris acuminata</i>	94	16-Jan-15	Mi,M		
Curlew Sandpiper	<i>Calidris ferruginea</i>	21	16-Jan-15	CE,Mi,M		en
Caspian Tern	<i>Hydroprogne caspia</i>	7	16-Jan-15	Mi,M	L	nt
Whiskered Tern	<i>Chlidonias hybrida</i>	76	25-Sep-15	M		nt
White-winged Black Tern	<i>Chlidonias leucopterus</i>	6	16-Nov-05	Mi,M		nt
Common Tern	<i>Sterna hirundo</i>	1	08-Jan-05	Mi,M		
Crested Tern	<i>Thalasseus bergii</i>	1	20-Jul-05	M		
Pacific Gull	<i>Larus pacificus</i>	34	14-Nov-12	M		
Silver Gull	<i>Chroicocephalus novaehollandiae</i>	234	25-Sep-15	M		
Yellow-tailed Black-Cockatoo	<i>Calyptorhynchus funereus</i>	2	23-Oct-11			
Galah	<i>Eolophus roseicapillus</i>	59	18-Aug-15			
Long-billed Corella	<i>Cacatua tenuirostris</i>	8	19-Sep-12			
Little Corella	<i>Cacatua sanguinea</i>	9	16-Jun-15			
Sulphur-crested Cockatoo	<i>Cacatua galerita</i>	80	18-Aug-15			
Rainbow Lorikeet	<i>Trichoglossus haematodus</i>	120	18-Aug-15			
Musk Lorikeet	<i>Glossopsitta concinna</i>	15	16-Mar-15			
Crimson Rosella	<i>Platycercus elegans</i>	1	17-Jul-13			
Eastern Rosella	<i>Platycercus eximius</i>	31	18-Aug-15			
Red-rumped Parrot	<i>Psephotus haematotus</i>	42	25-Sep-15			
Blue-winged Parrot	<i>Neophema chrysostoma</i>	4	16-Jun-15	M		
Horsfield's Bronze-Cuckoo	<i>Chalcites basalus</i>	60	25-Sep-15	M		
Black-eared Cuckoo	<i>Chalcites osculans</i>	1	20-Sep-96	M		
Pallid Cuckoo	<i>Cacomantis pallidus</i>	6	13-Feb-13	M		
Fan-tailed Cuckoo	<i>Cacomantis flabelliformis</i>	6	16-Mar-15	M		

Common Name	Scientific Name	No of sightings from 312 surveys	Most recent record	EPBC	FFG	DSE
Southern Boobook	<i>Ninox novaeseelandiae</i>	2	24-Apr-13	M		
Eastern Barn Owl	<i>Tyto javanica</i>	5	19-Sep-12			
Laughing Kookaburra	<i>Dacelo novaeguineae</i>	1	21-May-13			
Sacred Kingfisher	<i>Todiramphus sanctus</i>	8	22-Nov-13	M		
Superb Fairy-wren	<i>Malurus cyaneus</i>	279	25-Sep-15			
White-browed Scrubwren	<i>Sericornis frontalis</i>	109	25-Sep-15			
Yellow-rumped Thornbill	<i>Acanthiza chrysorrhoa</i>	40	16-Jun-15			
Brown Thornbill	<i>Acanthiza pusilla</i>	92	25-Sep-15			
Spotted Pardalote	<i>Pardalotus punctatus</i>	45	20-Jul-15			
Striated Pardalote	<i>Pardalotus striatus</i>	11	26-Jun-13			
Eastern Spinebill	<i>Acanthorhynchus tenuirostris</i>	14	18-Aug-15			
Yellow-faced Honeyeater	<i>Lichenostomus chrysops</i>	7	16-May-12			
White-plumed Honeyeater	<i>Lichenostomus penicillatus</i>	305	25-Sep-15			
Noisy Miner	<i>Manorina melanocephala</i>	73	25-Sep-15			
Spiny-cheeked Honeyeater	<i>Acanthagenys rufogularis</i>	17	22-Apr-15			
Little Wattlebird	<i>Anthochaera chrysoptera</i>	208	25-Sep-15			
Red Wattlebird	<i>Anthochaera carunculata</i>	270	25-Sep-15			
White-fronted Chat	<i>Epthianura albifrons</i>	12	23-Mar-02			
New Holland Honeyeater	<i>Phylidonyris novaehollandiae</i>	159	25-Sep-15			
White-naped Honeyeater	<i>Melithreptus lunatus</i>	6	21-Jun-10			
Black-faced Cuckoo-shrike	<i>Coracina novaehollandiae</i>	104	18-Aug-15	M		
Crested Shrike-tit	<i>Falcunculus frontatus</i>	3	10-Sep-04			
Golden Whistler	<i>Pachycephala pectoralis</i>	24	15-Sep-14			
Rufous Whistler	<i>Pachycephala rufiventris</i>	2	17-Oct-12			
Grey Shrike-thrush	<i>Colluricincla harmonica</i>	59	20-Jun-14			

Common Name	Scientific Name	No of sightings from 312 surveys	Most recent record	EPBC	FFG	DSE
Olive-backed Oriole	<i>Oriolus sagittatus</i>	1	15-Sep-06			
White-browed Woodswallow	<i>Artamus superciliosus</i>	1	18-Mar-14			
Grey Butcherbird	<i>Cracticus torquatus</i>	83	18-Aug-15			
Australian Magpie	<i>Cracticus tibicen</i>	194	25-Sep-15			
Pied Currawong	<i>Strepera graculina</i>	1	15-Sep-14			
Grey Fantail	<i>Rhipidura albiscapa</i>	80	16-Jun-15			
Willie Wagtail	<i>Rhipidura leucophrys</i>	291	25-Sep-15			
Little Raven	<i>Corvus mellori</i>	284	25-Sep-15	M		
Satin Flycatcher	<i>Myiagra cyanoleuca</i>	1	16-Mar-15	Mi,M		
Restless Flycatcher	<i>Myiagra inquieta</i>	1	15-Mar-96			
Magpie-lark	<i>Grallina cyanoleuca</i>	308	25-Sep-15			
Scarlet Robin	<i>Petroica boodang</i>	2	16-Jan-15			
Flame Robin	<i>Petroica phoenicea</i>	96	18-Aug-15	M		
*Eurasian Skylark	<i>Alauda arvensis</i>	127	28-May-14			
Golden-headed Cisticola	<i>Cisticola exilis</i>	276	25-Sep-15			
Australian Reed-Warbler	<i>Acrocephalus australis</i>	207	25-Sep-15	Mi,M		
Little Grassbird	<i>Megalurus gramineus</i>	292	25-Sep-15			
Rufous Songlark	<i>Cincloramphus mathewsi</i>	1	23-Oct-11			
Silvereeye	<i>Zosterops lateralis</i>	84	20-Jul-15	M		
Welcome Swallow	<i>Hirundo neoxena</i>	308	25-Sep-15			
Fairy Martin	<i>Petrochelidon ariel</i>	17	14-Aug-12			
Tree Martin	<i>Petrochelidon nigricans</i>	11	11-Mar-09	M		
*Common Blackbird	<i>Turdus merula</i>	309	25-Sep-15			
*Song Thrush	<i>Turdus philomelos</i>	5	10-Oct-08			
*Common Starling	<i>Sturnus vulgaris</i>	308	25-Sep-15			
*Common Myna	<i>Sturnus tristis</i>	303	25-Sep-15			
Mistletoebird	<i>Dicaeum hirundinaceum</i>	1	23-Mar-02			
Red-browed Finch	<i>Neochmia temporalis</i>	49	25-Sep-15			
*House Sparrow	<i>Passer domesticus</i>	259	25-Sep-15			
*Eurasian Tree Sparrow	<i>Passer montanus</i>	22	05-Feb-08			
Australasian Pipit	<i>Anthus novaeseelandiae</i>	4	18-Mar-12	M		
*European Goldfinch	<i>Carduelis carduelis</i>	226	25-Sep-15			

Common Name	Scientific Name	No of sightings from 312 surveys	Most recent record	EPBC	FFG	DSE
*Common Greenfinch	<i>Chloris chloris</i>	253	25-Sep-15			
*Domestic Goose	<i>Anser anser</i>	2	30-Dec-14			
*Domestic Duck	<i>Anas platyrhynchos</i>	7	13-Feb-15			

Seafood Wetlands

Common Name	Scientific Name	No of sightings from 252 surveys	Most recent record	EPBC	FFG	DSE
Stubble Quail	<i>Coturnix pectoralis</i>	3	25-Nov-11	M		
Brown Quail	<i>Coturnix ypsilophora</i>	7	27-Nov-13			
Musk Duck	<i>Biziura lobata</i>	8	26-Mar-13	M		vu
Freckled Duck	<i>Stictonetta naevosa</i>	18	31-Dec-14		L	en
Cape Barren Goose	<i>Cereopsis novaehollandiae</i>	4	26-Jul-10	M		
Black Swan	<i>Cygnus atratus</i>	225	29-Sep-15	M		
Australian Shelduck	<i>Tadorna tadornoides</i>	37	25-Mar-15			
Australian Wood Duck	<i>Chenonetta jubata</i>	38	23-Jul-15			
Pink-eared Duck	<i>Malacorhynchus membranaceus</i>	28	30-Apr-15			
Australasian Shoveler	<i>Anas rhynchotis</i>	54	30-Apr-15			vu
Grey Teal	<i>Anas gracilis</i>	219	29-Sep-15			
Chestnut Teal	<i>Anas castanea</i>	248	29-Sep-15			
*Northern Mallard	<i>Anas platyrhynchos</i>	4	24-Jun-05			
Pacific Black Duck	<i>Anas superciliosa</i>	242	29-Sep-15			
Hardhead	<i>Aythya australis</i>	103	26-Jun-15		L	en
Blue-billed Duck	<i>Oxyura australis</i>	106	20-Feb-15			
Australasian Grebe	<i>Tachybaptus novaehollandiae</i>	89	31-Aug-15			
Hoary-headed Grebe	<i>Poliiocephalus poliocephalus</i>	197	29-Sep-15			
Great Crested Grebe	<i>Podiceps cristatus</i>	3	31-Oct-13			
*Rock Dove	<i>Columba livia</i>	65	26-Apr-13			
*Spotted Dove	<i>Streptopelia chinensis</i>	251	29-Sep-15			

Common Name	Scientific Name	No of sightings from 252 surveys	Most recent record	EPBC	FFG	DSE
Common Bronzewing	<i>Phaps chalcoptera</i>	60	29-Sep-15			
Brush Bronzewing	<i>Phaps elegans</i>	1	21-Feb-07			
Crested Pigeon	<i>Ocyphaps lophotes</i>	137	29-Sep-15			
White-throated Needletail	<i>Hirundapus caudacutus</i>	2	03-Apr-96	Mi,M		vu
Australasian Darter	<i>Anhinga novaehollandiae</i>	13	28-Apr-14			
Little Pied Cormorant	<i>Microcarbo melanoleucos</i>	93	25-Mar-15			
Great Cormorant	<i>Phalacrocorax carbo</i>	73	25-Mar-15			
Little Black Cormorant	<i>Phalacrocorax sulcirostris</i>	112	25-Mar-15			
Australian Pelican	<i>Pelecanus conspicillatus</i>	177	29-Sep-15	M		
Australasian Bittern	<i>Botaurus poiciloptilus</i>	25	31-Aug-15	EN	L	en
Australian Little Bittern	<i>Ixobrychus dubius</i>	1	09-Dec-94		L	en
White-necked Heron	<i>Ardea pacifica</i>	15	31-Oct-14			
Eastern Great Egret	<i>Ardea modesta</i>	110	29-Sep-15	Mi,M	L	vu
Intermediate Egret	<i>Ardea intermedia</i>	1	24-Feb-12	M	L	en
Cattle Egret	<i>Ardea ibis</i>	11	26-Apr-11	Mi,M		
White-faced Heron	<i>Egretta novaehollandiae</i>	207	29-Sep-15			
Little Egret	<i>Egretta garzetta</i>	3	25-Feb-13	M	L	en
Nankeen Night-Heron	<i>Nycticorax caledonicus</i>	3	24-Feb-12	M		nt
Glossy Ibis	<i>Plegadis falcinellus</i>	2	29-Sep-15	Mi,M		nt
Australian White Ibis	<i>Threskiornis molucca</i>	118	29-Sep-15	M		
Straw-necked Ibis	<i>Threskiornis spinicollis</i>	112	29-Sep-15	M		
Royal Spoonbill	<i>Platalea regia</i>	128	25-Mar-15			nt
Yellow-billed Spoonbill	<i>Platalea flavipes</i>	64	31-Dec-14			
Black-shouldered Kite	<i>Elanus axillaris</i>	184	31-Aug-15			

Common Name	Scientific Name	No of sightings from 252 surveys	Most recent record	EPBC	FFG	DSE
White-bellied Sea-Eagle	<i>Haliaeetus leucogaster</i>	2	26-Jun-15	Mi,M	L	vu
Whistling Kite	<i>Haliastur sphenurus</i>	29	23-Jul-15	M		
Brown Goshawk	<i>Accipiter fasciatus</i>	106	29-Sep-15	M		
Collared Sparrowhawk	<i>Accipiter cirrocephalus</i>	27	30-Apr-15			
Swamp Harrier	<i>Circus approximans</i>	202	29-Sep-15	M		
Wedge-tailed Eagle	<i>Aquila audax</i>	1	26-Apr-11			
Little Eagle	<i>Hieraaetus morphnoides</i>	7	26-Apr-11			
Nankeen Kestrel	<i>Falco cenchroides</i>	100	29-Sep-15	Mi,M		
Brown Falcon	<i>Falco berigora</i>	27	27-Jan-15			
Australian Hobby	<i>Falco longipennis</i>	69	31-Dec-14			
Peregrine Falcon	<i>Falco peregrinus</i>	15	20-Feb-15			
Purple Swamphen	<i>Porphyrio porphyrio</i>	247	29-Sep-15			
Lewin's Rail	<i>Lewinia pectoralis</i>	2	27-Nov-13		L	vu
Buff-banded Rail	<i>Gallirallus philippensis</i>	10	27-Nov-13	M		
Baillon's Crake	<i>Porzana pusilla</i>	20	20-Feb-15	M	L	vu
Australian Spotted Crake	<i>Porzana fluminea</i>	34	29-Jan-14			
Spotless Crake	<i>Porzana tabuensis</i>	38	20-Feb-15	M		
Black-tailed Native-hen	<i>Tribonyx ventralis</i>	4	31-Jan-13			
Dusky Moorhen	<i>Gallinula tenebrosa</i>	127	27-Jan-15			
Eurasian Coot	<i>Fulica atra</i>	159	29-May-15			
Black-winged Stilt	<i>Himantopus himantopus</i>	109	26-Jun-15	M		
Red-necked Avocet	<i>Recurvirostra novaehollandiae</i>	17	31-Aug-15	M		
Red-capped Plover	<i>Charadrius ruficapillus</i>	35	23-Nov-02	M		
Double-banded Plover	<i>Charadrius bicinctus</i>	3	21-Mar-14	Mi,M		
Black-fronted Dotterel	<i>Euseyonis melanops</i>	105	31-Aug-15			
Red-kneed Dotterel	<i>Erythronyctes cinctus</i>	34	26-Jun-15			
Masked Lapwing	<i>Vanellus miles</i>	215	29-Sep-15			

Common Name	Scientific Name	No of sightings from 252 surveys	Most recent record	EPBC	FFG	DSE
Latham's Snipe	<i>Gallinago hardwickii</i>	93	29-Sep-15	Mi,M	N	nt
Black-tailed Godwit	<i>Limosa limosa</i>	2	19-May-05	Mi,M		vu
Common Greenshank	<i>Tringa nebularia</i>	60	20-Feb-15	Mi,M		vu
Marsh Sandpiper	<i>Tringa stagnatilis</i>	7	21-Mar-14	Mi,M		vu
Wood Sandpiper	<i>Tringa glareola</i>	2	06-Jan-06	Mi,M		vu
Red-necked Stint	<i>Calidris ruficollis</i>	15	27-Jan-15	Mi,M		
Long-toed Stint	<i>Calidris subminuta</i>	1	27-Feb-14	Mi,M		nt
Pectoral Sandpiper	<i>Calidris melanotos</i>	6	27-Jan-15	Mi,M		nt
Sharp-tailed Sandpiper	<i>Calidris acuminata</i>	61	25-Mar-15	Mi,M		
Curlew Sandpiper	<i>Calidris ferruginea</i>	15	27-Jan-15	CE,Mi,M		en
Red-necked Phalarope	<i>Phalaropus lobatus</i>	2	06-Jan-06	Mi,M		
Caspian Tern	<i>Hydroprogne caspia</i>	2	04-Jan-02	Mi,M	L	nt
Whiskered Tern	<i>Chlidonias hybrida</i>	34	29-Sep-15	M		nt
Common Tern	<i>Sterna hirundo</i>	1	07-Nov-99	Mi,M		
Pacific Gull	<i>Larus pacificus</i>	28	27-Feb-14	M		
Silver Gull	<i>Chroicocephalus novaehollandiae</i>	245	29-Sep-15	M		
Yellow-tailed Black-Cockatoo	<i>Calyptorhynchus funereus</i>	4	24-Oct-11			
Galah	<i>Eolophus roseicapillus</i>	82	29-Sep-15			
Long-billed Corella	<i>Cacatua tenuirostris</i>	1	26-Mar-13			
Little Corella	<i>Cacatua sanguinea</i>	4	25-Jul-13			
Sulphur-crested Cockatoo	<i>Cacatua galerita</i>	43	30-Apr-15			
Cockatiel	<i>Nymphicus hollandicus</i>	1	28-Nov-14			
Rainbow Lorikeet	<i>Trichoglossus haematodus</i>	242	29-Sep-15			
Scaly-breasted Lorikeet	<i>Trichoglossus chlorolepidotus</i>	1	18-Mar-11			
Musk Lorikeet	<i>Glossopsitta concinna</i>	84	23-Jul-15			
Purple-crowned Lorikeet	<i>Glossopsitta porphyrocephala</i>	1	18-Mar-11			

Common Name	Scientific Name	No of sightings from 252 surveys	Most recent record	EPBC	FFG	DSE
Crimson Rosella	<i>Platycercus elegans</i>	2	10-Apr-99			
Eastern Rosella	<i>Platycercus eximius</i>	197	29-Sep-15			
Swift Parrot	<i>Lathamus discolor</i>	4	25-Mar-15	En; M	L	en
Red-rumped Parrot	<i>Psephotus haematonotus</i>	3	26-Jun-14			
Blue-winged Parrot	<i>Neophema chrysostoma</i>	10	31-Oct-14	M		
Horsfield's Bronze-Cuckoo	<i>Chalcites basalis</i>	83	29-Sep-15	M		
Shining Bronze-Cuckoo	<i>Chalcites lucidus</i>	4	27-Sep-13	M		
Pallid Cuckoo	<i>Cacomantis pallidus</i>	22	27-Sep-13	M		
Fan-tailed Cuckoo	<i>Cacomantis flabelliformis</i>	29	30-Apr-15	M		
Eastern Barn Owl	<i>Tyto javanica</i>	3	30-May-12			
Laughing Kookaburra	<i>Dacelo novaeguineae</i>	4	28-May-13			
Sacred Kingfisher	<i>Todiramphus sanctus</i>	6	25-Nov-11	M		
Superb Fairy-wren	<i>Malurus cyaneus</i>	252	29-Sep-15			
White-browed Scrubwren	<i>Sericornis frontalis</i>	250	29-Sep-15			
Striated Fieldwren	<i>Calamanthus fuliginosus</i>	33	26-Jun-15			
Yellow Thornbill	<i>Acanthiza nana</i>	6	20-Jul-06			
Yellow-rumped Thornbill	<i>Acanthiza chrysorrhoa</i>	203	31-Aug-15			
Brown Thornbill	<i>Acanthiza pusilla</i>	245	29-Sep-15			
Spotted Pardalote	<i>Pardalotus punctatus</i>	205	29-Sep-15			
Striated Pardalote	<i>Pardalotus striatus</i>	31	30-Apr-15			
Eastern Spinebill	<i>Acanthorhynchus tenuirostris</i>	35	23-Jul-15			
Yellow-faced Honeyeater	<i>Lichenostomus chrysops</i>	21	28-Apr-14			
White-eared Honeyeater	<i>Lichenostomus leucotis</i>	11	30-Apr-15			
White-plumed Honeyeater	<i>Lichenostomus penicillatus</i>	251	29-Sep-15			
Noisy Miner	<i>Manorina melanocephala</i>	178	29-Sep-15			
Spiny-cheeked Honeyeater	<i>Acanthagenys rufogularis</i>	43	30-Apr-15			
Little Wattlebird	<i>Anthochaera chrysoptera</i>	248	29-Sep-15			

Common Name	Scientific Name	No of sightings from 252 surveys	Most recent record	EPBC	FFG	DSE
Red Wattlebird	<i>Anthochaera carunculata</i>	250	29-Sep-15			
White-fronted Chat	<i>Epthianura albifrons</i>	164	29-Sep-15			
Scarlet Honeyeater	<i>Myzomela sanguinolenta</i>	1	30-Oct-09			
New Holland Honeyeater	<i>Phylidonyris novaehollandiae</i>	233	29-Sep-15			
White-naped Honeyeater	<i>Melithreptus lunatus</i>	27	28-Apr-14			
Black-faced Cuckoo-shrike	<i>Coracina novaehollandiae</i>	199	29-Sep-15	M		
Crested Shrike-tit	<i>Falcunculus frontatus</i>	48	20-Feb-15			
Golden Whistler	<i>Pachycephala pectoralis</i>	50	23-Jul-15			
Rufous Whistler	<i>Pachycephala rufiventris</i>	4	25-Mar-09			
Grey Shrike-thrush	<i>Colluricincla harmonica</i>	213	29-Sep-15			
Olive-backed Oriole	<i>Oriolus sagittatus</i>	5	26-Nov-10			
Dusky Woodswallow	<i>Artamus cyanopterus</i>	126	29-Sep-15			
Grey Butcherbird	<i>Cracticus torquatus</i>	212	29-Sep-15			
Australian Magpie	<i>Cracticus tibicen</i>	251	29-Sep-15			
Grey Currawong	<i>Strepera versicolor</i>	1	26-Jul-10			
Rufous Fantail	<i>Rhipidura rufifrons</i>	2	25-Mar-15	Mi,M		
Grey Fantail	<i>Rhipidura albiscapa</i>	224	29-Sep-15			
Willie Wagtail	<i>Rhipidura leucophrys</i>	248	29-Sep-15			
Australian Raven	<i>Corvus coronoides</i>	1	02-Jul-00			
Little Raven	<i>Corvus mellori</i>	223	29-Sep-15	M		
Magpie-lark	<i>Grallina cyanoleuca</i>	250	29-Sep-15			
Scarlet Robin	<i>Petroica boodang</i>	10	25-Jul-13			
Red-capped Robin	<i>Petroica goodenovii</i>	1	21-Mar-14			
Flame Robin	<i>Petroica phoenicea</i>	113	31-Aug-15	M		
Pink Robin	<i>Petroica rodinogaster</i>	2	27-Jun-12	M		
Eastern Yellow Robin	<i>Eopsaltria australis</i>	220	29-Sep-15			
*Eurasian Skylark	<i>Alauda arvensis</i>	234	29-Sep-15			
Golden-headed Cisticola	<i>Cisticola exilis</i>	250	29-Sep-15			
Australian Reed-Warbler	<i>Acrocephalus australis</i>	151	29-Sep-15	Mi,M		

Common Name	Scientific Name	No of sightings from 252 surveys	Most recent record	EPBC	FFG	DSE
Tawny Grassbird	<i>Megalurus timoriensis</i>	1	21-Mar-14			
Little Grassbird	<i>Megalurus gramineus</i>	245	29-Sep-15			
Rufous Songlark	<i>Cincloramphus mathewsi</i>	6	29-Aug-12			
Brown Songlark	<i>Cincloramphus cruralis</i>	3	29-Aug-12			
Silvereeye	<i>Zosterops lateralis</i>	235	29-Sep-15	M		
Welcome Swallow	<i>Hirundo neoxena</i>	246	29-Sep-15			
Fairy Martin	<i>Petrochelidon ariel</i>	39	25-Mar-15			
Tree Martin	<i>Petrochelidon nigricans</i>	28	31-Dec-14	M		
*Common Blackbird	<i>Turdus merula</i>	252	29-Sep-15			
*Song Thrush	<i>Turdus philomelos</i>	21	09-Oct-01			
*Common Starling	<i>Sturnus vulgaris</i>	250	29-Sep-15			
*Common Myna	<i>Sturnus tristis</i>	246	29-Sep-15			
Mistletoebird	<i>Dicaeum hirundinaceum</i>	67	23-Jul-15			
Red-browed Finch	<i>Neochmia temporalis</i>	97	29-Sep-15			
*House Sparrow	<i>Passer domesticus</i>	220	29-Sep-15			
*Eurasian Tree Sparrow	<i>Passer montanus</i>	37	25-Mar-08			
Australasian Pipit	<i>Anthus novaeseelandiae</i>	47	27-Jan-15	M		
*European Goldfinch	<i>Carduelis carduelis</i>	248	29-Sep-15			
*Common Greenfinch	<i>Chloris chloris</i>	107	29-May-15			
*Domestic Duck	<i>Anas platyrhynchos domesticus</i>	82	02-Jul-03			
*Black Duck-Mallard hybrid	<i>Anas superciliosa – Anas platyrhynchos</i>	2	27-Jan-15			

Appendix 4 List of threatened and/or migratory species recorded at Edithvale Wetlands from 1989 to 2015 and at Seaford Wetlands from 1994 to 2015 during monthly surveys undertaken by Bird Life Australia (summary of surveys from BirdLife Australia Atlas database).

Key:

- * denotes introduced species
- X Recorded
- EPBC *Environment Protection and Biodiversity Conservation Act 1999*
 - CE Listed as Critically Endangered under the EPBC Act
 - EN Listed as Endangered under the EPBC Act
 - VU Listed as Vulnerable under the EPBC Act
- Mi Migratory
- M Marine overfly
- FFG *Flora and Fauna Guarantee Act 1988*
 - L Listed as threatened under the FFG Act
 - N Nominated for listing under the FFG Act
- DSE Status according to DSE (2013): Advisory List of Threatened Vertebrate Fauna in Victoria – 2013:
 - ce classified as critically endangered in DSE (2013)
 - en classified as endangered in DSE (2013)
 - vu classified as vulnerable in DSE (2013)
 - nt classified as near threatened in DSE (2013)
 - dd classified as data deficient in DSE (2013)

Common Name	Scientific Name	EPBC	FFG	DSE	Edithvale North	Edithvale South	Seaford
Stubble Quail	<i>Coturnix pectoralis</i>	M			X	X	X
Magpie Goose	<i>Anseranas semipalmata</i>	M	L	nt	X	X	
Musk Duck	<i>Biziura lobata</i>	M		vu	X	X	X
Freckled Duck	<i>Stictonetta naevosa</i>		L	en	X	X	X
Cape Barren Goose	<i>Cereopsis novaehollandiae</i>	M				X	X
Australasian Shoveler	<i>Anas rhynchotis</i>			vu	X	X	X
Hardhead	<i>Aythya australis</i>			vu	X	X	X
Blue-billed Duck	<i>Oxyura australis</i>		L	en	X	X	X
Short-tailed Shearwater	<i>Ardenna tenuirostris</i>	Mi,M				X	
Fork-tailed Swift	<i>Apus pacificus</i>	Mi,M				X	
White-throated Needletail	<i>Hirundapus caudacutus</i>	Mi,M		vu			X
Pied Cormorant	<i>Phalacrocorax varius</i>			nt	X	X	
Australian Pelican	<i>Pelecanus conspicillatus</i>	M			X	X	X
Australasian Bittern	<i>Botaurus poiciloptilus</i>	EN	L	en	X	X	X
Australian Little Bittern	<i>Ixobrychus dubius</i>		L	en	X	X	X
Eastern Great Egret	<i>Ardea modesta</i>	Mi,M	L	vu	X	X	X

Common Name	Scientific Name	EPBC	FFG	DSE	Edithvale North	Edithvale South	Seafood
Intermediate Egret	<i>Ardea intermedia</i>	M	L	en	X	X	X
Cattle Egret	<i>Ardea ibis</i>	Mi,M			X	X	X
Little Egret	<i>Egretta garzetta</i>	M	L	en	X	X	X
Nankeen Night-Heron	<i>Nycticorax caledonicus</i>	M		nt	X	X	X
Glossy Ibis	<i>Plegadis falcinellus</i>	Mi,M		nt	X	X	X
Australian White Ibis	<i>Threskiornis molucca</i>	M			X	X	X
Straw-necked Ibis	<i>Threskiornis spinicollis</i>	M			X	X	X
Royal Spoonbill	<i>Platalea regia</i>			nt	X	X	X
White-bellied Sea-Eagle	<i>Haliaeetus leucogaster</i>	Mi,M	L	vu		X	X
Whistling Kite	<i>Haliastur sphenurus</i>	M			X	X	X
Brown Goshawk	<i>Accipiter fasciatus</i>	M			X	X	X
Spotted Harrier	<i>Circus assimilis</i>			nt		X	
Swamp Harrier	<i>Circus approximans</i>	M			X	X	X
Nankeen Kestrel	<i>Falco cenchroides</i>	Mi,M			X	X	X
Lewin's Rail	<i>Lewinia pectoralis</i>		L	vu	X	X	X
Buff-banded Rail	<i>Gallirallus philippensis</i>	M			X	X	X
Baillon's Crake	<i>Porzana pusilla</i>	M	L	vu	X	X	X
Spotless Crake	<i>Porzana tabuensis</i>	M			X	X	X
Black-winged Stilt	<i>Himantopus himantopus</i>	Mi,M			X	X	X
Red-necked Avocet	<i>Recurvirostra novaehollandiae</i>	Mi,M			X	X	X
Red-capped Plover	<i>Charadrius ruficapillus</i>	,M			X	X	X
Double-banded Plover	<i>Charadrius bicinctus</i>	Mi,M			X		X
Australian Painted Snipe	<i>Rostratula australis</i>	CE; Mi,M	L	ce		X	
Latham's Snipe	<i>Gallinago hardwickii</i>	Mi,M	N	nt	X	X	X
Little Curlew	<i>Numenius minutus</i>	Mi,M				X	
Black-tailed Godwit	<i>Limosa limosa</i>	Mi,M		vu			X
Common Greenshank	<i>Tringa nebularia</i>	Mi,M		vu	X	X	X
Marsh Sandpiper	<i>Tringa stagnatilis</i>	Mi,M		vu	X	X	X
Wood Sandpiper	<i>Tringa glareola</i>	Mi,M		vu	X	X	X
Red Knot	<i>Calidris canutus</i>	Mi,M		en	X		
Red-necked Stint	<i>Calidris ruficollis</i>	Mi,M			X	X	X

Common Name	Scientific Name	EPBC	FFG	DSE	Edithvale North	Edithvale South	Seaford
Long-toed Stint	<i>Calidris subminuta</i>	Mi,M		nt	X	X	X
Pectoral Sandpiper	<i>Calidris melanotos</i>	Mi,M		nt	X	X	X
Sharp-tailed Sandpiper	<i>Calidris acuminata</i>	Mi,M			X	X	X
Curlew Sandpiper	<i>Calidris ferruginea</i>	CE,Mi,M		en	X	X	X
Red-necked Phalarope	<i>Phalaropus lobatus</i>	Mi,M					X
Caspian Tern	<i>Hydroprogne caspia</i>	Mi,M	L	nt		X	X
Whiskered Tern	<i>Chlidonias hybrida</i>	M		nt	X	X	X
White-winged Black Tern	<i>Chlidonias leucopterus</i>	Mi,M		nt	X	X	
Common Tern	<i>Sterna hirundo</i>	Mi,M				X	X
Crested Tern	<i>Thalasseus bergii</i>	M				X	
Pacific Gull	<i>Larus pacificus</i>	M			X	X	X
Silver Gull	<i>Chroicocephalus novaehollandiae</i>	M			X	X	X
Swift Parrot	<i>Lathamus discolor</i>	En; M	L	en			X
Blue-winged Parrot	<i>Neophema chrysostoma</i>	M			X	X	X
Horsfield's Bronze-Cuckoo	<i>Chalcites basalis</i>	M			X	X	X
Black-eared Cuckoo	<i>Chalcites osculans</i>	M				X	
Shining Bronze-Cuckoo	<i>Chalcites lucidus</i>	M					X
Pallid Cuckoo	<i>Cacomantis pallidus</i>	M			X	X	X
Fan-tailed Cuckoo	<i>Cacomantis flabelliformis</i>	M			X	X	X
Southern Boobook	<i>Ninox novaeseelandiae</i>	M				X	
Sacred Kingfisher	<i>Todiramphus sanctus</i>	M			X	X	X
Black-faced Cuckoo-shrike	<i>Coracina novaehollandiae</i>	M			X	X	X
Rufous Fantail	<i>Rhipidura rufifrons</i>	Mi,M					X
Little Raven	<i>Corvus mellori</i>	M			X	X	X
Satin Flycatcher	<i>Myiagra cyanoleuca</i>	Mi,M				X	
Flame Robin	<i>Petroica phoenicea</i>	M			X	X	X
Pink Robin	<i>Petroica rodinogaster</i>	M					X
Australian Reed-Warbler	<i>Acrocephalus australis</i>	Mi,M			X	X	X
Silvereye	<i>Zosterops lateralis</i>	M			X	X	X
Tree Martin	<i>Petrochelidon nigricans</i>	M			X	X	X
Australasian Pipit	<i>Anthus novaeseelandiae</i>	M			X	X	X

Appendix 5 Non-avian fauna species recorded within the Edithvale-Seafood Wetlands Ramsar Site (see Ecology Australia 2001; KBR 2009; DELWP 2016).

Key:

- * denotes introduced species
- EPBC *Environment Protection and Biodiversity Conservation Act 1999*
 - CE Listed as Critically Endangered under the EPBC Act
 - EN Listed as Endangered under the EPBC Act
 - VU Listed as Vulnerable under the EPBC Act
 - Mi Migratory
 - M Marine overfly
- FFG *Flora and Fauna Guarantee Act 1988*
 - L Listed as threatened under the FFG Act
 - N Nominated for listing under the FFG Act
- DSE Status according to DSE (2013): Advisory List of Threatened Vertebrate Fauna in Victoria – 2013:
 - ce classified as critically endangered in DSE (2013)
 - en classified as endangered in DSE (2013)
 - vu classified as vulnerable in DSE (2013)
 - nt classified as near threatened in DSE (2013)
 - dd classified as data deficient in DSE (2013)

Scientific Name	Common Name	DSE 2013	EPBC	FFG
Mammals				
<i>Tachyglossus aculeatus</i>	Short-beaked Echidna			
<i>Antechinus agilis</i>	Agile Antechinus			
<i>Isodon obesulus obesulus</i>	Southern Brown Bandicoot	nt	EN	L
<i>Trichosurus vulpecula</i>	Common Brushtail Possum			
<i>Pseudocheirus peregrinus</i>	Common Ringtail Possum			
<i>Petaurus breviceps</i>	Sugar Glider			
<i>Macropus giganteus</i>	Eastern Grey Kangaroo			
<i>Wallabia bicolor</i>	Swamp Wallaby			
<i>Pteropus poliocephalus</i>	Grey-headed Flying-fox	vu	VU	L
<i>Tadarida australis</i>	White-striped Freetail Bat			
<i>Nyctophilus geoffroyi</i>	Lesser Long-eared Bat			
<i>Scotorepens orion</i>	Eastern Broad-nosed Bat			
<i>Chalinolobus gouldii</i>	Gould's Wattled Bat			
<i>Chalinolobus morio</i>	Chocolate Wattled Bat			
<i>Vespadelus darlingtoni</i>	Large Forest Bat			
<i>Vespadelus regulus</i>	Southern Forest Bat			
<i>Vespadelus vulturnus</i>	Little Forest Bat			
<i>Rattus fuscipes</i>	Bush Rat			
<i>Rattus lutreolus</i>	Swamp Rat			
<i>Hydromys chrysogaster</i>	Water Rat (Rakali)			
* <i>Rattus rattus</i>	Black Rat		*	
* <i>Rattus norvegicus</i>	Brown Rat		*	
* <i>Mus musculus</i>	House Mouse		*	
* <i>Oryctolagus cuniculus</i>	European Rabbit		*	

Scientific Name	Common Name	DSE 2013	EPBC	FFG
* <i>Lepus capensis</i>	Brown Hare			
* <i>Felis catus</i>	Cat		*	
* <i>Vulpes vulpes</i>	European Fox			
* <i>Canis lupus familiaris</i>	Dog		*	
Reptiles				
<i>Chelodina longicollis</i>	Eastern Snake-necked Turtle	dd		
<i>Christinus marmoratus</i>	Marbled Gecko			
<i>Pygopus lepidopus</i>	Common Scaly-foot			
<i>Amphibolurus muricatus</i>	Tree Dragon			
<i>Acritoscincus duperreyi</i>	Eastern Three-lined Skink			
<i>Liopholis whitii</i> GROUP	White's Skink			
<i>Lampropholis delicata</i>	Delicate Skink			
<i>Eulamprus tympanum tympanum</i>	Southern Water Skink			
<i>Lampropholis guichenoti</i>	Garden Skink			
<i>Niveoscincus metallicus</i>	Metallic Skink			
<i>Saproscincus mustelinus</i>	Weasel Skink			
<i>Lerista bougainvillii</i>	Bougainville's Skink			
<i>Pseudemoia entrecasteauxii</i>	Southern Grass Skink			
<i>Tiliqua nigrolutea</i>	Blotched Blue-tongued Lizard			
<i>Tiliqua scincoides</i>	Common Blue-tongued Lizard			
<i>Notechis scutatus</i>	Tiger Snake			
<i>Pseudechis porphyriacus</i>	Red-bellied Black Snake			
<i>Pseudonaja textilis</i>	Eastern Brown Snake			
<i>Austrelaps superbus</i>	Lowland Copperhead			
<i>Drysdalia coronoides</i>	White-lipped Snake			
<i>Suta flagellum</i>	Little Whip Snake			
Frogs				
<i>Limnodynastes dumerilii</i>	Southern Bullfrog (ssp. unknown)			
<i>Limnodynastes peronii</i>	Striped Marsh Frog			
<i>Limnodynastes tasmaniensis</i>	Spotted Marsh Frog (race unknown)			
<i>Neobatrachus sudellae</i>	Common Spadefoot Toad			
<i>Paracrinia haswelli</i>	Haswell's Froglet			
<i>Crinia signifera</i>	Common Froglet			
<i>Pseudophryne semimarmorata</i>	Southern Toadlet	vu		
<i>Litoria ewingii</i>	Southern Brown Tree Frog			
<i>Litoria raniformis</i>	Growling Grass Frog	en	Vu	L
<i>Litoria verreauxii verreauxii</i>	Verreaux's Tree Frog			
Fish				
<i>Anguilla australis</i>	Southern Shortfin Eel			
<i>Galaxias truttaceus</i>	Spotted Galaxias			

Scientific Name	Common Name	DSE 2013	EPBC	FFG
<i>Galaxias brevipinnis</i>	Climbing Galaxias			
<i>Galaxias maculatus</i>	Common Galaxias			
* <i>Gambusia holbrooki</i>	Eastern Gambusia			

Appendix 6 Details of microbat surveys undertaken by Parson Brinckerhoff (2009 and 2010), Edithvale-Seaford Wetlands Ramsar Site.

Key:

- X Positively identified
 P Potential call detection
 FFG *Flora and Fauna Guarantee Act 1988*
 L Listed as threatened under the FFG Act
 DSE Status according to DSE (2013): Advisory List of Threatened Vertebrate Fauna in Victoria – 2013:
 vu classified as vulnerable in DSE (2013)
 dd classified as data deficient in DSE (2013)

Common name	Species name	FFG	DSE 2013	Edithvale Wetlands	Seaford Wetlands
Chocolate Wattled Bat	<i>Chalinolobus morio</i>			X	
Gould's Wattled Bat	<i>Chalinolobus gouldii</i>			X	X
Large Forest Bat	<i>Vespadelus darlingtoni</i>			X	X
Southern Forest Bat	<i>Vespadelus regulus</i>			X	X
Little Forest Bat	<i>Vespadelus vulturinus</i>			X	
Southern Freetail Bat	<i>Mormopterus</i> sp 4			X	X
Eastern Freetail Bat / Southern Freetail Bat	<i>Mormopterus</i> sp 2 / <i>Mormopterus</i> sp 4			X	X
White-striped Freetail Bat	<i>Tadarida australis</i>			X	X
Lesser Long-eared Bat	<i>Nyctophilus geoffroyi</i>			X	
Gould's Wattled Bat/Mormopterus sp.	<i>Chalinolobus gouldii</i> / <i>Mormopterus</i> sp.				X
Large Forest Bat/ Eastern Bent-wing Bat	<i>Vespadelus darlingtoni</i> / <i>Miniopterus schreibersii</i> <i>oceanensis</i>				X
Forest Bat spp.	<i>Vespadelus</i> spp.			X	X
Long-eared Bat spp.	<i>Nyctophilus</i> spp.			X	X
Eastern Bent-wing Bat	<i>Miniopterus schreibersii</i> <i>oceanensis</i>	L	vu	P	P
Eastern Broad-nosed Bat	<i>Scotorepens orion</i>			P	P
Yellow-bellied Sheathtail Bat	<i>Saccolaimus flaviventris</i>	L	dd		P

Appendix 7 Hydrology of wetland cells at Edithvale-Seaford Wetlands (see Section 6.3).

Table 17 Hydrology of the wetland cells at Edithvale Wetlands (summarised from GHD 2006, KBR 2009, SKM 2011, DSE 2012 and Jacobs 2016 and updated from data provided by Paul Rees, Melbourne Water).

Hydrological Cell	Water sources	Water Discharges	Morphology	Watering regime	Habitat
Edithvale South					
ES1 (Main Wetland)	Three drains from the developed urban catchment to the east (via sediment ponds)	To Edithvale North (EN1) via a siphon under Edithvale Road when water levels reach 0.00 m AHD	Natural dish-shaped cross-section underlain by a thick layer of peat which thins towards the edges.	Inundated in winter and spring (standing water level generally sits at 0.00 m AHD, but rises to 0.78 m AHD during rain events)	Shallow and deep fresh-brackish marsh
	Overland flows	During rain events discharges to Edithvale North Wetlands at Centre Swamp Drain when water levels are above the invert level of the weir (-0.02m AHD)	Depth is normally c. 0.45 m (the deepest part is 0.3 m below sea level and water level is usually 0.2 m AHD at peak level, but up to 1.25 m AHD in a 1 in 100 year average return interval (ARI) event (Paul Rees, Melbourne Water, pers. comm.).	Drawdown and drying occurs in summer and autumn to below 0.00 m AHD	Drawdown during summer and autumn creates critical mudflat foraging habitat for migratory shorebirds
	Overflows from Centre Swamp Drain during storms with 1 in 2 year annual return interval (ARI)			Maximum drying is generally by late-January, but may occur earlier during drought conditions	Supports a heavy growth of Salt/Marsh Club-rush (<i>Bolboschoenus caldwellii</i>) in late-summer and autumn – the key to maintain peats and nutrient cycling and control <i>Typha</i> invasion
	Flood overflows from Centre Swamp Drain are generally controlled by inflows into Edithvale North (EN1)			Unseasonal event flows may partially or totally rewet the area	
ES1a, ES1b and ES1c (Drought Refuges)	ES1a and ES1b - pumping from Centre Swamp Drain during prolonged dry periods (enters via ES1a and overflows to ES1b) in addition to the sources to ES1 (Main Wetland)	None known	Excavated pools are deeper than the main wetland (ES1)	Inundated in winter and spring (standing water level generally at 0.00 m AHD), but rises to 0.78 m AHD during rain events)	Drought refuges support permanent water, except during drought
	ES1c - Groundwater in addition to the sources to ES1 (Main Wetland)			Generally dry out every year, except during unusually wet years (level will sit at -0.40 m AHD or lower depending on rainfall)	
Edithvale North					
EN1	Mostly from ES1 via a siphon crest in ES1 at -0.01 m AHD under Edithvale Road	To EN2 at Weir 1 at -0.15 m AHD	Constructed within the former floodplain	Generally inundated in winter and spring (level will sit at 0.00 m AHD or below)	Fresh-brackish
	Also via Weir 5 from Centre Swamp Drain at 0.2 m AHD	To EN2 via overflow at -0.13 m AHD	Shallow, peat-lined cell with concentric marsh zones and an island in the centre	Drying in summer and autumn	Supports a heavy growth of Salt/Marsh Club-rush (<i>Bolboschoenus caldwellii</i>) in late-summer and autumn
	Leakage from groundwater			Generally dries-up every year, but some wet summers will result in a residual pool in autumn	The management objective is to provide habitat for waders, dabbling and filter-feeding waterbird species
				A moist area is maintained during dry periods by groundwater	
EN2	From EN1 at Weir 1 at -0.15 m AHD	To EN3 at Weir 2 at -0.37 m AHD, but is dependent on the ponded level in EN3; excess flows can be directed to the Dog Pond	Constructed within the former floodplain and reach deep into the sandy substrates that underlie the area	During winter and spring, will fill from EN1 – level is determined by level in EN3 with a minimum of -0.40 m AHD	Fresh-brackish
			The invert level of EN2 is -1.95 m AHD		
			Data for EN2 suggest that it ranges in depth from 0.00 m (i.e. empty) (-1.95 m AHD) recorded in Feb 2016, to more than 2.00 m (gauge under water) (>0.15 m AHD) (Paul Rees, Melbourne Water, pers. comm.).		

Hydrological Cell	Water sources	Water Discharges	Morphology	Watering regime	Habitat
	Groundwater	To Dog Pond via a high level overflow channel at c. 0.4 m AHD	A series of weirs between EN3, EM3a and EN2 limit the total drawdown under prolonged dry or drought conditions	Drawdown generally to c. -0.5 m AHD during average years, but may drawdown to -1.2 m AHD during drought years	
				System responds to groundwater in summer which is controlled by area saturation and drawdown to invert low flow pipe in Centre Swamp Drain floodway to south	
EN3 (main open water pond)	From EN2 at Weir 2 at -0.37 m AHD	To Centre Swamp Drain via two 150 mm outlet pipes	The invert level of EN3 is -1.9 m AHD	Will fill to -0.2 to 0.3 m in wet years, but may be much lower during low rainfall'	EN3 is the main open water pond
	From EN3a via Weir 3 at -0.23 m AHD	To Centre Swamp Drain via a high level overflow channel (overflows top of outlets to be set at -0.2 to 0.3 m AHD)	Data for EN3 suggest that it ranges in depth from 0.42 m (-1.48 m AHD) to 2.00 m (-0.1 m AHD) (Paul Rees, Melbourne Water, pers. comm.)	Filling of pond to full supply level generally only occurs in July-October	EN3 water levels are responsive to groundwater. Levels can be -1.2 m AHD during drought years with salinity levels of 10,000-12,000 us/cm, or -0.2-0.4 m AHD in wetter periods with salinities of 4,000-5,000 us/cm.
	From EN4 via Weir 3 at -0.98 m AHD			Drawdown generally to c. -0.5 m AHD during average years, but may drawdown to -1.2 m AHD during drought years	
	From groundwater			During summer, is occasionally filled from subdivision stormwater and overflow from EN1 and EN2	
EN3a	From Stormwater drains to the north via litter traps and Sediment Ponds E4 and E5	To EN3 via Weir 3 at -0.23 m AHD and Weir 4 at -0.98 m AHD		Filling of pond to full supply level generally only occurs in July-October	EN3a water levels are responsive to groundwater. Levels can be -1.2 m AHD during drought years with salinity levels of 10,000-12,000 us/cm, or -0.2-0.4 m AHD in wetter periods with salinities of 4,000-5,000 us/cm.
	From stormwater from an overland flow-path via litter traps and Sediment Ponds EN4 and EN5			Drawdown via groundwater and evaporation response generally to c. -0.5 m AHD during average years, but may drawdown to -1.2 m AHD during drought years	
	From groundwater		Water levels in EN3 and EN3a are generally responsive to the groundwater table, which varies in height, but can draw down to -1.95 m AHD during drought years. Salinity is therefore a problem in EN3 and EN3a e.g. 10,000 to 12,000 us/cm (GHD 2006). In wetter seasons, the standing water can reach 0.00 to 0.15 m AHD and salinity is lower e.g. 4,000 to 5,000 us/cm (GHD 2006).		
Dog Pond	Water level is controlled by groundwater	None	Constructed within the former floodplain	Relies on inflows from EN2 to be above the standing groundwater height	
	Dog Pond can receive water from EN2 via a high level overflow channel at c. -0.4 m AHD		Deep, reaching into the sandy substrates that underlie the area	Generally dry by late-January	
EN4 (Sediment Pond)	Stormwater from stormwater drains to the north	To EN3a		During winter and spring, the water level and overflow is controlled by a sill at c. -0.2 m AHD	Deep pool generally supports freshwater and provides good habitat for Blue-billed Ducks, Musk Ducks and Australasian Bittern
	Overland flow			Drawdown to the weir at -0.2 m AHD, and then evaporation and inflows control the level	

Hydrological Cell	Water sources	Water Discharges	Morphology	Watering regime	Habitat
EN5 (Sediment Pond)	Stormwater from stormwater drains to the north	To EN3a		During winter and spring, water levels respond to main pond height in EN3	
	Overland flow			Drawdown occurs during summer and is generally dry by late-December	
EC1 and EC2 (Edithvale Common Wetlands)	EC1 – Ephemeral Wetland fills to 0.1 m AHD from a diversion weir in the outfall pipeline from Centre Swamp Drain	None		During winter and spring, the ephemeral wetland fills to 0.1 m AHD from a diversion weir in the outfall pipeline from Centre Swamp Drain	The ephemeral wetland provided good habitat for Latham's Snipe and duck species prior to improvements to drainage (Will Steele, Melbourne Water, pers. comm.)
				The wetland is generally dry by the end of November	
	EC2 – Shallow depression is only filled by local run-off	None	Shallow depression located to the east of the "Duck Inn"	Only fills in wet years	The ephemeral wetland provides good Latham's Snipe and frog habitat
				The wetland is dry by the end of November	

Table 18 Hydrology of the wetland cells at Seaford Wetlands (summarised from GHD 2006, KBR 2009, SKM 2011, DSE 2012 and Jacobs 2016).

Hydrological Cell	Water sources	Water Discharges	Morphology	Watering regime	Habitat
Seaford North					
SN1 (Main Header Pool)	Rising main transfers base flows from Wadsleys Road Drain to the north	Discharges via a central pipe (c. 600 mm) which can feed into either SCW1 or SCE2. variable level regulator - 0.145m AHD to 0.8m AHD	Standing water levels during winter and spring are generally at 0.6 m AHD and are controlled via overflow orifices in earth bund	Standing water levels during winter and spring are fill and maintain at 0.6 m AHD	Fresh-brackish
	The centre drain (old Seaford Drain) runs from north to south through the site and carries inflows to SN1 through higher ground (also see Aurecon 2011). Gates enable the direction of water flow to the east or west.		Level of side caste overflow structure in east is at 0.80 m AHD and/or sag point in west is at 0.80 m AHD (levee was upgraded in 2013, Paul Rees, pers. comm.).	Summer water levels are at c. 0.45 m AHD	Maintains permanent water
		Overflows to the west into SCW1		Summer-autumn maintenance flows occur via Wadsleys Drain	
		Overflows to the east into SCE2			
		System is manually operated to redirect and discharge flows to a centre drain after November to allow for drying			
SN2	Local rainfall	None	Ephemeral wetlands that form part of the areas original morphology- ephemeral areas are blocked by landforms of the former swamp that are disrupted by Eel Race Drain	Ephemeral wetlands wet during winter and spring Generally dry during summer and autumn	Ephemeral wetlands providing important habitat for waders and insectivorous birds in late-spring
*SN Downs Land	Local rainfall	None	Seasonal wetlands in a series of old watercourses	Ephemeral wetlands only fill during a wet winter	Ephemeral wetlands with potential to be managed for late-winter and early-spring habitat for waders
Seaford Central					
SCW1	Stormwater fed by Seaford North local drainage system to the west via undiverted drains	Discharges to the centre drain	Deeper substrate under shallow peats was disturbed historically during extensive cropping	SCW1 is wetter than natural due to the drainage inputs, cessation of the overall through drainage, a lack of saline groundwater interception on the drains from	Long-term running of freshwater has resulted in excessive growth of Common Reed,

Hydrological Cell	Water sources	Water Discharges	Morphology	Watering regime	Habitat
				Seaford North and the long-term running of freshwater through the pool in summer months	Cumbungi and Spiny Rush
	The surrounding levee is not totally impermeable and therefore the drain is not fully regulated; it overtops to the east when water levels reach 14 cm in height and to the west at 12 cm (Paul Rees, Melbourne Water, pers. comm.).		SCW1 has an invert of 0.14 m AHD, with a range of 0.00m (0.14 m AHD) to 0.70 m (0.84 m AHD)	SCW1 and SCE2 operate independently below 0.14 m, but are connected above this point as essentially one large pool, as the levee on either side of the main drain has low points of 0.28 m AHD on the west and 0.31 m AHD on the east. In an average year, SCW1 and SCE2 will be disconnected initially in autumn, they will slowly fill-up until they reach 0.31 m AHD (0.12 to 0.14 m deep) at which point they will become connected over most of the winter and spring, until in late-spring and early-summer, they draw down below 0.31 m AHD and are no longer connected.	
	Overflow sill from SN1 at 0.80 m AHD		The western margin of the cell was historically filled for subdivision with a steep interface between the swamp and fill	Water levels in winter and spring sit at c. 0.4 m AHD and are controlled at two locations by sills to the centre drain, but will pond higher than this level during flood events	
	From the central pipe (c. 600 mm) from SN1				
	Groundwater				
SCE2	Overflow from SN1 at 0.8 m AHD	Overflows via a sill in the south corner of the cell to SCE2a at c. 0.7 m AHD	Is the least disturbed of the cells and supports original morphology	Supports a relatively natural wetting and drying regime and dries in summer and autumn with exception of residual pools and unseasonal events	Management regime aims to reflect a natural cycle
	From the central pipe (c. 600 mm) from SN1		SCE2 has an invert of 0.19 m AHD, with a range of 0.00m (0.19 m AHD) to 0.60m (0.79 m AHD) and generally dries out in summer.	SCW1 and SCE2 operate independently below 0.14 m, but are connected above this point as essentially one large pool, as the levee on either side of the main drain has low points of 0.28 m AHD on the west and 0.31 m AHD on the east. In an average year, SCW1 and SCE2 will be disconnected initially in autumn, they will slowly fill-up until they reach 0.31 m AHD (0.12 to 0.14 m deep) at which point they will become connected over most of the winter and spring, until in late-spring and early-summer, they draw down below 0.31 m AHD and are no longer connected.	Important habitat for waders with a large productive mudflat available from September to December
			Peat layers are predominantly intact	Water levels pond to 0.45 m AHD in late-spring following which supply is halted, except for unseasonal event inflows.	Supports a heavy growth of Marsh Club-rush (<i>Bolboschoenus caldwellii</i>) in late-summer and autumn
			Water levels are controlled by sills in the southwest corner adjacent to the centre drain	Water levels should desiccate to mudflats by mid-December (inflows from SN1 cease by late-October). Unseasonal flows could refill the cell.	
SCE2a	From SCE2	Overflows to SSE4	This cell is part of SCE2, but is separated from it by a low level weir on the northern margin	Minor ponding occurs to 0.55 m AHD during winter due to the old fill mound of the centre drain	Shallow brackish wetland
	The surrounding levee is not totally impermeable and therefore the drain is not fully regulated; it overtops to the east when water levels reach 14 cm in height and to the west at 12 cm (Paul Rees, Melbourne Water, pers. comm.).	Overflows to SSW1	Is the least disturbed of the cells and supports original morphology	Separation from SCE2 by the low level weir results in draining and drying separately in late-spring	
			Peat layers are predominantly intact	Generally dry in summer and autumn, except for some residual pool areas in natural landform	

Hydrological Cell	Water sources	Water Discharges	Morphology	Watering regime	Habitat
Seaford South					
SSW1	Numerous inlets in the form of high level surcharges from local drains to the west (most smaller events do not surcharge because they are intercepted by the James Street Pump Station system)	Via the Austin Road outlet (water levels are controlled by an outlet orifice in the southeast corner of SSW1 which allows pondage up to 0.3 m AHD or drainage down to 0.1 m AHD and is connected to the Austin Road outlet)	Constructed within the former floodplain in 1989 and is the lowest point in the Seaford system	During winter and spring, water levels are generally maintained at 0.18 – 0.35 m AHD by the outlet orifice to Austin Road outlet	Highly saline and impacted by acid sulphate oxidation (low pH)
		A higher level overflow weir to SSW3 on the southern side of the cell is set at 0.5 m AHD	Excavation has intercepted the sub-stratum of the wetland	During summer and autumn, water levels will be maintained at 0.25-0.30 m AHD by base flows	Limited productivity and lower diversity of flora and fauna due to
			Ponds to 0.35 m AHD and drains at 0.1 m AHD		Deep wetland used by a range of diving duck species
			SSW1 has an invert level of 0.08 m AHD and ranges from 0.00 (0.08 m AHD) to 0.68 (0.76 m AHD) m.		A saline water tolerant species of mosquito (<i>Aedes camphotricus</i>) is a risk at higher salinities
SSE2	Fed by local drains to the east	Overflows to SSE4	Perched cell that has been leveed-off from the remainder of the Swamp to receive a fresher water regime in the centre of the Swamp	During winter and spring, water levels can reach 0.7 - 0.8 AHD	Semi-permanent wetland generally supports water of c. 3,000 us/cm, but can be fresher when local drains supply run-off
				In summer, this wetland generally dries to one or two residual pools at 0.3 m AHD	The cell has been colonised by Cumbungi
					The pool supports good habitat, especially for frogs
SSW3	From SSW1 via the overflow at 0.18 – 0.35 m AHD controlled by the outlet orifice to Austin Road outlet	The outlet control occurs at the Austin Road end and deliver water to the local drainage system via the Weatherstone Road Drain	Modified cell	During winter and spring, this cell operates at c. 0.45 m AHD by overflow weir on central pipe	The salinity of water in this cell is highly variable, but generally is above 20,000 us/cm
	From SSW1 via the higher level overflow weir at 0.5 m AHD	Variable crest regulator at 0.045 to 0.55 m AHD	SSW3 has an invert level of 0.21 m AHD and ranges from 0.00 m (0.21m AHD) to 0.66 m (0.87m AHD).	During summer and autumn, water levels hold to a pool height of 0.45 m AHD, or lower depending on event inflows and tidal back flooding	This cell is dominated by Common Reed
	This cell can receive tidal intrusion from Kananook Creek via the Bardia Avenue and Weatherstone Road Drain if the floodgate and pump system fail	If the flood gates on Kananook Creek get blocked, water backs-up and flows back into Seaford Wetlands (Paul Rees, Melbourne Water, pers. comm.)			This cell supports habitat for a variety of waterfowl
SSE4	Receives overflows from SCE2a	To drains in the southwest corner of the cell	Cell has two components: (i) the major component occurs adjacent to Francis Street and Austin Road; and (ii) a smaller component lies on the western side of SSE2	Only wet in winter when will hold to 0.45 -0.50 m AHD, but also responsive to water levels in SSW3	Ephemeral wetland
	From Stormwater drains to the east			During summer and autumn, this cell is generally dry, but will receive major flood inflows from Austin Road	Wetland dominated by halophytes, such as Salt Marsh Rush/Sea Rush (<i>Juncus kraussii</i>)
	Receives major flood inflows from Austin Road				

*Not currently part of the Ramsar site

Appendix 8 The risk assessment for threats at Edithvale-Seafood Wetlands Ramsar Site.

The risk assessment process adopted for this project is consistent with the ISO 31000:2009, Risk management – Principles and guidelines and the Standards Australia Handbook: Environmental risk management - principles and process (HB 203-2000; Standards Australia and Standards New Zealand 2006). The risk assessment approach follows a structured and iterative process, with the following steps:

4. Establish the context – existing values and environmental conditions;
5. Identify risks – threats and associated potential impacts; and
6. Analyse risks – assign likelihoods and consequences to determine level of risk

Establishing the context

A review of existing published and unpublished information relevant to the Edithvale-Seafood Wetlands Ramsar Site was undertaken to identify and summarise the important environmental, social and cultural values; current condition and potential threats to ecological character.

The risk assessment was based on a desktop review of existing information, supplemented by stakeholder knowledge; the latter gained through a stakeholder workshop and site visit.

Identifying risks

The approach uses a hierarchical process to identify potential risks as follows:

- Pressures – activities in the Edithvale-Seafood Wetlands or surrounding area that could affect ecological character
- Stressors – the physical or chemical changes that could arise as a result of an activity
- Effects – the potential responses caused by the stressors.

This allows for clear identification of the underlying causes of risks and threats to ecological character of the Ramsar site, separating the threat from the impact.

Analyse risks

Impact pathways were developed that integrated each level of the hierarchy and these formed the basis of a formal risk analysis process. Likelihood and consequence were assigned to each **impact pathway** in its entirety. See below for an example for an impact pathway:

Pressures	Stressors	Impact
Pollution: Urban stormwater	Increased toxicants	Impacts waterbirds

Likelihood and consequence descriptions used in this assessment are provided in Table 19 and Table 20, respectively, with the risk matrix (Table 21) showing how they combine to score the overall risk. A number of ground rules for the risk assessment were established:

- The risk assessment was focussed on assessing risks to ecological character over the next 15 years.
- In assessing each impact pathway all likely future changes (population, land use, climate change) were considered.
- Where possible all decisions were based on multiple lines of evidence (Table 22).

Table 19 Likelihood

Almost certain	Likely	Possible	Unlikely	Rare
Is expected to occur in most circumstances	Will probably occur in most circumstances	Could occur	Could occur but not expected	Occurs only in exceptional circumstances

Table 20 Consequence

Category	Insignificant	Minor	Moderate	Major	Extreme
Habitat, communities and / or assemblages	Alteration or disturbance to habitat within natural variability. Less than 1% of the area of habitat affected or removed.	1 to 5% of the area of habitat affected in a major way or removed.	5 to 30% of the area of habitat affected in a major way or removed.	30 to 90% of the area of habitat affected in a major way or removed.	Greater than 90% of the area of habitat affected in a major way or removed.
Species and / or groups of species (including protected species)	Population size or behaviour may have changed but it is unlikely that there would be any detectable change outside natural variation / occurrence.	Detectable change to population size and / or behaviour, with no detectable effect on population viability (recruitment, breeding, recovery) or dynamics. Recovery in less than 1 year.	Detectable change to population size and / or behaviour, with no effect on population viability (recruitment, breeding, recovery) or dynamics. Recovery in 1 to 2 years	Detectable change to population size and / or behaviour, with an impact on population viability and or dynamics.	Local extinctions are imminent / immediate or population no longer viable.
Cultural /social	Short-term interruptions in recreational use (days) and perception as a high amenity place unaltered.	Recreational activities restricted and perceptions of amenity altered in a localised area for short-term (weeks)	Recreational activities restricted and perceptions of amenity altered in a localised area for medium term (months).	Long-term disruption to recreational activities and perceptions of amenity altered for 1- 5 years.	Long-term disruption to recreational activities and perceptions of amenity altered for > 5 years.

Table 21 Risk matrix

		Consequence				
Category		Insignificant	Minor	Moderate	Major	Extreme
Likelihood	Almost certain	Negligible	Medium	High	Extreme	Extreme
	Likely	Negligible	Medium	Medium	High	Extreme
	Possible	Negligible	Low	Medium	High	High
	Unlikely	Negligible	Low	Low	Medium	Medium
	Rare	Negligible	Negligible	Negligible	Low	Medium

The risk assessment was undertaken by Jenny Hale (Ramsar specialist) and checked by Andrew McMahon.

Table 22 Risk assessment for the Edithvale-Seaford Wetlands Ramsar Site. Cells highlighted in blue provide a description of the pressure / stressor that is applicable to the relevant impact pathways below

Pressures	Stressors	Impact	Likelihood of impact	Consequence of impact	Risk	Evidence / comments
Pollution: urban stormwater	Increased nutrients and sediments					The Edithvale-Seaford Wetlands are located within a highly urbanised area and the dominant water source for the system is stormwater and drainage water (SKM 2011). Urban water sources are known to be high in nutrient and sediment loads, particularly carried in the first flushes after heavy rainfall (ANZECC and ARMCANZ 2000). Monitoring of water column nutrient concentrations indicates eutrophic conditions at both Seaford and Edithvale Wetlands and periodic high turbidity (> 100 NTU) (Melbourne Water unpublished data). However, this is not surprising for urban wetlands receiving primarily stormwater inflows and there is no indication of a sustained rising trend. In addition, sediments and particulate nutrients are controlled by sediment traps at major drain inflows.
Pollution: urban stormwater	Increased nutrients	Adversely impacts vegetation and habitat	Unlikely	Minor	Low	Temporary Australian wetlands are adapted to periods of high nutrients and the storage of nutrients in the sediment (McComb and Qiu 1998). It is likely that some species of emergent vegetation benefit from nutrient inflows.
Pollution: urban stormwater	Increased nutrients	Adversely impacts waterbird diversity and abundance	Unlikely	Minor	Low	Based on the number of waterbirds supported at highly eutrophic wetlands (e.g. Western Treatment Plant; Menkhorst et al. 2015) it does not seem likely that birds at the site are adversely impacted by increased nutrients and sediments and may even benefit from increased productivity.
Pollution: urban stormwater	Increased nutrients	Adversely impacts waterbird breeding	Unlikely	Minor	Low	As above
Pollution: urban stormwater	Increased nutrients	Adversely impacts threatened species (Australasian bittern and curlew sandpiper)	Unlikely	Minor	Low	As above
Pollution: urban stormwater	Increased nutrients	Adversely affects other fauna	Possible	Minor	Low	Although obligate aquatic species such as fish and amphibians may be more sensitive to changes in water quality. Australian fauna are adapted to temporary wetland systems that exhibit periodic eutrophic conditions.
Pollution: urban stormwater	Increased nutrients	Adversely affects wetland function (drainage and flood mitigation services)	Possible	Minor	Low	Increased sediments and sedimentation may result in a decline in storage capacity at the wetlands. This is currently managed with sediment traps that are periodically emptied to remove sediment.
Pollution: urban stormwater	Increased nutrients	Adversely affects recreation and amenity values	Possible	Minor	Low	Short term increases in odour or unsightly visible algal blooms may impact wetland users for short periods of time.
Pollution: urban stormwater	Increased nutrients	Adversely affects scientific and monitoring values	Rare	Negligible	Negligible	Possibly not even a plausible impact pathway.
Pollution: urban stormwater	Increased nutrients	Adversely affects cultural values	Possible	Minor	Low	Short term increases in odour or unsightly visible algal blooms may impact cultural values for short periods of time.
Pollution: urban stormwater	Toxicants					Studies from elsewhere indicate that urban stormwater and drainage can contain be a source of a number of toxicants such as heavy metals, petroleum derivatives, herbicides and pesticides (Gobbel et al. 2007). Data collected by Melbourne Water from sediments and the water column indicate concentrations of toxicants within ANZECC trigger values, with occasional exceptions for mercury and Total recoverable hydrocarbons. Sediment traps are installed and have been proven effective elsewhere in reducing toxicant loads to urban wetlands (Birch et al. 2005).
Pollution: urban stormwater	Toxicants	Adversely impacts vegetation and habitat	Unlikely	Minor	Low	Concentrations of toxicants are largely within guideline levels and unlikely to affect the growth of wetland vegetation. There is a small likelihood of a spill of herbicides, which could have a measurable, but short term impact.
Pollution: urban stormwater	Toxicants	Adversely impacts waterbird diversity and abundance	Unlikely	Minor	Low	Concentrations are likely too low to have a sustained impact.
Pollution: urban stormwater	Toxicants	Adversely impacts waterbird breeding	Unlikely	Minor	Low	Concentrations are likely too low to have a sustained impact.
Pollution: urban stormwater	Toxicants	Adversely impacts threatened species (Australasian bittern and curlew sandpiper)	Unlikely	Minor	Low	Concentrations are likely too low to have a sustained impact.
Pollution: urban stormwater	Toxicants	Adversely affects other fauna	Possible	Minor	Low	Amphibians are known to be highly sensitive to toxicants in their environment (Mann and Bidwell 1999; Mann et al. 2003). Higher likelihood score reflects this.
Pollution: urban stormwater	Toxicants	Adversely affects wetland function (drainage and flood mitigation services)	Rare	Negligible	Negligible	Not a plausible impact pathway.
Pollution: urban	Toxicants	Adversely affects recreation and	Rare	Minor	Negligible	Concentrations of toxicants are well below those for primary contact recreation (ANZECC and ARMCANZ 2000). Main impact

Edithvale-Seaford Wetlands Ramsar Site Management Plan

Pressures	Stressors	Impact	Likelihood of impact	Consequence of impact	Risk	Evidence / comments
stormwater		amenity values				would be from decreased amenity values through impacts to other biota.
Pollution: urban stormwater	Toxicants	Adversely affects scientific and monitoring values	Rare	Minor	Negligible	Not a plausible impact pathway.
Pollution: urban stormwater	Toxicants	Adversely affects cultural values	Rare	Minor	Negligible	As for recreational and amenity values.
Water management: urban stormwater	Altered water regimes					An assessment of water regime requirements and current water regime at Edithvale-Seaford Wetlands indicated that Edithvale maintains an adequate water regime (except during extreme drought when it dries too quickly). The stability of the shallow wetlands at Seaford was, however, not ideal (SKM 2011). A capital works program was developed with construction of capital works in 2013. The site is now managed to slowly drawdown water levels during late spring and summer to both control invasive plants and provide habitat for shorebirds. The ability to manage water levels at the Ramsar sites is, however, limited. The site acts as flood mitigation and control for stormwater and is largely reliant on water inflows following heavy rainfall. Risks are assessed under the new management regime which includes the new infrastructure.
Water management: urban stormwater	Altered water regimes	Adversely impacts vegetation and habitat	Likely	Moderate	Medium	Control of invasive plants and domination of the site by a few aggressive species has been an ongoing issue. It is likely that this will continue into the future given the limitations on water regime management at the site.
Water management: urban stormwater	Altered water regimes	Adversely impacts waterbird diversity and abundance	Possible	Moderate	Medium	Hydrology and vegetation type have been identified as the most important habitat components for supporting waterbirds at the Ramsar site (Tzaros and Silcocks 2004). Any changes to habitat are likely to affect waterbird abundance and diversity.
Water management: urban stormwater	Altered water regimes	Adversely impacts waterbird breeding	Possible	Moderate	Medium	Breeding of many species is tightly linked to duration of inundation. Any sudden drawdowns could affect waterbird breeding success.
Water management: urban stormwater	Altered water regimes	Adversely impacts threatened species (Australasian bittern and curlew sandpiper)	Possible	Moderate	Medium	These two wading species use shallow water feeding habitat that may be affected by altered water regimes and prolonged inundation or drying.
Water management: urban stormwater	Altered water regimes	Adversely affects other fauna	Possible	Moderate	Medium	Particularly important for obligate aquatic fauna, if drying occurs over long periods.
Water management: urban stormwater	Altered water regimes	Adversely affects wetland function (drainage and flood mitigation services)	Rare	Negligible	Negligible	Not a plausible pathway
Water management: urban stormwater	Altered water regimes	Adversely affects recreation and amenity values	Unlikely	Minor	Low	Wetland and other amenity values exist in both the presence and absence of water. It is possible that a fluctuating water regime provides educational opportunities about the importance and values of temporary wetland systems.
Water management: urban stormwater	Altered water regimes	Adversely affects scientific and monitoring values	Rare	Negligible	Negligible	Not a plausible pathway
Water management: urban stormwater	Altered water regimes	Adversely affects cultural values	Unlikely	Minor	Low	As above for amenity values.
Water management: urban stormwater	Increased salinity					Prolonged drying results in saline water entering the system from groundwater (GHD 2006). There is also some concentration of salts in the system as wetlands dry. Water quality data from the site indicate periodic rises in salinity (linked to water regime) but no sustained trend over time (Melbourne Water unpublished data).
Water management: urban stormwater	Increased salinity	Adversely impacts vegetation and habitat	Possible	Moderate	Medium	Has been known to lead to an increase in the invasive spiny rush at the site. Altered salinity could result in a decline in freshwater taxa at the expense of more salt tolerant species.
Water management: urban stormwater	Increased salinity	Adversely impacts waterbird diversity and abundance	Unlikely	Minor	Low	Most of the species that occur at the site can tolerate a wide range of salinity conditions. The main impact pathway would be from loss of habitat.
Water management: urban stormwater	Increased salinity	Adversely impacts waterbird breeding	Unlikely	Minor	Low	Breeding of many species is tightly linked to duration of inundation. Any sudden drawdowns could affect waterbird breeding success.
Water management: urban stormwater	Increased salinity	Adversely impacts threatened species (Australasian bittern and curlew sandpiper)	Likely	Major	High	Curlew sandpiper feeds in both saline and freshwater environments. The Australasian bittern, however, is a freshwater species and declines elsewhere have been recorded when systems became secondary salinised (Jaensch, 2004).
Water management: urban stormwater	Increased salinity	Adversely affects other fauna	Likely	Moderate	Medium	Particularly important for obligate aquatic fauna, such as frogs and fish that are not adapted to saline conditions, particularly if salinity exceeds species tolerances.
Water management: urban stormwater	Increased salinity	Adversely affects wetland function (drainage and flood mitigation services)	Rare	Negligible	Negligible	Not a plausible pathway

Edithvale-Seaford Wetlands Ramsar Site Management Plan

Pressures	Stressors	Impact	Likelihood of impact	Consequence of impact	Risk	Evidence / comments
Water management: urban stormwater	Increased salinity	Adversely affects recreation and amenity values	Possible	Minor	Low	Altered amenity values from altered vegetation, increasing saline conditions.
Water management: urban stormwater	Increased salinity	Adversely affects scientific and monitoring values	Rare	Negligible	Negligible	Not a plausible pathway
Water management: urban stormwater	Increased salinity	Adversely affects cultural values	Possible	Moderate	Medium	Altered cultural values from altered vegetation, increasing saline conditions.
Urban development and recreation	Litter					Litter is a common occurrence in urban stormwater systems. Melbourne Water has installed litter traps to manage the problem and limit litter entering the wetlands.
Urban development and recreation	Litter	Adversely impacts vegetation and habitat	Rare	Negligible	Negligible	Very low likelihood of impact on vegetation and habitat given the extent of litter at the site.
Urban development and recreation	Litter	Adversely impacts waterbird diversity and abundance	Unlikely	Moderate	Low	Entanglement is a problem for some birds and waterbirds are also susceptible to ingestion of litter and micro-plastics with effects on nutrition and toxicity reported (Sutherland et al. 2012). However, the level of litter is low, making for low likelihoods.
Urban development and recreation	Litter	Adversely impacts waterbird breeding	Unlikely	Moderate	Low	As above
Urban development and recreation	Litter	Adversely impacts threatened species (Australasian bittern and curlew sandpiper)	Unlikely	Moderate	Low	As above
Urban development and recreation	Litter	Adversely affects other fauna	Unlikely	Moderate	Low	As above
Urban development and recreation	Litter	Adversely affects wetland function (drainage and flood mitigation services)	Rare	Negligible	Negligible	Not a plausible pathway?
Urban development and recreation	Litter	Adversely affects recreation and amenity values	Possible	Minor	Low	Litter does not extend across the site, but localised build-ups at drains and traps are unsightly.
Urban development and recreation	Litter	Adversely affects scientific and monitoring values	Rare	Negligible	Negligible	Not a plausible impact pathway.
Urban development and recreation	Litter	Adversely affects cultural values	Possible	Moderate	Medium	As per amenity value impacts.
Disturbance of Acid Sulfate Soils (ASS)	Acidity and liberation of heavy metals					Areas of ASS and Potential ASS are present in the Edithvale-Seaford Wetlands (O'Neill 2013). If disturbed due to prolonged drying of wetland areas or physical disturbance of the soil surface, then sulphuric acid is formed and can liberate metals from the sediments. Risks are associated with both acidification and the liberation of metals from the sediments (noting that sediment metal concentrations are generally low and within ANZECC sediment guidelines). Low pH (3–5) has been periodically recorded at Seaford Wetlands, but not Edithvale (Melbourne Water unpublished data). Melbourne Water has developed and implemented an ASS management plan (O'Neill 2013) to minimise risks associated with works in the wetlands.
Disturbance ASS	Acidity and liberation of metals	Adversely impacts vegetation and habitat	Possible	Moderate	Medium	Acidity is a risk to a number of wetland plant species (Sammut et al. 1996). The management of ASS at the site mitigates the likelihood and extent of impacts.
Disturbance ASS	Acidity and liberation of metals	Adversely impacts waterbird diversity and abundance	Possible	Minor	Low	Unlikely to be directly impacted from acidity and low level metal release, but impacts may be felt through the food chain from effects on prey.
Disturbance ASS	Acidity and liberation of metals	Adversely impacts waterbird breeding	Possible	Minor	Low	As above
Disturbance ASS	Acidity and liberation of metals	Adversely impacts threatened species (Australasian bittern and curlew sandpiper)	Possible	Minor	Low	As above
Disturbance ASS	Acidity and liberation of metals	Adversely affects other fauna	Possible	Moderate	Medium	There is a high potential for disturbed acid sulfate soils to destroy eggs and resting/estivating stages of frogs, reptiles and macroinvertebrates (Fitzpatrick et al. 2008).
Disturbance ASS	Acidity and liberation of metals	Adversely affects wetland function (drainage and flood mitigation services)	Possible	Minor	Low	Not a plausible pathway?
Disturbance ASS	Acidity and liberation of metals	Adversely affects recreation and amenity values	Possible	Moderate	Medium	Disturbance of ASS can result in unsightly conditions and odours. Likely to persist only for short durations in localised areas.

Edithvale-Seaford Wetlands Ramsar Site Management Plan

Pressures	Stressors	Impact	Likelihood of impact	Consequence of impact	Risk	Evidence / comments
Disturbance ASS	Acidity and liberation of metals	Adversely affects scientific and monitoring values	Rare	Negligible	Negligible	Not a plausible pathway?
Disturbance ASS	Acidity and liberation of metals	Adversely affects cultural values	Possible	Moderate	Medium	As above for amenity values.
Invasive species	Invasive native species: <i>Phragmites</i> and <i>Typha</i>					Vegetation monitoring indicates that common reed has contracted from 2013 to 2015 in low lying areas at Seaford Wetlands, but expanded in other areas, reflecting water regime changes during and post Millennium drought (Greet 2015). Melbourne Water has committed significant resources to the management of common reed within the Ramsar site, having undertaken trials of different control methods and implemented a dedicated management program.
Invasive species	Invasive native species: <i>Phragmites</i> and <i>Typha</i>	Adversely impacts vegetation and habitat	Likely	Major	High	This is actively managed by Melbourne Water and the risk has been assessed considering this intensive management regime.
Invasive species	Invasive native species: <i>Phragmites</i> and <i>Typha</i>	Adversely impacts waterbird diversity and abundance	Likely	Major	High	Impacts through altered habitat, particularly for birds that require open water or mudflats for foraging.
Invasive species	Invasive native species: <i>Phragmites</i> and <i>Typha</i>	Adversely impacts waterbird breeding	Possible	Minor	Low	Most waterbirds prefer the protective cover of reed beds for breeding, providing they're not overly dense.
Invasive species	Invasive native species: <i>Phragmites</i> and <i>Typha</i>	Adversely impacts threatened species (Australasian Bittern and Curlew Sandpiper)	Likely	Major	High	Australasian bittern requires cover of emergent vegetation, but density must be controlled to facilitate movement and hunting behaviours. Curlew sandpiper requires open mudflat habitat and control of emergent vegetation is essential.
Invasive species	Invasive native species: <i>Phragmites</i> and <i>Typha</i>	Adversely affects other fauna	Likely	Moderate	Medium	The mosaic of wetland habitats is important for diversity of fauna. Moves towards a monoculture of <i>Phragmites</i> would reduce the sites capacity to support a diversity of species.
Invasive species	Invasive native species: <i>Phragmites</i> and <i>Typha</i>	Adversely affects wetland function (drainage and flood mitigation services)	Likely	Major	High	Increases in plant biomass limit the wetlands storage capacity and increase the risk of localised flooding.
Invasive species	Invasive native species: <i>Phragmites</i> and <i>Typha</i>	Adversely affects recreation and amenity values	Possible	Moderate	Medium	The amenity values of the site would be diminished if the habitat mosaic including open water was no longer present and biodiversity values were reduced.
Invasive species	Invasive native species: <i>Phragmites</i> and <i>Typha</i>	Adversely affects scientific and monitoring values	Possible	Moderate	Medium	The scientific value of a less diverse site would be lower than a diverse one.
Invasive species	Invasive native species: <i>Phragmites</i> and <i>Typha</i>	Adversely affects cultural values	Possible	Moderate	Medium	As per amenity value impacts.
Invasive species	Spiny rush (<i>Juncus acutus</i>)					Vegetation monitoring from 2013 to 2015 indicates that spiny rush cover has remained relatively stable over that period (Greet 2015). Melbourne Water has an active invasive plant control program that targets spiny rush (among other invasive species).
Invasive species	Spiny rush (<i>Juncus acutus</i>)	Adversely impacts vegetation and habitat	Likely	Moderate	Medium	Although spiny rush is highly invasive it will be limited to salinised areas of the site.
Invasive species	Spiny rush (<i>Juncus acutus</i>)	Adversely impacts waterbird diversity and abundance	Likely	Minor	Medium	Related to loss of habitat values, however, providing the rush does not become too dense, and is limited to saline areas, it likely to be minor.
Invasive species	Spiny rush (<i>Juncus acutus</i>)	Adversely impacts waterbird breeding	Likely	Minor	Medium	As above
Invasive species	Spiny rush (<i>Juncus acutus</i>)	Adversely impacts threatened species (Australasian Bittern and Curlew Sandpiper)	Likely	Moderate	Medium	Australasian Bittern is unlikely to use the saline areas of the site. An expansion of spiny rush into mudflat habitat could impact the Curlew Sandpiper foraging habitat.
Invasive species	Spiny rush (<i>Juncus acutus</i>)	Adversely affects other fauna	Likely	Minor	Medium	Lower number of other fauna species using the saline areas of the site.
Invasive species	Spiny rush (<i>Juncus acutus</i>)	Adversely affects wetland function (drainage and flood mitigation services)	Unlikely	Minor	Low	Minor reductions in storage capacity possible.
Invasive species	Spiny rush (<i>Juncus acutus</i>)	Adversely affects recreation and amenity values	Possible	Minor	Low	As long as contained to small areas of salinised habitat, impact is small.
Invasive species	Spiny rush (<i>Juncus acutus</i>)	Adversely affects scientific and monitoring values	Possible	Negligible	Negligible	As long as contained to small areas of salinised habitat, impact is small.
Invasive species	Spiny rush (<i>Juncus acutus</i>)	Adversely affects cultural values	Possible	Minor	Low	As long as contained to small areas of salinised habitat, impact is small.

Edithvale-Seaford Wetlands Ramsar Site Management Plan

Pressures	Stressors	Impact	Likelihood of impact	Consequence of impact	Risk	Evidence / comments
	<i>acutus</i>)					
Invasive species	Predators (foxes, cats and rats)					While there are no trend data available, foxes and cats are regularly observed during bird surveys at the Ramsar site (Silcocks et al. 2006, Silcocks 2013). Melbourne Water undertakes periodic pest control in Edithvale South. Kingston Council have been undertaking pest control near Edithvale North and Melbourne Water together with City of Frankston undertake pest control at Seaford.
Invasive species	Predators (foxes, cats and rats)	Adversely impacts waterbird diversity and abundance	Almost certain	Moderate	High	Predation by foxes, cats and rats is a significant threat to foraging waterbirds, particularly waders that are not out on deep, open water.
Invasive species	Predators (foxes, cats and rats)	Adversely impacts waterbird breeding	Almost certain	Moderate	High	Nesting birds and eggs are vulnerable to predation.
Invasive species	Predators (foxes, cats and rats)	Adversely impacts threatened species (Australasian Bittern and Curlew Sandpiper)	Almost certain	Major	Extreme	The consequences of mortality of a few birds from small threatened populations are higher than for more common species.
Invasive species	Predators (foxes, cats and rats)	Adversely affects other fauna	Likely	Moderate	Medium	Other fauna at the site is also vulnerable to predation, but perhaps at lower risk than waterbirds.
Invasive species	Native species: kangaroos					Recent surveys indicated a density of 1.8 kangaroos per hectare at Edithvale Wetlands. This is two to three times the sustainable population (Hynes 2015a). Melbourne Water has implemented control strategies.
Invasive species	Native species: kangaroos	Adversely impacts vegetation and habitat	Possible	Minor	Low	Primary impact is through trampling not grazing and impact is low (Hynes 2015b).
Invasive species	Native species: kangaroos	Adversely affects recreation and amenity values	Possible	Minor	Low	Risk of injury from fleeing / disturbed kangaroos is minor as visitors to the wetland centre are low.
Invasive species	Native species: kangaroos	Adversely affects other fauna	Likely	Moderate	Medium	Impacts to bush birds, nesting birds and ground mammals are expected from grazing of terrestrial grasslands in the site (Hynes 2015b).
Invasive species	Grazing animals (pigs, rabbits)					Rabbits are regularly observed within the site, and recently there has been evidence of a pig in the Seaford Wetlands. Melbourne Water undertakes pest control within the site.
Invasive species	Grazing animals (pigs, rabbits)	Adversely impacts vegetation and habitat	Almost certain	Minor	Medium	While it is almost certain that rabbits (and the single pig) are grazing on vegetation, this is likely to be limited to dryland areas with respect to rabbits.
Invasive species	Grazing animals (pigs, rabbits)	Adversely impacts waterbird diversity and abundance	Possible	Minor	Low	Based on impacts to vegetation and habitat.
Invasive species	Grazing animals (pigs, rabbits)	Adversely impacts waterbird breeding	Possible	Minor	Low	Based on impacts to vegetation and habitat.
Invasive species	Grazing animals (pigs, rabbits)	Adversely impacts threatened species (Australasian Bittern and Curlew Sandpiper)	Possible	Minor	Low	Based on impacts to vegetation and habitat.
Invasive species	Grazing animals (pigs, rabbits)	Adversely affects other fauna	Possible	Minor	Low	Based on impacts to vegetation and habitat.
Invasive species	Grazing animals (pigs, rabbits)	Adversely affects wetland function (drainage and flood mitigation services)	Rare	Negligible	Negligible	Not a plausible pathway?
Invasive species	Grazing animals (pigs, rabbits)	Adversely affects recreation and amenity values	Unlikely	Minor	Low	Minor impacts to amenity value over small areas.
Invasive species	Grazing animals (pigs, rabbits)	Adversely affects scientific and monitoring values	Rare	Negligible	Negligible	Not a plausible pathway?
Invasive species	Grazing animals (pigs, rabbits)	Adversely affects cultural values	Unlikely	Minor	Low	Minor impacts to amenity value over small areas.
Recreational activities	Human disturbance (noise and visual disturbance)					Edithvale-Seaford Wetlands are within an urban setting and there is ambient noise and visual disturbance from traffic and other surrounding urban activities, as well as from recreational users of the site, including those that use the site for casual walking, birdwatching and dog walking on- and off-leash. There is also increasing pressure to use the site for events (e.g. running events). The centre of the wetlands, however, has restricted access.

Pressures	Stressors	Impact	Likelihood of impact	Consequence of impact	Risk	Evidence / comments
Recreational activities	Human disturbance (noise and visual disturbance)	Adversely impacts waterbird diversity and abundance	Possible	Moderate	Medium	Human presence impacts on shorebirds is well documented (e.g. Weston 2003; Weston and Elgar 2005 and 2007; Antos et al. 2007; Weston et al. 2012; Martin et al. 2015). For migratory shorebirds impacts may include displacement from foraging areas, reduced feeding and unnecessary energy use feared to impact birds abilities to successfully make the return journey to the northern hemisphere to breed. For breeding waterbirds, the disturbance may result in periods of reduced parental care and absences from the nest, exposing eggs or nestlings to increased risk of predation or weather effects potentially resulting in a lower chance of survivorship of young and reduced breeding success. Restricted access at the site reduces both the likelihood and magnitude of disturbance.
Recreational activities	Human disturbance (noise and visual disturbance)	Adversely impacts waterbird breeding	Possible	Moderate	Medium	As above.
Recreational activities	Human disturbance (noise and visual disturbance)	Adversely impacts threatened species (Australasian bittern and curlew sandpiper)	Possible	Moderate	Medium	As above.
Recreational activities	Human disturbance (noise and visual disturbance)	Adversely affects other fauna	Possible	Moderate	Medium	Other resident fauna are likely to be more habituated to urban environment and therefore less likely to be impacted by noise and visual disturbance in an urban setting than waterbirds.
Climate change						Regional climate projections have recently been released by CSIRO for sub-cluster regions in Australia. The relevant region for Edithvale-Seaford Wetlands is "Southern Slopes Victoria West" http://www.climatechangeinaustralia.gov.au/en/climate-projections/future-climate/regional-climate-change-explorer/sub-clusters/?current=SSVWC&tooltip=true&popup=true. Of concern for the Ramsar site is the potential increased fire risk under hotter and drier conditions, the implications of sea level rise on Seaford Wetlands and altered water regimes in a future climate of more extreme rainfall events.
Climate change	Increased frequency / intensity of fire					The intensity and frequency of fires in south eastern Australia is predicted to rise under future climate predictions (Williams et al. 2013). The risk of unplanned fires will increase and it is likely that if commenced the fire would spread rapidly and burn the peat layer (Terramatrix 2013). Melbourne Water has a fire management plan in place. Terramatrix (2013) undertook a risk assessment for this pressure and stressor and those have been adapted to the consequence and likelihood tables used here.
Climate change	Increased frequency / intensity of fire	Adversely impacts vegetation and habitat	Possible	Major	High	As per Terramatrix (2013).
Climate change	Increased frequency / intensity of fire	Adversely impacts waterbird diversity and abundance	Possible	Major	High	As per Terramatrix (2013).
Climate change	Increased frequency / intensity of fire	Adversely impacts waterbird breeding	Possible	Major	High	As per Terramatrix (2013).
Climate change	Increased frequency / intensity of fire	Adversely impacts threatened species (Australasian Bittern and Curlew Sandpiper)	Possible	Major	High	As per Terramatrix (2013).
Climate change	Increased frequency / intensity of fire	Adversely affects other fauna	Possible	Major	High	As per Terramatrix (2013).
Climate change	Increased frequency / intensity of fire	Adversely affects wetland function (drainage and flood mitigation services)	Possible	Minor	Low	As per Terramatrix (2013).
Climate change	Increased frequency / intensity of fire	Adversely affects recreation and amenity values	Unlikely	Major	Medium	As per Terramatrix (2013).
Climate change	Increased frequency / intensity of fire	Adversely affects scientific and monitoring values	Rare	Minor	Negligible	As per Terramatrix (2013).
Climate change	Increased frequency / intensity of fire	Adversely affects cultural values	Unlikely	Major	Medium	As per Terramatrix (2013).
Climate change	Sea level rise					Sea levels are predicted to increase by 0.08 to 0.18 m by 2030 with a very high degree of confidence (Grose et al. 2015). Local assessments indicate that sea water could rise up into the wetlands through connecting channels and rising groundwater over the period 2030 – 2070, substantially altering the character of the wetlands (DELWP 2015).
Climate change	Sea level rise	Adversely impacts vegetation and habitat	Almost certain	Major	Extreme	Saline intrusion likely to transition habitat from freshwater to brackish and reed dominated to halophyte (saltmarsh) dominance, including potential expansion of Sea Rush, Shrubby Glasswort, and various halophytic herbs. Melbourne Water suggest that there

Edithvale-Seaford Wetlands Ramsar Site Management Plan


Pressures	Stressors	Impact	Likelihood of impact	Consequence of impact	Risk	Evidence / comments
						is greater potential to mitigate the impacts of sea level rise at Edithvale (feasible) than Seaford (much less feasible).
Climate change	Sea level rise	Adversely impacts waterbird diversity and abundance	Almost certain	Major	Extreme	Loss or much reduced breeding habitat for freshwater dependent species e.g. Australasian Bittern, Blue-billed Duck, Hardyheads and Baillon's Crake.
Climate change	Sea level rise	Adversely impacts waterbird breeding	Almost certain	Major	Extreme	Loss or much reduced breeding habitat for freshwater dependent waterbirds – as above.
Climate change	Sea level rise	Adversely impacts threatened species (Australasian bittern and curlew sandpiper)	Almost certain	Major	Extreme	Loss or much reduced habitat for Australasian Bittern, potentially neutral to low impact for Curlew Sandpiper.
Climate change	Sea level rise	Adversely affects other fauna	Almost certain	Major	Extreme	Loss or reduced habitat for seed-dependant passerines e.g. Little Grassbird, Australian Reed-warbler.
Climate change	Sea level rise	Adversely affects wetland function (drainage and flood mitigation services)	TBD	TBD	#N/A	Discuss implications with Melbourne Water.
Climate change	Sea level rise	Adversely affects recreation and amenity values	Possible	Minor	Low	Change in wetland character may have a minor impact on amenity values.
Climate change	Sea level rise	Adversely affects scientific and monitoring values	Likely	Moderate	Medium	Change in ecological character will diminish its value as a remnant of Carrum Swamp.
Climate change	Sea level rise	Adversely affects cultural values	Possible	Moderate	Medium	On-going change from its original state will likely reduce its cultural values.
Climate change	Increased frequency & intensity of storms					Extreme events (storms and high rainfall events) are predicted to occur with high confidence (Grose et al. 2015). As the dominant water source at Edithvale-Seaford Wetlands is urban stormwater, it is likely that this will arrive less frequently, but with higher magnitudes.
Climate change	Increased frequency & intensity of storms	Adversely impacts vegetation and habitat	Possible	Moderate	Medium	Storm surges (tides) with added wind force plus sea level rise is predicted to increase the area of tidal inundation (DELWP 2015) potentially damaging habitat and enhancing a trend of ecological change.
Climate change	Increased frequency & intensity of storms	Adversely impacts waterbird diversity and abundance	Possible	Moderate	Medium	Increased storms and flooding could reduce the wetlands role as a storm refuge/roosting habitat, further impacting on waterbirds.
Climate change	Increased frequency & intensity of storms	Adversely impacts waterbird breeding	Possible	Moderate	Medium	Increased spring-summer storms could impact on waterbird breeding.
Climate change	Increased frequency & intensity of storms	Adversely impacts threatened species (Australasian bittern and curlew sandpiper)	Possible	Moderate	Medium	Increased storms and tidal inundation would further impact on an already compromised habitat for Australasian Bittern. Storms would impact on the foraging capacity for migratory shorebirds.
Climate change	Increased frequency & intensity of storms	Adversely affects other fauna	Possible	Moderate	Medium	Storm damage to habitat would impact on most fauna.
Climate change	Increased frequency & intensity of storms	Adversely affects wetland function (drainage and flood mitigation services)	Possible	Major	High	Any information about whether this will go over capacity?
Climate change	Increased frequency & intensity of storms	Adversely affects recreation and amenity values	Possible	Moderate	Medium	Potential loss of amenity and recreation values due to storm damage.
Climate change	Increased frequency & intensity of storms	Adversely affects scientific and monitoring values	Possible	Moderate	Medium	Increased frequency of severe perturbations generally deleterious for scientific and monitoring values
Climate change	Increased frequency and intensity of storms (increased intensity of stormwater)	Adversely affects cultural values	Possible	Moderate	Medium	Potential change to land forms and geomorphological process.


Appendix 9 Works undertaken by Melbourne Water since the production of the last Ramsar Management Plan (KBR 2009).

Works	Melbourne Water/Company	Timeframe
Reports		
Hydrology / Enviro Water Requirements	SKM	from 2010 to 2011
Weeds and Overabundant Natives	Australian Ecosystems	2011
Bird Diversity and Abundance	Birds Australia	2010 and 2011
	BirdLife Australia	2013a, 2013b, 2015 and 2016
Kangaroo Management Plan	Ecoplan	2015
Kangaroo Control	VWSA	2016
Fire Management Plan (Seaford Wetlands)	Terramatrix	2013
<i>In Progress</i>		
Hydrology (Update REALM, Risk Assessment, and Operational Plan)	Jacobs	2014-current
Water Quality Review and Direction	Jacobs	2016
Habitat Hectare and trajectories	Australian Ecosystems	2015
Monitoring		
Birds and Frogs (frog calls only)	BirdLife Australia	Monthly from 2008-current
Frog Census	Volunteers	2012–2013
Bat Monitoring	Parsons Brinckerhoff	2009 and 2010
Kangaroo Count	Ecoplan	2013-2015
	VWSA	2016
Mosquitos	Ecowise	2008–2010,
	ALS	2011
	GHD	2012-current
Water Quality -	Waterwatch	from 2009 to2010,
	GHD	from 2011– current
Groundwater Monitoring	GHD	2014-current
Water Level	Thiess	from 2008–2012
	BirdLife Australia	from 2013-current
Weed Mapping	Australian Ecosystems	in 2011 and 2015
Habitat Hectares	Australian Ecosystems	in 2010 and 2015
Tall Marsh/ <i>Phragmites</i> Mapping	Australian Ecosystems in and	2011
	Melbourne Water	2013
Vegetation Quadrats (Seaford)	University of Melbourne	from 2013 –current
Photo-points (Seaford)	Australian Ecosystems	in 2010
	University of Melbourne	from 2013– current
Photo-points (Edithvale)	Australian Ecosystems	in 2010
	Melbourne Water	in 2014
Capital Works		
Edithvale Education Centre	Melbourne Water	2011

Works	Melbourne Water/Company	Timeframe
Upgrade to Edithvale South Wetlands Bird Hide	Friends of Edithvale-Seaford Wetlands	2016
Seaford Hydrology Works –	Melbourne Water	from 2011 to 2014
Weed Removal and Revegetation of Seaford (east)	Melbourne Water	from 2008–2010
Maintenance		
Weed Control – Edithvale	Melbourne Water	Monthly from 2008 by
Weed Control – Seaford	Melbourne Water	Monthly from 2008
	FCC	On-going
Pest Control	Melbourne Water contractor	2013 and 2015
Nest Box Clearing	Melbourne Water	2012/2013, 2013/2014 and 2014/2015
Kangaroo Management (Sterilisation)	VWSA	2009 and 2016
Mosquito Management (Spray or Briquettes) –	Melbourne Water	monthly and on-going from 2008
Grooming of <i>Phragmites</i> –	CSA	2011 to current
Desilting -	Melbourne Water	on-going and annual
Community		
ESW Community Liaison Committee (CLC)	CLC	Quarterly meetings on-going

Appendix 10 Mapping of weeds of management concern at Edithvale and Seaford Wetlands, undertaken by Australian Ecosystems (2016).





Seaford Wetlands

Legend

Survey Effort

- Study Area Extent

High Priority Weeds

- Acacia longifolia (Sallow Wattle)
- Chrysanthemoides monilifera (Boneseed)
- Coprosma repens (Mirror Bush)
- Juncus acutus (Spiny Rush)
- Lycium ferocissimum (African Box-thorn)
- Mesembryanthemum crystallinum (Common Ice Plant)
- Phytolacca octandra (Red-ink Weed)
- Pinus radiata (Radiata Pine)
- Pittosporum undulatum (Sweet Pittosporum)
- Rosa rubiginosa (Sweet Briar)
- Rubus anglocandicans (Common Blackberry)
- Salpichroa origanifolia (Pampas Lily-of-the-valley)
- Ulex europaeus (Gorse)
- Coprosma repens (Mirror Bush)
- Casuarina cunninghamiana (River Oak)
- Scattered Acacia longifolia (Sallow Wattle), Rubus anglocandicans (Common Blackberry) and Ulex europaeus (Gorse)
- Leptospermum laevigatum (Coast Tea-tree)

Hydrology

- Major River, Creek, Channel or Drain
- Minor Creek, Channel or Drain

Transportation

- Freeway
- Highway
- Sealed Arterial Road
- Sealed Road



Coordinate System: GDA 1994 MGA Zone 55
 Projection: Transverse Mercator
 Datum: GDA 1994



Compilation Notes: Vicmap Products (Copyright The State of Victoria, Department of Environment, Land, Water and Planning 2015) have been used in preparing this map. 2013 Aerial Imagery courtesy of Melbourne Water.

Project: Melbourne Water Sites of Biodiversity Significance: 2014 -2015 Surveys
 Final Map prepared by Holocene Environmental Science 28th December 2015
 Using GIS layers prepared by Amanda Featham (GeoEccentric) September 2015
 Surveyors: Bryan Mole (Australian Ecosystems Pty Ltd) and Damien Cook
 Survey Period: 3rd - 4th February 2015

Disclaimer: while every care has been taken to ensure the accuracy of this product, no representations or warranties about its accuracy, completeness or suitability for any particular purpose is made. Liability of any kind for any expenses, losses, damages and/or costs which are or may be incurred as a result of this product being inaccurate, incomplete or unsuitable in any way and for any reason will not be accepted.



Appendix 11 Additional weed control and revegetation information.

Herbicide use

All herbicide usage must be in accordance with the following:

- All use of herbicides (and associated additives) will be in accordance with the product label. Off-label use of herbicides may be permitted where approval has been granted from a state government department (e.g. Department of Environment, Land, Water and Planning).
- Site-specific planning for herbicide use (application methods, chemicals used, weather conditions, plant phenology, etc.) will be employed to avoid off-target herbicide damage to indigenous vegetation. Off-target herbicide damage is the detrimental application of herbicide to plant species that have not been targeted for control. While this generally applies to plants in and around the point of herbicide application, it may also refer to organisms (flora and fauna) some distance away.

Revegetation / supplementary planting

The process of revegetation requires planning, documentation, implementation, monitoring and maintenance, all of which are essential in ensuring success. Each of these stages is discussed below in relation to the reserve. More information can also be found in *Melbourne Water Provenance Policy*, *Melbourne Water Standard for Plant Installation* and *Melbourne Water Standard for Plant Supply* (Melbourne Water 2013c, 2014a, b, respectively).

Appendix 12 Plant species suitable for revegetation in the various habitat types of Edithvale-Seaford Wetlands Ramsar Site (brackish wetlands, freshwater wetlands, Swamp Scrub and Damp Sands Herb-rich Woodland).

Key:

Structural role

- A Structural dominant of the vegetation stratum
- B Localised structural co-dominant in vegetation stratum
- C Scattered thinly or discontinuously as small groups or isolated individuals with low overall cover

Planting zone (wetlands only)

- 1 Seasonally wet margins to permanently moist; shallow seasonal inundation in lower part of zone
- 2 Shallow inundation; minimum depth of inundation c. 10 cm; amphibious and emergent aquatic herbs, some
- 3 Permanent water; submergent and emergent aquatic-herbs, some straddling Zones 2 and 3

Freshwater Wetlands

Scientific Name	Common Name	Structural role	Planting zone
Grasses and graminoids			
<i>Alisma plantago-aquatica</i>	Water Plantain	C	2
<i>Amphibromus fluitans</i>	River Swamp Wallaby-grass	C	1, 2
<i>Baumea arthropphylla</i>	Fine Twig-sedge	A	2
<i>Baumea articulata</i>	Jointed Twig-sedge	C	2
<i>Bolboschoenus medianus</i>	Marsh Club-sedge	C	1, 2
<i>Cladium procerum</i>	Leafy Twig-rush	C	2
<i>Cycnogeton procerum</i> (broad erect leaves)	Water Ribbons	B	2
<i>Eleocharis acuta</i>	Common Spike-sedge	B	1, 2
<i>Eleocharis sphacelata</i>	Tall Spike-sedge	A	3
<i>Juncus amabilis</i>	Hollow Rush	B	1
<i>Juncus australis</i>	Austral Rush	B	1
<i>Juncus holoschoenus</i>	Joint-leaf Rush	C	1
<i>Juncus pallidus</i>	Pale Rush	C	1
<i>Juncus pauciflorus</i>	Loose-flower Rush	C	1
<i>Juncus procerus</i>	Tall Rush	C	1, 2
<i>Potamogeton cheesemanii</i>	Small-fruit Pondweed	C	2, 3
<i>Potamogeton crispus</i>	Curly Pondweed	C	3
<i>Potamogeton ochreatus</i>	Blunt Pondweed	C	3
<i>Schoenoplectus tabernaemontani</i>	River Club-sedge	B	3
<i>Schoenus brevifolius</i>	Zig-zag Bog-sedge	C	1
<i>Stuckenia pectinata</i>	Fennel Pondweed	C	3
<i>Triglochin striata</i>	Streaked Arrowgrass	C	1, 2
<i>Vallisneria australis</i>	Eel Grass	C	3
Dicot herbs			
<i>Centella cordifolia</i>	Centella	B	1
<i>Crassula helmsii</i>	Swamp Crassula	C	1, 2
<i>Epilobium billardierianum</i> ssp. <i>billardierianum</i>	Smooth Willow-herb	C	1
<i>Epilobium billardierianum</i> ssp. <i>cinereum</i>	Grey Willow-herb	C	1
<i>Epilobium hirtigerum</i>	Hairy Willow-herb	C	1
<i>Lilaeopsis polyantha</i>	Australian Lilaeopsis	B	2

Scientific Name	Common Name	Structural role	Planting zone
<i>Liparophyllum exaltatum</i>	Erect Marsh-flower	C	2, 3
<i>Lobelia anceps</i>	Angled Lobelia	C	1, 2
<i>Lycopus australis</i>	Australian Gipsywort	C	1, 2
<i>Lythrum hyssopifolia</i>	Small Loosestrife	C	1, 2
<i>Myriophyllum caput-medusae</i>	Coarse Water-milfoil	C	3
<i>Myriophyllum crispatum</i>	Upright Water-milfoil	C	2, 3
<i>Myriophyllum simulans</i>	Amphibious Water-milfoil	b	2
<i>Ornduffia reniformis</i>	Running Marsh-flower	C	2, 3
<i>Ottelia ovalifolia</i>	Swamp Lily	C	2, 3
<i>Persicaria decipiens</i>	Slender Knotweed	C	
<i>Philydrum lanuginosum</i>	Woolly Waterlily	C	1
<i>Selliera radicans</i>	Shiny Swamp-mat	B	1

Brackish Wetlands

Scientific Name	Common Name	Structural role	Planting zone
Grasses and graminoids			
<i>Baumea arthropophylla</i>	Fine Twig-sedge	C	2
<i>Baumea articulata</i>	Jointed Twig-sedge	C	2
<i>Baumea juncea</i>	Bare Twig-sedge	C	1
<i>Bolboschoenus caldwellii</i>	Salt Club-sedge	B	1, 2
<i>Cladium procerum</i>	Leafy Twig-rush	C	2
<i>Cycnogeton procerum</i> (broad erect leaves)	Water Ribbons	C	2
<i>Distichlis distichophylla</i>	Australian Salt-grass	B	1
<i>Eleocharis acuta</i>	Common Spike-sedge	A	1, 2
<i>Juncus kraussii</i> ssp. <i>australiensis</i>	Sea Rush	A	1, 2
<i>Juncus pallidus</i>	Pale Rush	C	1, 2
<i>Poa poiformis</i> var. <i>poiformis</i>	Coast Tussock-grass	C	1
<i>Potamogeton crispus</i>	Curly Pondweed	C	3
<i>Potamogeton ochreateus</i>	Blunt Pondweed	C	3
<i>Stuckenia pectinata</i>	Fennel Pondweed	B	3
<i>Schoenoplectus tabernaemontani</i>	River Club-sedge	C	3
<i>Triglochin striata</i>	Streaked Arrowgrass	A	1, 2
Dicot herbs			
<i>Apium prostratum</i> ssp. <i>prostratum</i>	Sea Celery	C	1
<i>Chenopodium glaucum</i>	Glaucous Goosefoot	C	1
<i>Crassula helmsii</i>	Swamp Crassula	C	1, 2
<i>Disphyma crassifolium</i> ssp. <i>clavellatum</i>	Rounded Noon-flower	C	1
<i>Lilaeopsis polyantha</i>	Australian Lilaeopsis	B	1
<i>Lobelia anceps</i>	Angled Lobelia	B	1
<i>Thyridia repens</i>	Creeping Monkey-flower	B	1, 2
<i>Myriophyllum salsugineum</i>	Lake Water-milfoil	B	3

Scientific Name	Common Name	Structural role	Planting zone
<i>Myriophyllum verrucosum</i>	Red Water-milfoil	C	2, 3
<i>Samolus repens</i>	Creeping Brookweed	C	1, 2
<i>Selliera radicans</i>	Shiny Swamp-mat	A	1, 2

Swamp Scrub

Scientific name	Common name	Structural role	Notes
Trees			
<i>Acacia melanoxylon</i>	Blackwood	B	
<i>Eucalyptus camaldulensis</i>	River Red-gum	C	
<i>Eucalyptus ovata</i> var. <i>ovata</i>	Swamp Gum	C	
Shrubs			
<i>Acacia stricta</i>	Hop Wattle	C	
<i>Acacia verticillata</i>	Prickly Moses	B	
<i>Bursaria spinosa</i> ssp. <i>spinosa</i>	Sweet Bursaria	C	
<i>Cassinia aculeata</i>	Common Cassinia	C	
<i>Goodenia ovata</i>	Hop Goodenia	B	
<i>Hakea nodosa</i>	Yellow Hakea	C	
<i>Leptospermum lanigerum</i>	Woolly Tea-tree	B	
<i>Leptospermum scoparium</i>	Manuka	B	
<i>Melaleuca ericifolia</i>	Swamp Paperbark	A	
<i>Melaleuca squarrosa</i>	Scented Paperbark	C	
<i>Ozothamnus ferrugineus</i>	Tree Everlasting	B	
<i>Ozothamnus rosmarinifolius</i>	Rosemary Everlasting	C	
<i>Rhagodia candolleana</i> ssp. <i>candolleana</i>	Seaberry Saltbush	C	
<i>Rubus parvifolius</i>	Small-leaf Bramble	B	
<i>Viminaria juncea</i>	Golden Spray	C	
Grasses and graminoids			
<i>Apodasmia brownii</i>	Coarse Twine-rush	C	
<i>Baumea arthropphylla</i>	Fine Twig-sedge	C	Permanently moist soils
<i>Baumea articulata</i>	Jointed Twig-sedge	C	Permanently moist soils
<i>Baumea juncea</i>	Bare Twig-sedge	C	
<i>Carex appressa</i>	Tall Sedge	B	Seasonally moist soils
<i>Carex fascicularis</i>	Tassel Sedge	C	Permanently moist soils
<i>Carex gaudichaudiana</i>	Fen Sedge	C	Seasonally moist soils
<i>Carex tereticaulis</i>	Hollow Sedge	C	Seasonally moist soils
<i>Chorizandra australis</i>	Heron Bristle-sedge	C	Permanently moist soils
<i>Cladium procerum</i>	Leafy Twig-rush	C	Permanently moist soils
<i>Cycnogeton procerum</i> (broad erect leaves)	Water Ribbons	B	Permanently moist soils
<i>Cyperus lucidus</i>	Leafy Flat Sedge	C	Permanently moist soils
<i>Distichlis distichophylla</i>	Australian Salt-grass	C	Brackish environments
<i>Empodisma minus</i>	Spreading Rope-rush	C	
<i>Gahnia radula</i>	Thatch Saw-sedge	C	
<i>Gahnia sieberiana</i>	Red-fruit Saw-sedge	B	
<i>Gahnia trifida</i>	Coast Saw-sedge	C	Brackish environments
<i>Gleichenia microphylla</i>	Scrambling Coral-fern	C	
<i>Hemarthria uncinata</i> var. <i>uncinata</i>	Mat Grass	C	

Scientific name	Common name	Structural role	Notes
<i>Juncus amabilis</i>	Hollow Rush	B	Seasonally moist soils
<i>Juncus australis</i>	Austral Rush	B	Seasonally moist soils
<i>Juncus flavidus</i>	Gold Rush	C	
<i>Juncus holoschoenus</i>	Joint-leaf Rush	C	Seasonally moist soils
<i>Juncus kraussii ssp. australiensis</i>	Sea Rush	C	Brackish environments
<i>Juncus pallidus</i>	Pale Rush	C	Seasonally moist soils
<i>Juncus pauciflorus</i>	Loose-flower Rush	C	Seasonally moist soils
<i>Juncus procerus</i>	Tall Rush	C	Permanently moist soils
<i>Lepidosperma longitudinale</i>	Pithy Sword-sedge	C	Moist soils
<i>Lepyrodia muelleri</i>	Common Scale-rush	C	Moist soils
<i>Lomandra longifolia ssp. longifolia</i>	Spiny-headed Mat-rush	C	
<i>Poa labillardierei var. labillardierei</i>	Common Tussock-grass	C	
<i>Schoenus brevifolius</i>	Zig-zag Bog-sedge	C	Moist soils
<i>Schoenus lepidosperma</i>	Slender Bog-sedge	C	
Other (dicot herbs, scramblers/climbers, ferns)			
<i>Acaena novae-zelandiae</i>	Bidgee-widgee	B	
<i>Clematis microphylla</i>	Small-leaved Clematis	C	
<i>Lycopus australis</i>	Australian Gipsywort	C	Permanently moist soils
<i>Pteridium esculentum</i>	Austral Bracken	C	
<i>Pteris tremula</i>	Tender Brake	C	
<i>Tetragonia implexicoma</i>	Bower Spinach	C	

Damp Sands Herb-rich Woodland

Scientific name	Common name	Structural role
Trees		
<i>Acacia mearnsii</i>	Black Wattle	C
<i>Acacia melanoxylon</i>	Blackwood	B
<i>Allocasuarina littoralis</i>	Black Sheoak	B
<i>Allocasuarina verticillata</i>	Drooping Sheoak	C
<i>Banksia integrifolia ssp. integrifolia</i>	Coast Banksia	B
<i>Eucalyptus camaldulensis</i>	River Red-gum	C
<i>Eucalyptus ovata var. ovata</i>	Swamp Gum	C
<i>Eucalyptus radiata ssp. radiata</i>	Narrow-leaf Peppermint	C
<i>Eucalyptus viminalis ssp. pryoriana</i>	Coast Manna-gum	A
Shrubs		
<i>Acacia oxycedrus</i>	Spike Wattle	C
<i>Acacia paradoxa</i>	Hedge Wattle	C
<i>Acacia stricta</i>	Hop Wattle	C
<i>Acacia suaveolens</i>	Sweet Wattle	C
<i>Acacia ulicifolia</i>	Juniper Wattle	C
<i>Allocasuarina paradoxa</i>	Green Sheoak	C
<i>Amperea xiphoclada var. xiphoclada</i>	Broom Spurge	C
<i>Aotus ericoides</i>	Common Aotus	C
<i>Banksia marginata</i>	Silver Banksia	B
<i>Bossiaea cinerea</i>	Showy Bossiaea	B
<i>Bursaria spinosa ssp. spinosa</i>	Sweet Bursaria	B
<i>Cassinia aculeata</i>	Common Cassinia	C
<i>Correa reflexa var. reflexa</i>	Common Correa	B

Scientific name	Common name	Structural role
<i>Dillwynia glaberrima</i>	Smooth Parrot-pea	C
<i>Dillwynia sericea s.l.</i>	Showy Parrot-pea	C
<i>Epacris impressa</i>	Common Heath	B
<i>Indigofera australis</i>	Austral Indigo	C
<i>Leptospermum myrsinoides</i>	Heath Tea-tree	B
<i>Leptospermum scoparium</i>	Manuka	B
<i>Leucopogon ericoides</i>	Pink Beard-heath	C
<i>Leucopogon virgatus var. virgatus</i>	Common Beard-heath	B
<i>Monotoca scoparia</i>	Prickly Broom-heath	C
<i>Olearia lirata</i>	Snowy Daisy-bush	C
<i>Olearia ramulosa</i>	Twiggy Daisy-bush	C
<i>Ozothamnus ferrugineus</i>	Tree Everlasting	C
<i>Persoonia juniperina</i>	Prickly Geebung	C
<i>Platylobium obtusangulum</i>	Common Flat-pea	C
<i>Ricinocarpos pinifolius</i>	Wedding Bush	C
Grasses and graminoids		
<i>Apodasmia brownii</i>	Coarse Twine-rush	C
<i>Gahnia radula</i>	Thatch Saw-sedge	C
<i>Hemarthria uncinata var. uncinata</i>	Mat Grass	C
<i>Lomandra longifolia ssp. longifolia</i>	Spiny-headed Mat-rush	C
<i>Poa labillardierei var. labillardierei</i>	Common Tussock-grass	C
<i>Poa sieberiana var. sieberiana</i>	Grey Tussock-grass	C
Other (dicot herbs, scramblers/climbers, ferns)		
<i>Acaena novae-zelandiae</i>	Bidgee-widgee	B
<i>Dianella brevicaulis</i>	Small-flower Flax-lily	C
<i>Clematis microphylla</i>	Small-leaved Clematis	C
<i>Pteridium esculentum</i>	Austral Bracken	C




Figure 18 Edithvale Wetlands revegetation zones.










Figure 19 Seaford Wetlands revegetation zones.

Appendix 13 Revegetation prescriptions for Edithvale-Seafood Wetlands from TBLA and Australian Ecosystems (2005).

The Edithvale-Seafood Wetlands Revegetation Prescriptions (TBLD and Australian Ecosystems 2005) provides a detailed revegetation strategy for Edithvale-Seafood Wetlands. This information is still useful and for the most part relevant. The figures illustrating the details of the revegetation prescriptions are provided below (TBLD and Australian Ecosystems 2005).



LA ZONE H
Eastern interface of Edithvale North with Branganan Drive from Edithvale Road north to Lewellyn Avenue

Existing conditions

- Width of the zone varies, but is generally between 30 to 40 metres.
- Revegetation areas have been established in scattered clumps between residential properties and the wetland.
- This area is noted as being dominated by introduced bird species including the Common Myna, Common Starling and Spotted Turtle-Dove. The residential land use is the primary influence on this along with the vegetation style and type (Birds Australia, 2004).

Revegetation Prescription Aims for LA Zone H

- Plant additional indigenous vegetation reflective of the Plains Grassland/Grassy Woodland EVC character in this zone.
- Improve the shrub and field layer species in the existing and new planting areas to improve habitat diversity.
- Retain selective view sheds over the wetlands from adjoining properties and council open space reserves.

LA ZONE I
North-eastern interface of Edithvale North extending from Lewellyn Avenue open space reserves, north west to Nurten Parade.

Existing conditions

- Reserve between wetland fence and residential properties widens to approximately 50 metres in width.
- Bird habitat values are similar to those described in LA Zone H.

Revegetation Prescription Aims for LA Zone I

- Plant additional indigenous vegetation reflective of the Plains Grassland/Grassy Woodland EVC character in this zone.
- Improve the shrub and field layer species in the existing and new planting areas to improve habitat diversity.
- Retain selective view sheds over the wetlands from adjoining properties and council open space reserves.

LA ZONE J
North-western interface of Edithvale North adjacent to Atherton Close

Existing conditions

- Reserve between wetland perimeter fence and residential properties is approximately 30 to 40 metres wide. The lack of existing vegetation between housing and the wetland increases their visual prominence, impact on wetland habitat values and visitor amenity values on the existing maintenance access track.
- The bird habitat values are similar to those described in LA Zone I, although there is generally a lack of indigenous overstorey vegetation at all through this zone.

Revegetation Prescription Aims for LA Zone J

- Softan the interface between the housing and the wetlands by planting additional scattered indigenous overstorey trees and selective red-strey.
- As part of the new path alignment along the northern boundary, revegetation is to retain selective views of the wetlands.

LA ZONE K
Western interface with Centre Swamp Drain and adjoining open space

Existing conditions

- Narrow reserve of approximately 20 metres between fence and Centre Swamp Drain.
- Viewing platform with expansive views over the northern wetlands to the Dandenongs in the east.
- Bird habitat values are similar to those described in LA Zone J.

Revegetation Prescription Aims for LA Zone K

- Plant additional scattered indigenous overstorey trees to establish a link with the reserve vegetation in Rosedale Golf Course.
- As part of the proposed path connecting the viewing platform to the maintenance track around the north of the wetland, ensure revegetation works retain selected views over the wetland whilst providing a screening function.


LA ZONE L
Dog pond area on the western boundary

Existing conditions

- The dog pond is saline with large exposed sand areas that have no vegetation.
- The intent of the dog pond reducing the incidents of people walking dogs through the wetlands is a positive initiative.
- Threatened and notable species in this zone include Flame Robin and Black-billed Dotterel. Despite the lack of fencing a few wetland species use the saline ponds including the Black-billed Dotterel and Masked Lapwing and the small marsh next to the saline pond supports Spotted Fat-venter and Golden-headed Cuckoo (Birds Australia, 2004).

Revegetation Prescription Aims for LA Zone L

- Improve the landscape and community educational values of this zone with additional appropriate scattered overstorey tree planting in the terrestrial zone and additional planting.
- Retain the community and dog owner use of this zone, however, where possible establish saltmarsh vegetation to the perimeter.



LA ZONE M
Edithvale North Wetlands (modified ponds)

Existing conditions

- Constructed in the late 1980s.
- Northern ponds are deeper than Edithvale South (LA Zone N) and have intercepted saline ground water.
- This area of the wetland is characterised by deep ponds fringed by tall reeds, and are important for many species, most notably Blue-tailed Duck and Masked Duck which favour the deeper water. This is particularly important in late summer when these ponds can hold numbers of Australian Shearwaters and Hareheads and occasionally Common Greenshank along the edges. Latham's Snipe will feed along the marshy banks, except where there is Tall Marsh and can be flushed from the grassy banks. Flame Robins frequent the area in winter where they prefer the open conditions such as bare wetland edges and shorter grass (Birds Australia, 2004).

Revegetation Prescription Aims for LA Zone M

- Increase habitat diversity with species enrichment planting to the wetlands and some revegetation of terrestrial areas with Plains Grassland/Grassy Woodland vegetation.
- Monitor and control Spiny Rush which is present in low numbers.
- Monitor water quality in conjunction with vegetation to ensure diversity of vegetation is retained and eutrophic conditions do not occur.


LA ZONE N
Edithvale South Wetlands


Existing conditions

- Purchased by DVA in the mid-1970s.
- Relatively natural morphology, subject to a few minor modifications including some deepening and creation of small inlets.
- Peat layer remains in-lieu in the wetland.
- These ephemeral wetlands are extremely important for migratory waders, especially when water levels are receding and damp mud is exposed. All wader species utilise the whole wetland area, though Latham's Snipe stay from the cover of marshy vegetation (Birds Australia, 2004).


Revegetation Prescription Aims for LA Zone N

- Maintain seasonal wetting and drying cycle.
- Monitor water quality in conjunction with vegetation to ensure diversity of vegetation is retained and eutrophic conditions do not occur.
- Biologise the diversity of frog populations in the wetland, in addition to the anura habitat values.
- Establish Swamp Scrub along some of the wetland margins where it will not impact on wader bird habitat values to improve the diversity of habitat values, and be representative of the former vegetation regime of Carrum Carrum Swamp.





1 SKETCH SECTION THROUGH LANDSCAPE ZONE SA AND SL
Open space on the north side of Seaford Wetland
Not to scale



LA ZONE SB
Rural Land, with potential to be acquired by Council


Existing Conditions

- Freshland land currently grazed, with potential to be acquired by Frankston City Council.
- There is a combination of Pines and introduced vegetation and scattered remnant vegetation including Red Gum, Swamp Gums and Banksia along the eastern boundary.
- The dominant habitat type in this area is introduced pasture which is favoured by flocks of Flame Robins during the winter as they prefer the open and short grassy conditions, along with other species including Straw-necked and Australian White Ibis which prefer grazing in an open habitat (Birds Australia, 2004).

Revegetation Prescription Aims for LA Zone SB

- Create freshwater wetlands (ie Herb-rich Plains Grassland) in the old creek course depressions through a combination of revegetation and increasing seasonal freshwater flows.
- Retain the majority of the zone as open slashed pasture with some scattered low shrubs and field layer species from Swamp Scrub EVC for shelter.
- Retain and improve the remnant Damp Sands Herb-rich Woodland vegetation along the eastern boundary with progressive removal of Pines.

2 SKETCH SECTION THROUGH LANDSCAPE ZONE SC
Located immediately south of Seaford North Primary School
Scale 1:100



LA ZONE SC
Western boundary north of Armstrong Road


Existing Conditions

- No views over wetland due to extensive area of Tall Marsh.
- Stormwater outfall at the northern end of zone contributes to permanent wetness of this area and favourable conditions for Tall Marsh.
- The trees provide habitat for overstorey species including Spotted Pardalote and Brown Thornbill. During flowering, the mature trees support large numbers of common nectar-feeding birds including White-plumed and New Holland Honeyeaters and the Red and Little Whitebills (Birds Australia, 2004).

Revegetation Prescription Aims for LA Zone SC

- Increase vegetation between housing and wetland including reinstatement of Swamp Scrub margin. This will also improve wetland habitat and vegetation diversity reflective of the original Caman Caman Swamp.
- Address weed invasion into the Wetland in the northern extent of the zone.

3 SKETCH SECTION THROUGH LANDSCAPE ZONE SD
Located south of Armstrong Road
Scale 1:100



LA ZONE SD
Western boundary from Mona Street to Armstrong Road


Existing Conditions

- Linear reserve in the north section of this zone with predominantly planted native and indigenous overstorey trees. The current position of the shared trail prevents vegetation being established between residential properties and the path. The visual dominance of side boundary fences reduces the recreational value and landscape character in this zone.
- Bird habitat values are similar to those described in LA Zone SC.

Revegetation Prescription Aims for LA Zone SD

- Increase planting of overstorey trees to provide additional screening to residential properties and provide shade to shared path.
- Improve the recreational and landscape character and value of this zone with additional trees for shade and create a natural bushland character.
- Increase diversity of indigenous species on 88 mound over time.

4 SKETCH SECTION THROUGH LANDSCAPE ZONE SD
Located south of Denigh Rd
Scale 1:100



LA ZONE SE
Western boundary from Seaford Primary School to Mona Street

Existing Conditions

- Open water areas close to the wetland perimeter fence. There is a viewing mound with a bench seat just south of Mona Street.
- Bird habitat values are similar to those described in LA Zone SC.

Revegetation Prescription Aims for LA Zone SE

- Increase planting of overstorey trees to provide additional screening to residential properties and provide shade to shared path.
- Improve the recreational and landscape character and value of this zone with additional trees for shade and create a natural bushland character.
- Improve the environmental and landscape amenity of the viewing mound, including potential for a screened approach and interpretive information about the birds.
- Improve the environmental and landscape value of grassed overland flow swales on the western side of perimeter fence by planting edge and rush species.

5 SKETCH SECTION THROUGH LANDSCAPE ZONE SA
North west area of the site adjacent to Patterson River Secondary College and Seaford North Primary School



LA ZONE SF
Eastern boundary to Seaford Primary School and proposed residential use

Existing Conditions

- Established trees including a mix of native and introduced trees including Cypress trees.
- Path is close to the wetland perimeter fence and affords good views northwards over wetland.
- Bird habitat values are similar to those described in LA Zone SC.

Revegetation Prescription Aims for LA Zone SF

- Retain established vegetation as a buffer between proposed residences and the wetland.
- Screen the path from the wetland to potentially improve bird habitats and in an area of the wetland that currently has low presence of bird species.



LA ZONE SI
Eastern boundary to residential courts including Rossett and Bennett Courts

Existing Conditions

- Housing adjacent, poor visibility and access.
- Stormwater outfall creates a freshwater meadow that has good habitat values at the end of Rossett Court. The area adjoining this is slightly elevated and provides good views over the wetland.
- Bird habitat values similar to those described in LA Zone SB, although there are no dense plantings of Coast Banksia in this zone.

Revegetation Prescription Aims for LA Zone SI

- Improve the landscape character by selective thinning and removal of some trees, and strengthening the low shrub and field layer species to establish an open woodland character.
- Provide seating viewing area over the freshwater meadow, along with interpretation of the localised change in vegetation in this area.
- Increase wetland habitat diversity and selective screening with planting of Swamp Scrub along the wetland margin.

LA ZONE SJ
Eastern boundary from Bennett Court North to Armstrong Road

Existing Conditions

- This zone has been largely revegetated over recent years with dense plantings of predominantly indigenous species.
- The wider reserve in this zone is not apparent to the shared trail user, as revegetation works have screened views into and out of this larger area of open space.
- Bird habitat values are similar to those described in SH.

Revegetation Prescription Aims for LA Zone SJ

- Achieve a scattered open woodland character of the Damp Sands Herb-rich Woodland.
- Increase wetland habitat diversity and selective screening by planting Swamp Scrub EVC along the wetland margin.

LA ZONE SK
Eastern boundary to Wells Road

Existing Conditions

- Remnant Damp Sands Herb-rich Woodland along the eastern boundary of this zone, which contains remnant vegetation including a large number of large remnant Swamp Gum, Red Gum and Coast Banksia. There are extensive shelter belt plantings of Pines amongst this remnant vegetation.
- The eastern extent of this zone contains the best quality woodland habitat for bush birds with all types of bush bird fauna represented. It is the only site where Red-browed Finches, Yellow-headed Honeyeaters and Striped Pardalotes occur. The variety of microhabitats allows for a range of different foraging requirements and Spotted Pardalotes, Eastern Yellow Robins and Dusky Woodswallows are all known to nest in this area which may be important locally as a site where bush birds can breed. The major threats to the bird habitat values are motorbike and BMX bike access and weed invasion (Birds Australia, 2004).

Revegetation Prescription Aims for LA Zone SK

- Restore the Damp Sands Herb-rich Woodland vegetation present along the eastern boundary with progressive removal of existing Pines and allowing natural regeneration of indigenous species, in preference to revegetation.
- Expand the area of Damp Sands Herb-rich Woodland to the wetland perimeter to establish a natural connection between the open woodland and the wetland.
- North of the levee bank only, reintroduce fringing Swamp Scrub between the wetland margin and the Damp Sands Herb-rich Woodland vegetation. South of the levee bank the wetland margins are to remain free of Swamp Scrub.

LA ZONE SL
Seaford Wetland, north east-west access bank/levee bank

Existing Conditions

- The area seasonally inundates.
- There is no specific description of bird habitat values for this zone. It is anticipated if the area is restored to a similar condition to LA Zone SA, the bird habitat values will be similar.

Revegetation Prescription Aims for LA Zone SL

- Minor adjustments to hydrological regime to allow seasonal wetting of this area to an approximate depth of 300mm, and active revegetation to establish Brackish Wetland/Herbland which will expand the wetland habitat.
- Undertake Spiny Rush control and ongoing monitoring.
- Establish Swamp Scrub along the east and west margins, leaving the northern extent open for uninhabited arboreal flight path.


LA ZONE SM
Seaford Wetland, south of east-west access bank/levee bank

Existing Conditions


- The northern extent of zone retains its relatively natural morphology.
- The southern part of the zone has been modified with areas excavated to create deeper open water areas. Where this has occurred some ground layer water has been intercepted.
- Extensive areas of Spiny Rush infestation in the north-eastern section of wetland.
- Tall Marsh dominates the north-western part of the wetland.
- The wetlands are noted as being extremely important for migratory waders, especially when water levels are receding and damp mud is exposed (usually December to February). All wader species recorded at Seaford Wetlands use this zone, and these include Latham's Snipe, Whiskered Tern, Common Greenshank, Red-necked Stint, Sharp-tailed Sandpiper, Spotted Crake and Rock-kneed Oystercatcher. Latham's Snipe are only recorded in the southern area of this zone. This zone is also important habitat for ducks, swans and other waterbirds such as herons, egrets and spoonbills. The ponds that are most saline appear to support a reduced number of species and individuals. The extensive areas of Tall Marsh provide ideal habitat for Little Grasshoppers, Clamorous Reed-Warblers and Golden-headed Cuckoos (Birds Australia, 2004).

Revegetation Prescription Aims for LA Zone SM


- Minor improvements to the hydrology to achieve desired seasonal wetting and drying cycles.
- Control Spiny Rush and re-vegetate Brackish Aquatic Herbland fringed by Brackish Wetland/Herbland in the eastern areas of the wetland.
- Control other high priority weed species and regularly monitor for weed infestations.
- Increase species diversity in freshwater wetland communities.
- Provide adequate east west fire breaks to control wildfire and pest fire outbreaks.
- Control predators, particularly foxes, dogs and cats.




5A SKETCH SECTION THROUGH LANDSCAPE ZONE SA
EXISTING CONDITIONS - South of Mona Street
Not to scale



5B SKETCH SECTION THROUGH LANDSCAPE ZONE SA
PROPOSED REVEGETATION - South of Mona Street
Not to scale



6 SKETCH SECTION THROUGH LANDSCAPE ZONE SM
South eastern area of Seaford Wetland
Scale 1:100



7 SKETCH SECTION THROUGH LANDSCAPE ZONE SM AND SK
Eastern side immediately north of Armstrong Road
Not to scale



8 SKETCH SECTION THROUGH LANDSCAPE ZONE SL AND SK
Eastern side north of the central levee bank
Not to scale

RAMRSAR MANAGEMENT PLAN GOAL

To maintain and enhance the natural and cultural heritage of the wetlands and associated terrestrial areas through effective management and active community ownership.


RAMRSAR MANAGEMENT PLAN OBJECTIVES
relevant to the Revegetation Prescription Project

- To maintain and enhance the diversity of wetland vegetation, where practical, to its pre-European state, except where other needs (eg. keeping fire risk at acceptable levels) dictate an alternative approach.
- To maintain and enhance the diversity and abundance of fauna in the wetland.
- To prevent wildfire and pest fires, and to use fire for vegetation and fauna habitat management, consistent with protection of property and visitors, and in a manner that does not start a pest fire.
- To ensure that landscaping is done in a manner that resembles the floristic and structure of pre-European vegetation of the area (using plant material of local provenance), including extensive open areas, and that landscaping products and enhances the hydrogeological and ecological functioning of the wetlands.
- To ensure that community use of the wetlands is consistent with maintaining ecological values by catering for a suitable range of activities in areas where the wetlands and their fauna will not be disturbed.
- To provide opportunities for environmental education for schools, tertiary institutions, special interest groups and general public, consistent with the protection and interpretation of the natural and cultural heritage of the wetlands.
- To generate adaptive information on the key environmental variables to ensure well documented and adaptive management.

REVEGETATION PRESCRIPTIONS PROJECT AIMS:

- reinforce the revegetation works that have been undertaken over the years with some changes to landscape a relatively natural open woodland character in the terrestrial areas; establish some sheltered areas of the fringing Swamp Scrub on the wetland margins whilst retaining extensive areas of open woodland character;
- continue the Spiny Rush control program and control of other weeds in the wetland and undertake species enrichment where required;
- improve the avifauna values with additional vegetation in screen adjoining activities from the wetland and provide a diversity of habitats for a range of birds;
- protect the significant vegetation communities;
- continue to encourage community involvement in revegetation works and in particular recognise the ongoing work and involvement of the Friends of Edithvale-Seaford Wetlands; and
- create an inviting natural setting which reflects the environmental significance of the wetlands and engenders a respect for these values from visitors and local residents.


LAND OWNERSHIP DIAGRAM



VEGETATION COMMUNITIES KEY



DRAWING KEY



Appendix 14 Edithvale-Seafood Ramsar Wetlands Desktop Cultural Heritage Assessments for the Edithvale Section (Heritage Insight 2016a) and Seafood Section (Heritage Insight 2016b).

This page is intentionally blank.

Edithvale-Seaford Ramsar Wetlands
Edithvale Section
Desktop Cultural Heritage Assessment



Cultural Heritage Advisor: Bianca Di Fazio

Author: Renee McAlister and Bianca Di Fazio

June 10, 2016

**Heritage
Insight**
Pty Ltd



Heritage Insight P/L
53 Stubbs Street Kensington VIC 3031

PHONE: +61 3 9376 6569 INQUIRIES: archaeology@heritageinsight.com
FAX: +61 3 9376 6526 WEB: www.heritageinsight.com

ABN: 73 116 621 884 ACN: 116 621 88

Acknowledgements

Heritage Insight Pty Ltd would like to acknowledge and thank the following people for their assistance and participation in this study:

Ecology Australia

Darren Quin

Jonathan Ricciardello

Heritage Insight

Shannah Anderson Background Research

Katarina Audy GIS Mapping

Traditional Owner Groups

Wurundjeri Tribe Land Compensation and Cultural Heritage Council Inc (WTCCHC)

Boon Wurrung Foundation (BWF)

Bunurong Land Council Aboriginal Corporation (BLCAC)

Bunurong Land and Sea Association (BLSA)

Heritage
Insight
Pty Ltd



Heritage Insight P/L
53 Stubbs Street Kensington VIC 3031

PHONE: +61 3 9376 6569 INQUIRIES: archaeology@heritageinsight.com
FAX: +61 3 9376 6526 WEB: www.heritageinsight.com

ABN: 73 116 621 884 ACN: 116 621 88

Disclaimer

The information contained in this Cultural Heritage Assessment has been compiled from the standard heritage database sources and is accurate as far as Heritage Insight Pty Ltd is aware. However, within the timeframes available for technical heritage reporting, it is not possible to carry out comprehensive research of all published or unpublished manuscripts, journals, maps or oral history which may pertain to the study area. No responsibility can be taken for errors or omissions in primary and secondary source material cited in this report. Any opinions expressed in this report are those of Heritage Insight Pty Ltd only.

Copyright Notice

This report is copyright. Apart from any fair dealing for the purposes of private study, research, criticism or review as permitted under the Copyright Act, no part may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, whether electronic, mechanical, photocopying, recording or otherwise without prior written permission. Enquiries should be made to Heritage Insight Pty Ltd.

Executive Summary

This report outlines the results of an investigation into the heritage values of the Edithvale-Seaford Ramsar Wetlands, and has been conducted as part of the larger Ramsar Management Plan undertaken by Ecology Australia for Melbourne Water.

The purpose of this report is to identify the heritage values associated with the Edithvale section of the Edithvale-Seaford Ramsar wetlands, formulate a site prediction model and provide recommendations for the ongoing management of cultural heritage issues within the area.

Section 2 outlines information about the formation and hydrology of the Edithvale section of the Edithvale-Seaford Ramsar wetlands along with discussing the geology and geomorphology of the study area and providing information on the climate and available flora, fauna and stone resources within the study area prior to European contact. The Edithvale wetland basin is the last and deepest remnant of Carrum Swamp. This area is now managed by Melbourne Water as part of a larger drainage and storm water management system. The area is also home to a wide variety of bird life. The geology of the activity area characterised by coastal lagoon deposits, representing formerly inundated swampland.

Section 3 presents a review of previous archaeological assessment within the study area and surrounds. This information is combined with the environmental data and the land use history to assist in the formulation of a site prediction model. A search of the VAHR shows that the study area contains no registered Aboriginal sites. A review of previous archaeological research concludes that the study area is unlikely to contain registered Aboriginal sites as it is located within the boundary of the former Carrum Swamp, and Aboriginal sites are more likely to be located around the margins of the swamp on elevated sandy landforms.

Section 4 details the land use history of the study area and provides an ethnography for the *Bun wurrung* people. A review of the land use history demonstrates that the study area is located within a landform that has undergone substantial modification and disturbance over the years; from being part of Carrum Swamp, through drainage and utilisation for farming, to its role in managing drainage and storm water today.

Due to the fact that the study area is located in an area that was formerly within the wetlands of Carrum Swamp it is unlikely to contain Aboriginal cultural material, as Aboriginal activity was more likely to have been focused around the margins of the swamp on more elevated sandy landforms. Additionally the land has undergone substantial disturbance which is known to disturb and destroy Aboriginal sites. As such the study area is considered to contain extremely low archaeological potential. A site prediction model is presented in Section 5.1.

Section 5.2 outlines the responsibilities and requirements under the Aboriginal Heritage Regulations 2007. It notes that planned projects will need to be assessed on an individual basis to determine if they require a CHMP or if an archaeological due diligence assessment will be sufficient to manage any potential heritage management issues.

Table of Contents

<i>Executive Summary</i>	<i>i</i>
<i>Part One: Assessment</i>	<i>1</i>
<i>1.0 Introduction</i>	<i>1</i>
<i>2. Existing Conditions, Environment and Geology</i>	<i>4</i>
2.1 Existing Conditions.....	4
2.2 Landforms and Geomorphology of the Study Area.....	6
2.3 Resources Available to Aboriginal People within the Study Area.....	8
<i>3. Review of Previous Archaeological Research</i>	<i>11</i>
3.1 VAHR search.....	11
3.2 Previous Archaeological Assessment within 2km of the Study Area.....	14
<i>4. Land Use History and Ethnography</i>	<i>19</i>
4.1 Historical and Ethno-Historical Accounts in the Geographic Region.....	19
4.2 Land-Use History of the Study Area.....	22
<i>5. Conclusions and Recommendations</i>	<i>32</i>
5.1 Site Prediction Model.....	32
5.2. Recommendations and Obligations under the Aboriginal Heritage Act.....	33
<i>References</i>	<i>34</i>
<i>Appendix 1: Division 5 of the Aboriginal Heritage Regulations 2007</i>	<i>37</i>
<i>Appendix 2: Glossary</i>	<i>45</i>
<i>Appendix 3: OAAV Practice Note for Determining Significant Ground Disturbance</i>	<i>50</i>

Maps

Map 1: Location of the study area.....	3
Map 2: Map showing existing conditions.....	5
Map 3: Map showing geology of the study area and surrounds.....	7
Map 4: Map showing 1750s EVC mapping of the study area and surrounds.....	9
Map 5: Map showing location of registered Aboriginal places within 200m of the study area.....	13

Figures

Figure 1. 1840 map by William Thomas indicating routes taken on his travels with <i>Bun wurrung</i> tribes. (Reproduced from Presland 1994, p. 74).....	21
Figure 2: Map showing location of Edithvale-Seaford Wetlands in blue and extent of the former Carrum Swamp in grey.	23
Figure 3: 1871 Parish Map of Carrum Swamp (source: TROVE).....	24
Figure 4: 1945 map showing location of study area and surrounds.....	26
Figure 5: 1956 map showing location of study area and surrounds.....	27
Figure 6: 1963 map showing location of study area and surrounds.....	28
Figure 7: 1972 map showing location of study area and surrounds.....	29
Figure 8: 1980 map showing location of study area and surrounds.....	30
Figure 9: 2009 map showing location of study area and surrounds.....	31

Tables

Table 1: Summary of registered Aboriginal Places within the geographic region	11
Table 2: Summary details for Aboriginal Places within 2km of the activity area (sites within 200m shown in orange).....	12
Table 5: Relevant archaeological studies within 2km of the activity area	15

Abbreviations

ACHRIS – Aboriginal Cultural Heritage Register and Information System
ASL – Above Sea Level
BLCAC - Bunurong Land Council Aboriginal Corporation
BLSA – Bunurong Land and Sea Association
BWF – Boon Wurrung Foundation
CHA – Cultural Heritage Advisor
CHMP – Cultural Heritage Management Plan
DPC – Department of Premier and Cabinet
EMP – Environmental Management Plan
EVC – Ecological Vegetation Class
LDAD – Low Density Artefact Distribution
LGM – Last Glacial Maximum
OAAV – Office of Aboriginal Affairs Victoria
PAS – Potential Archaeological Sensitivity
PSP – Precinct Structure Plan
RTP – Radial Test Pit
STP – Shovel Test Pit
TP – Test Pit
UFZ – Urban Floodway Zone
VAHR – Victorian Aboriginal Heritage Register
WTLCCHC – Wurundjeri Tribe Land Compensation and Cultural Heritage Council Inc

Please note that all maps and plans in this CHMP are prepared using Victorian Government Standard GDA94 MGA (Zone 55) coordinates.

Part One: Assessment

1.0 Introduction

This report outlines the results of an investigation into the heritage values of the Edithvale-Seaford Ramsar Wetlands, and has been conducted as part of the larger Ramsar Management Plan undertaken by Ecology Australia for Melbourne Water.

The purpose of this report is to identify the heritage values associated with the Edithvale section of the Edithvale-Seaford Ramsar wetlands, formulate a site prediction model and provide recommendations for the ongoing management of cultural heritage issues within the area.

The Edithvale-Seaford Wetland is the largest natural wetland of its type in the Port Phillip and Westernport basins. It is all that remains of Carrum Swamp, which once covered more than 5,000 hectares from Mordialloc in the north to Frankston in the south. This site comprises two separate wetland areas, Edithvale and Seaford. This report will deal with the Edithvale section of the Edithvale-Seaford Wetlands.

The wetlands were listed on 29 August 2001 under the Ramsar Convention, as Ramsar site 1096, Australia's 11th site, in recognition of their international importance, and specifically because they:

- Are the last remaining examples of the Carrum Swamp, containing a variety of permanent and seasonal, freshwater and saline wetlands,
- Support populations of the Australasian bittern, considered to be of state significance and threatened in Victoria,
- Support more than 1% of the East Asian – Australasian Flyway population of sharp-tailed sandpipers (or over 2000 birds) in up to one year in three; and
- They are also considered to be of exceptional significance as examples of cost-effective management of wetlands in an urban setting to provide conservation benefits, manage storm water, and encourage environmental education and research.

This assessment has utilised standard heritage databases and historic sources to provide a detailed land use history of the study area. A review of previous archaeological assessments has been completed and this along with geological and environmental information has been utilised to formulate a site prediction model and establish the heritage values of the study area.

Location of the Study Area

The Edithvale site covers an area of 103 hectares.


Section 1 is the northern part of the Edithvale site. It is comprised of parcels 1\TP820840, 1\TP131999, 1\TP225777, 2\TP225777, 1\TP83139, 1\TP820843, 2\TP820843, 1\TP385644, 1\TP414444 and 1\TP82835 (DEPI 2013).

Section 2 is the southern part of the Edithvale site. It is comprised of parcels 1\TP138507, 1\TP132070, 1\TP366503, 1\TP370109, and 1\TP95924 (DEPI 2013).

The Edithvale wetlands are located in the suburbs of Aspendale Gardens (section 1) and Chelsea Heights (section 2) approximately 1.6km south of Mordialloc Creek and 2.7km north of the Patterson River, in the Parish of Lyndhurst, City of Kingston approximately 28km south of the Melbourne CBD. They are surrounded by residential development, with sporting facilities lying to the west and Edithvale road running through the centre.

Land Manager

The land is managed by Melbourne Water.



Map 1: Location of the study area

2. Existing Conditions, Environment and Geology

2.1 Existing Conditions

The study area is the Edithvale section of the Edithvale Seaford Wetlands. These wetlands represent the remnants of the former Carrum Swamp that extended over 5000 hectares, stretching along the eastern shore of Port Phillip Bay from Mordialloc to Frankston.

The Edithvale Wetland basin is the last and deepest remnant of Carrum Carrum Swamp, and it still has an intact underlying peat layer in places. The wetland system is usually divided into Edithvale North and Edithvale South for management purposes, and hydrology differs between and across these systems (KBR 2009).

Edithvale South Wetlands have retained most of their natural morphology (GHD 2005). The system consists of one main shallow pool (usually less than 1 m in depth) and three smaller, deeper and more permanent drought refuge pools. Apart from the drought refuge pools, the majority of the system is managed such that it is allowed to dry out over late summer and autumn. Drying of the wetlands facilitates the growth of salt club-sedge (*Bolboschoenus caldwellii*), the plant species that produces and sustains the peat layer in the system. The system is seasonally flooded by stormwater from the local residential areas (KBR 2009).

The Edithvale North Wetlands is a partially constructed system. Some of the wetlands area has recreated itself, while a series of pools have been constructed to assist with water management. The pool to the north of Edithvale is a shallow constructed pool that is seasonally inundated. There are a series of other pools to the north that reach through the peat layers and into the sandy substrate. There are a series of weirs that control water movement between the pools. These pools are filled through stormwater drainage and overflow which enters the system through sediment ponds (KBR 2009).

The Edithvale section of the wetlands is utilised for both drainage purposes, stormwater management and the outer areas are used as an open public space popular with local dog walkers, bird watchers and for other recreational use. The wetlands are home to a significant population of birdlife.



Map 2: Map showing existing conditions

2.2 Landforms and Geomorphology of the Study Area

DEPI mapping shows that the study area is located within the footprint of the former Carrum Swamp (Figure 2) and in fact is a remnant of that former swamp. The Carrum Swamp was an extensive freshwater wetlands stretching between Mordialloc and Frankston and as far east as Bangholme. The water from the Carrum Swamp and the Dandenong and Eumemmerring Creeks drained into Port Phillip Bay via Mordialloc and Kananook Creeks. This wetland has now been drained via the Mordialloc and Dandenong Creeks/drains. The current alignment of the Mordialloc drain was formerly the channel of the Mordialloc Creek, and the current alignment of Dandenong Creek has been cut through the former Tea Tree Swamp (upper part of Carrum Swamp).


Carrum Swamp was drained in the 1870s by cutting channels through natural waterways and wetlands to increase discharge into Port Phillip Bay (Murphy & Dugay-Grist 2008). Patterson's Cut (now Patterson River) was excavated in 1876 to assist with drainage of the swamp. The Carrum Swamp along with the Koo Wee Rup Swamp provided a barrier to early European exploration and settlement of the region and drainage works began in order to drain the land and provide more accessible farmland. The Carrum Swamp would have represented a valuable resource for Aboriginal people provide freshwater and access to a wide range of flora and faunal resources.

The study area is located in the geomorphic region known as the Barrier Complexes – Discovery Bay, Gippsland Lakes. This is an extensive geomorphic region covering part of eastern Victoria. The landform of the wetlands area has been described as part of the Carrum Lowlands or Sunklands.

Geological mapping of the region shows that the study area is located on Coastal Lagoon Deposits (Qg), which are described as dark grey to black silt and clay, and represent deposits that were previously inundated by swamp waters.

This area has undergone significant change overtime as a result of fluctuating sea levels, climate change and irrigation programs. Between 14,000 and 10,000 years ago climatic conditions changed to become warmer and wetter resulting in rising sea levels. Port Phillip Bay flooded around 10,000 years ago though formation of the current coastline did not stabilise until 4,000 years ago. Inundation occurred over several thousand years, with French and Phillip Islands not formed until the highest sea level of 5,000-6,000 years ago (Coutts 1977). As current dune barrier systems developed, previous discharge points for the region's rivers and creeks were blocked resulting in a series of freshwater swamps collectively known as Carrum Swamp.

The Carrum Swamp originally stretched from Port Phillip Bay across to Western Port Bay.




Map 3: Map showing geology of the study area and surrounds

2.3 Resources Available to Aboriginal People within the Study Area

Plant Resources and Pre-Contact Vegetation

The pre-1750 Ecological Vegetation Community (EVC) present within the study area has been identified as EVC 125 (Plains Grassy Wetland) (Department of Environment and Primary Industries Biodiversity Interactive Map, accessed 02/02/2016).

Plains Grassy Wetland is described as usually treeless, but in some instances can include sparse River Red Gum (*Eucalyptus camaldulensis*) or Swamp Gum (*Eucalyptus ovata*). A sparse shrub component may also be present. The characteristic ground cover is dominated by grasses and small sedges and herbs. The vegetation is typically species-rich on the outer verges but is usually species-poor in the wetter central areas. As the study area would have been either partially or completely inundated, non-aquatic vegetation would have been restricted to the edges of the swamp.



Map 4: Map showing 1750s EVC mapping of the study area and surrounds

Information on Fauna of the Activity Area

A number of animals, water birds, fish and eels would have been present within the study area and the wider region of Carrum Swamp and are likely to have been hunted by Aboriginal people. However, due to the inundated nature of the study area, Aboriginal activity would have been focused around the swamp margins.

Water Resources

Carrum Swamp would have been the closest fresh water source to the study area

Stone Resources

The lithology of the proposed activity area and surrounding region indicates that quartz would be a highly accessible stone material available to Aboriginal people for the manufacture of stone artefacts. Outcrops of basalt occur at Berwick and to the north siltstone and mudstone exists. Silcrete and chert, which were materials highly sought after by Aboriginal people, were probably obtained from coastal areas such as the Mornington Peninsula (Smith 1991)

3. Review of Previous Archaeological Research

The Victorian Aboriginal Heritage Register (VAHR), accessed through Aboriginal Cultural Heritage Register and Information System (ACHRIS), was searched to identify any previously registered Aboriginal Places within a 2km radius of the study area. This search shows that one CHMP has been conducted within the study area (Murphy & Dugay-Grist 2008). No recorded Aboriginal sites have been located within the study area.

3.1 VAHR search

A search of the Victorian Aboriginal Heritage Register identified that there are 13 registered Aboriginal Places within a 2km radius of the study area, comprising a total of 19 components (Table 1). Of these 13 sites, four were located within 200m of the study area (Table 2). The majority of sites in the area are small deposits of stone artefacts located in both surface and sub-surface contexts. Artefact raw material in these sites is primarily quartz and quartzite, with some silcrete. These sites have all been located on the lowland plain on the former swamp landform and likely represent transitory use of the landscape rather than substantial long term campsites. It should also be noted that contour mapping shows all of these sites are located at a higher elevation than the study area.

Table 1: Summary of registered Aboriginal Places within the geographic region

Component Type	Frequency (No.)	Frequency (%)
Artefact Scatter	9	47
Low Density Artefact Distribution	9	47
Scarred Tree	1	6
Total Components	19	
Total Registered Places	13	

Table 2: Summary details for Aboriginal Places within 2km of the activity area (sites within 200m shown in orange).

VAHR ID	Site Name	Site Type	Description
7921-0038	CHELSEA HEIGHTS 1	Artefact Scatter	Six surface quartz artefacts on a lowland plain
7921-0039	CHELSEA HEIGHTS 2	Artefact Scatter	Small surface scatter of stone artefacts on a lowland plain
7921-0040	CHELSEA HEIGHTS 3	Artefact Scatter	Eight surface quartz artefacts on a low land plain
7921-0266	KEYS 1	Scarred Tree	Red gum tree with cultural scar in poor health on lowland plain
7921-1379	Chelsea Heights 4	Artefact Scatter	Two artefacts retrieved from a depth of 600-700mm, watertable encountered at 850mm.
7921-1391	44 First Avenue 1	Artefact Scatter	Five sub-surface stone artefacts retrieved from a depth of 600-700mm on a former swamp landform, watter table encountered at 800mm.
7921-1392	44 First Avenue 2 IA	Artefact Scatter	One isolated quartzite artefact retrieved from a depth of 600mm, watertable encountered at 600mm.
7921-1393	44 First Avenue 3 IA	Artefact Scatter	One isolated silcrete artefact retrieved from a depth of 600mm, watertable encountered at 600mm.
7921-1396	Chelsea Heights 5	Artefact Scatter	Five sub-surface quartz and quartzite artefacts retrieved from a depth of 400-600mm on a former swamp landform, watter table encountered at 900mm.
7921-1400	Chelsea Heights 6	Artefact Scatter	Two stone artefacts retrieved from a depth of 500 -700mm, water table encountered at 800mm.
7921-1447	58 Fourth Avenue Chelsea Heights LDAD	Low Density Artefact Distribution	Five stone artefacts
7921-1520	40 Ella Grove, Chelsea LDAD	Low Density Artefact Distribution	One isolated artefact
7921-1530	58-60 Berry Ave LDAD	Low Density Artefact Distribution	Three stone artefacts



Map 5: Map showing location of registered Aboriginal places within 200m of the study area.

3.2 Previous Archaeological Assessment within 2km of the Study Area

A number of reports have been undertaken within the geographic region. The results of relevant regional and localised studies, including CHMPs are presented in Table 3. The results of these reports are utilised to assist in the formulation of the site prediction model.

Discussion of Previous Archaeological Work

DEPI mapping shows that the study area is located within the footprint of the former Carrum Swamp (Figure 2) and in fact is a remnant of that former swamp. The Carrum Swamp was an extensive freshwater wetlands stretching between Mordialloc and Frankston and as far east as Bangholme. As discussed above, Aboriginal sites are commonly found around the margins of the Swamp and on elevated ground within the area. However, recent archaeological work has also located sites underneath swamp deposits. CHMP 11958 (Kennedy, Wheeler & Foley 2012) challenged the assumption that Aboriginal sites will not be found within the former swamp, as they identified archaeological deposits within a sandy dune layer located underneath the swamp deposits which is believed to predate the inundation of Carrum Swamp, dating the finds to at least 7000 years old. However, this sandy layer has not been identified in other CHMPs conducted nearby.

More broadly however, archaeological work conducted in Edithvale and surrounds has shown that Aboriginal archaeological sites located within the margins of the former Carrum Swamp are extremely rare. While Carrum Swamp was known to have been an important resource for Aboriginal people and was heavily utilised, Aboriginal activity was focused around the margins of the swamp on elevated land forms. Low lying floodplain and frequently inundated areas made poor camp sites and were more rarely utilised by Aboriginal people, who preferred higher well drained areas adjacent to fresh water (Murphy & Dugay-Grist 2008). As the reviews in Table 3 below demonstrate, this landform has been shown to have extremely low archaeological potential.

Table 3: Relevant archaeological studies within 2km of the activity area

Study Name	Investigation Type	Results
<p>Residential Subdivision 1 Bristol Avenue, Edithvale</p> <p>(White 2015)</p>	CHMP	<p>This CHMP (13482) was conducted for a small residential subdivision. The desktop assessment noted that the activity area was dominated by two landforms, the coastal barrier dune system and the coastal lagoonal deposits that form the wetlands of the Carrum Swamp. For this reason this it was considered that the area contained potential for the discovery of shell middens and artefact scatters. The standard assessment noted that the activity area had undergone high levels of disturbance through its land use history of residential development. The activity area is considered to retain no archaeological potential.</p>
<p>Subdivision Lot 1, 2, 3 TP49429 and Lot A PS605114 Springvale Road Aspendale Gardens</p> <p>(Murphy & Thomson 2015)</p>	CHMP	<p>This CHMP (13226) was prepared for a large residential subdivision. This land was previously zoned green wedge and had been subject to minimal disturbance from agricultural use. The desktop assessment noted that the activity area was within the boundaries of the former Carrum Swamp and as such contained low potential for the discovery of Aboriginal cultural heritage material. The complex assessment comprised the excavation of three 1m² test pits and 31 shovel test pits. Sub-surface testing identified that the area contained very little topsoil and comprised dense clay layers suggesting that the area had previously been permanently inundated by the Carrum Swamp. No Aboriginal cultural heritage material was located and no areas of archaeological potential were identified.</p>
<p>Proposed Residential Development at 58-60 Berry Avenue, Edithvale</p> <p>(Mitchell 2014)</p>	CHMP	<p>This CHMP (13201) was conducted for a small residential subdivision. The desktop assessment noted that the activity area was within the boundary of the former Carrum Swamp and also includes a section of the Koo Wee Rup Plain and foreshore dune deposits. This intersection of landforms would have provided a range of resources for Aboriginal people. The complex assessment comprised the excavation of a 1m² test pit and nine shovel test pits. This revealed a low density artefact distribution of three silcrete flakes (VAHR 7921-1530).</p>
<p>40 Ella Grove, Chelsea: Residential Subdivision</p> <p>(Matic 2014)</p>	CHMP	<p>This CHMP (12961) was conducted for a small residential subdivision. The desktop assessment noted that the activity area had been impacted by residential development. The complex assessment comprised the excavation of a 1m² test pit and nine shovel test pits. One Aboriginal place was located during testing, VAHR 7921-1520, a single silcrete flake. This flake was located on a section of a former dune landform located close to the boundaries of the former Carrum Swamp.</p>

Study Name	Investigation Type	Results
<p>Perry Road and Mordialloc Main Drain: Drainage Infrastructure, Keysborough</p> <p>(Kennedy & Crocker 2013)</p>	CHMP	This CHMP (12053) was conducted for the installation of drainage infrastructure along Perry Road. The site prediction model noted that the activity area contains potential for the discovery of scarred trees and artefact scatters. Sandy rises and dune crests were considered areas of high archaeological potential, while low lying and formerly waterlogged/wetlands had low archaeological potential. The standard assessment noted that the activity area comprised the low-lying plain landform and as such was unlikely to contain Aboriginal sites.
<p>Residential Development, 85 Berry St Edithvale</p> <p>(Murphy & Thomson 2013)</p>	CHMP	This CHMP (12395) was conducted for a residential subdivision. The desktop assessment demonstrated that the activity area was located within the margins of the former Carrum Swamp. The complex assessment comprised a 1m ² test pit, four 50x50cm shovel test pits and three auger holes. Sub-surface testing revealed that disturbance across the top 30cm of topsoil. No Aboriginal cultural heritage material was located.
<p>Springvale Road, Aspendale Gardens and Nursery Development</p> <p>(Green 2013)</p>	CHMP	This CHMP (12851) was conducted for the construction of a commercial nursery. This activity area is located within the former Carrum Swamp. The complex assessment comprised a 1m ² test pit and 21 shovel test pits excavated in a grid across the activity area. Sub-surface testing revealed sediments consistent with former swamp deposits and no Aboriginal cultural material was located.
<p>Drainage Works, Edithvale</p> <p>(Matic, MacCulloch & Falvey 2013)</p>	CHMP	This CHMP (12363) was conducted for proposed drainage works. The desktop assessment noted that while the activity area is located within the margins of the former Carrum Swamp, the complex assessment should aim to identify if the activity area contains the landform identified in CHMP 11958 as sensitive. The complex assessment comprised four 1m ² test pits, one shovel test pit and a 5.6m x 8m machine excavated pit. The soil profile was identified as swampy deposits with a layer of sand identified as Pleistocene dune deposits identified at depths of approximately 80-85cm. No Aboriginal cultural heritage material was located.
<p>58 Fourth Avenue, Chelsea Heights</p> <p>(Dugay-Grist, Cowled & Maher 2012)</p>	CHMP	This voluntary CHMP (12450) was conducted for the sale of land on the Koo Wee Rup plain. The activity area was considered to contain moderate archaeological potential due to its location within a Neogene dune system. The complex assessment comprised the excavation of two 1m ² test pits and seven shovel test pits. Sub-surface testing revealed five stone artefacts, VAHR 7921-1447 in sandy dune deposits.

Study Name	Investigation Type	Results
<p>Sporting Pavilion and Early Years Hub, Edithvale</p> <p>(Matic & Falvey 2012)</p>	CHMP	This CHMP (12313) was conducted for the construction of a sporting facility and childcare centre. The desktop assessment noted that the activity area lay within the footprint of the former Carrum Swamp. The complex assessment comprised of a 1m ² test pit and one 50x50cm test pit. The test pits were excavated down to the water table. No Aboriginal cultural material was located.
<p>142-143 Nepean Highway, Aspendale</p> <p>(Thomas & Howell-Meurs 2012)</p>	CHMP	This CHMP (12048) was conducted for the construction of multiple dwellings on a residential allotment. This area is considered culturally sensitive due to its proximity to the coastline. Sub-surface testing comprised a 1m ² test pit, six shovel test pits and five hand auger probes. Testing revealed the activity area had undergone extensive ground disturbance, all excavations terminated due to the presence of construction rubble and material suspected of being asbestos. No Aboriginal cultural material was located.
<p>44 First Avenue, Chelsea Heights</p> <p>(Kennedy, Wheeler & Foley 2012)</p>	CHMP	This CHMP (11958) was conducted for a residential subdivision. The desktop assessment noted that the activity area was located within the margins of the former Carrum Swamp and as such contained low potential for the discovery of Aboriginal cultural material, which is more commonly located on elevated rises around the margins of the swamp. The standard assessment confirmed the activity area was a broad flat open landform that comprised former swamp land. The complex assessment comprised the excavation of thirteen 1m ² test pits and two 3x3m test trenches. Sub-surface testing identified archaeological deposits within a buried B horizon sand unit underneath an A horizon swamp profile. A total of sixteen stone artefacts were located and recorded as six low density stone artefact scatters VAHR 7921-1391, 7921-1379, 7921-1392, 7921-1393, 7921-1396, and 7921-1400. As the activity area was inundated between 7000 BP until 2000 BP, these sites likely dates to pre-7000 BP, making the sites of high significance due to their rarity and scientific, public and cultural values. The discovery of these sites also challenges the previously held assumption that sites are unlikely to be located within the swamp itself.
<p>Proposed Residential Subdivision at 88 Berry Ave, Edithvale</p> <p>(Schultz & Donati 2011)</p>	CHMP	This CHMP (11883) was conducted for a residential subdivision. The standard assessment noted evidence of ground disturbance associated with the dwelling, drive way and shed however the balance of the property appeared relatively undisturbed. The complex assessment comprised a 1m ² test pit and five shovel test pits. Sub-surface testing revealed a fairly disturbed soil profile and no Aboriginal cultural material was located.

Study Name	Investigation Type	Results
<p>Residential Development 8-10 Maury Road, Chelsea</p> <p>(Kiddell, Watson & Howes 2011)</p>	CHMP	<p>This CHMP (11673) was conducted for a small residential subdivision. The desktop assessment noted the activity area was located on barrier dunes and contained potential for stone artefact scatters. The complex assessment comprised six test pits (1x2m, two 50x50cm and three 30x30cm). Test Pit 1 was excavated to a depth of 2550mm, the other test pits were excavated to depths between 300mm and 1450mm. No cultural heritage material was identified.</p>
<p>Edithvale-Seaford Wetland Discovery Center</p> <p>(Murphy & Dugay-Grist 2008)</p>	CHMP	<p>This CHMP (10081) was conducted within the current activity area. This area is considered sensitive as it is a Ramsar wetlands. The site prediction model noted that elevated land associated with Carrum Swamp should be considered to contain high archaeological potential. The standard assessment noted extremely poor ground surface visibility. The complex assessment comprised four 1m² test pits and two machine transects. The test pits revealed dark peat deposits, identified as former swamp deposits. Shell material was identified within one of the machine transects, however the shell deposits were identified as natural rather than cultural. No Aboriginal cultural material was identified and as the test pits showed the area comprised former swamp deposits, the area was not considered to contain archaeological potential.</p>
<p>An archaeological assessment of a proposed subdivision at Patterson River Country Club</p> <p>(Webb 2000)</p>	Field Survey	<p>This archaeological assessment was conducted for a residential subdivision of a strip of land adjacent to Patterson River. Prior to being drained, the Carrum Swamp had two natural outlets, Mordialloc Creek to the north and Kananook Creek to the south. The Patterson Cut was an artificial opening built in 1879 to assist with the drainage of Carrum Swamp. The field survey noted variable ground surface visibility dependant on vegetation coverage. Areas of good visibility were limited to tracks and bare ground under trees. Some natural shell deposits were identified however no Aboriginal sites were located. As the land was previously covered by Carrum Swamp, it was not considered to contain archaeological potential.</p>

4. Land Use History and Ethnography

4.1 Historical and Ethno-Historical Accounts in the Geographic Region

Indigenous people have likely occupied and utilised the landscape around Melbourne for over 40,000 years, although there is very little known about human occupation and palaeo-environments in the Melbourne Region until 10-15,000 years ago.

Prior to European settlement, the Indigenous population of Victoria was divided into a number of groups. In central Victoria, the languages of each of these groups were dialectically similar, sharing up to 93% of their vocabularies. These dialect groups, named the *Woiworung*, *Bunurong*, *Taungurong*, *Ngurainwurung*, *Wada wurrung* and *Dja dja wurrung*, comprised a 'regional cultural confederacy known as the Kulin' (Barwick 1984, pp.104–105). Members of the Kulin confederacy shared economic and social relationships, common religious beliefs, creation legends and Dreamtime ancestors. Each person also belonged to one of two moieties named after the Dreamtime ancestors, Bunjil, the eaglehawk, and Waa, the crow. Moiety affiliation was determined at birth with the child adopting the moiety of the father's clan. Kulin people were exogamous, meaning they married outside their clan and to a person from the opposite moiety (Barwick 1984, p.106; Clark 1990).

The study area is located within the land of the *Bun wurrung* people. The *Bun wurrung* (Bunurong) were one of four 'dialectal tribes' comprising the East Kulin language group (Barwick 1984). William Thomas, Assistant Protector of Aborigines in the Western Port district (1839 – 1849), spent much of his time travelling with *Bun wurrung* people between his hut at Arthur's Seat, the Aboriginal reserve which he established at Narre Narre Warren during 1841 and the Aboriginal camps around Melbourne (Sullivan 1981, p.25; Cannon 1983). This extensive travel through *Bun Wurrung* territory enabled Thomas to argue that the *Bun Wurrung* claimed "all the country south of the Yarra River, whose creeks and inlets fall into the sea from the Werribee River west to the Tarwin River, east of Cape Patterson" (Thomas papers Vol.7 17/1/1860).

The study area was the country of the *Mayone buluk* meaning 'people of the swamp'. Their territory is thought to have been "Carrum Swamp, the coastal strip at the head of Western Port Bay and the upper portion of the Mornington Peninsular" (Barwick 1984: 177).. The *arweet* (clan head) was identified as Motrungo / Budgery Tom (c. 1797/8-1848). His heir was Buggup / Buckup (c. 1820/3-1848), a corporal in the Native Police Corps (Clark 1990, p.367; Barwick 1984, p.117). The moiety affiliation of the *Mayone bulluk* was Bunjil (Clark 1990, p.367).

Post-contact historical accounts

Much of the available information about traditional *Bun wurrung* culture has been provided by William Thomas, Assistant Protector of Aborigines for the Western Port District. Other sources of information are the journals and letters of G. A. Robinson, Chief Protector of Aborigines (1839 – 1850); A. W. Howitt and early settlers such as Georgiana McCrae and Maurice Meyrick (Bride 1969; Presland 1977; McCrae 1992).

First contact between *Bun wurrung* people and Europeans was with whalers and sealers prior to 1803 (sealing began in Bass Strait in 1798), as European huts were discovered by Grimes at Boneo in January of that year (Rogers 1957). The fact that whalers and sealers had arrived early in the history of the colony is further substantiated by entries in Robinson's journal (26 December 1836 cited in (Plomley 1987, p.405), where he records an account of the early kidnapping of *Bun wurrung* women:

Matilda the VDL native woman pointed out the spot a few miles down the harbour at Point Nepean where she said George Meredith and his crew of sealers stole the native women. The men's names were Brown, Mr. West the master of the schooner, and a man named Billy...said there was plenty of black fellows, some on the Port Phillip side some outside, sea coast. Said the sealers were afraid of the Port Phillip natives. Said they employed her to entice them. George Meredith stole the, I think she said, four women, took them...and then sold them to the [other] sealers there. I am informed that Munro bought one. (G. A. Robinson Journal entry, 26 December 1836, in (Plomley 1987, p.405).

During 1839, the British Colonial Government established an Aboriginal Protectorate in what is now Victoria. A Chief Protector, G.A. Robinson and four Assistant Protectors were appointed to administer the Protectorate. William Thomas was appointed Assistant Protector for the Port Phillip and Westernport Districts and had extensive contact with traditional *Bun wurrung* people during the early years of European settlement (Cotter 2005; 1-2). Thomas travelled with groups of *Bun wurrung* and related Woiworung people on seasonal movements around the Mornington Peninsula and Westernport, producing a map in 1841, showing the locations of *Bun wurrung* campsites and routes of movement.

Thomas also noted the frequent exploitation of the Nepean Peninsula coast. Sullivan ascertains that some of the places mentioned by Thomas could be base camps including *Boniong* and *Turtguruk* located in the hinterland of the Nepean Peninsula (1981, p.29).

Shellfish gathering was observed by Thomas in Port Phillip Bay, near Melbourne, when he reported that a group of women went at least three times a week to collect shellfish (Thomas cited in (Sullivan 1981, p.25). Cockle and mussel shells were observed in Aboriginal huts on the Nepean Peninsula (Knopwood cited in (Sullivan 1981, p.36). Rough waves pound the Bass Strait coastline of the Nepean Peninsula, particularly during winter, and this may have made the collection of shellfish difficult at this time of year and restricted activities to the warmer months (Sullivan 1981, p.8). Thomas also observed shellfish being collected by women diving in creeks and rivers (Thomas cited in (Sullivan 1981, p.28).

In 1840, Thomas (cited in (Sullivan 1981, p.30) observed that about 44 people were still exploiting the country between Mount Martha and Cape Schanck, divided into small groups of about six to seven people. Larger groups of 20 to 30 people would have gathered where resources were plentiful or concentrated (Sullivan 1981, p.32).

Thomas considered that the *Bun wurrung* economy was predominantly terrestrial in orientation despite their close association with the coast (Gaughwin & Sullivan 1984, p.89). However, it is possible that this was partially a reflection of the time of year that he made his observations. He noted that favourite foods were kangaroo and possum, and that they had the 'greatest abhorrence' for snakes (Thomas cited in (Sullivan 1981, p.22). Robinson was told by an eyewitness that in about 1834, "there was a tribe of Natives on the Point hunting kangaroo" (Robinson cited in (D'Arcy 2005, p.28). Women caught many smaller creatures such as bandicoots, rats and lizards (Thomas cited in (Sullivan 1981, p.22). It is probable that many women's subsistence and other activities were not seen or recorded by Thomas, either through his lack of interest or because the women carried them out away from the presence of men.

Fires were commonly lit by Aboriginal people in the coastal area of Port Phillip and were seen by early explorers (Sullivan 1981, p.23). While in the vicinity of Arthurs Seat in January 1803, James Flemming noted that the area was 'all newly burnt' (Shillinglaw 1972, p.24). In December 1836, George Augustus Robinson also noted at Point Nepean that "the bush was on fire, as indeed it was in all parts of the country" (Plomley 1987, p.405). Fires were mainly lit to maintain pathways through dense scrub, to increase the fertility of the land, to drive game and quite probably as a smoke screen to hide behind or to warn off or confuse intruders.

Further historical accounts of *Bun wurrung* traditions were recorded by Georgiana McCrae. Andrew McCrae and his wife Georgiana built a homestead at Arthur's Seat. Georgiana recounts that *Bun wurrung* people camped on the "edge of Cape Schanck Road" (McCrae 1934, p.209) and also mentions that large groups of *Bun wurrung* people made seasonal camps between the Cape Schanck Road and the bay beneath tea trees or coastal banksia (McCrae 1934, p.194). In 1851, Georgiana observed a *Bun wurrung* burial outside the paddock fence on the edge of the Cape Schanck Road:

I watched the grave being dug by some, while others wrapped a possum-rug about the corpse, which they interred in a sitting position, the elbows on the knees, the chin supported in the left hand, and the opposite one laid, with the fingers open, along the angle of the jaw. Cords were drawn tightly across the shoulders and around the waist, then a new pannikin and the last of the bottle of medicine I had sent him having been put into the grave, the father, and (fifth) stepmother filled the hole with sand. (McCrae 1992, p.213)

Indigenous travel routes were located throughout Bunurong country. Thomas' 1840 map below (Fig. 1) shows some of these tracks, it also shows Carrum Swamp, labelled 'lagoon'.




Figure 1. 1840 map by William Thomas indicating routes taken on his travels with *Bun wurrung* tribes. (Reproduced from Presland 1994, p. 74)

There is considerable emphasis in the ethnographic literature on the use of creeks and waterways as corridors of movement. Gunson (1974: 10) stated that members of the *Mayune balug* usually camped beside waterholes, creeks, and at coastal locations. Early settlers of the Western Port region also noted

that Aboriginal campsites containing huts were often found beside rivers and creeks (Sullivan 1981: 33). Carrum Swamp would have provided important resources and been a regular stopping place.

Just prior to, and overlapping, the period of British exploration and settlement, the *Bun wurrung* were involved in a long-running dispute with the *Gunai/Kurnai* people from Gippsland. According to William Barak, the conflict was a dispute over resources, which resulted in heavy casualties being suffered by the *Bun wurrung*. Many *Gunai/Kurnai* raids occurred to abduct Bunurong women. According to Barwick (1984), the *Yowengerra* had almost been completely annihilated by 1836, largely as a result of attacks from the *Gunai/Kurnai*. During 1833 - 1834 around 60-70 Bunurong people were killed in a raid by *Gunai/Kurnai* while they were camped to the north of Carrum Swamp (Rhodes 2003).

Warfare, disease and kidnapping all conspired to precipitate a rapid decline in the population of *Bun wurrung* peoples inhabiting their traditional lands. These numerous impacts would also have disrupted the economic and social ties binding the *Bun wurrung* to neighbouring *Kulin* peoples. Consequently, much of the *Bun wurrung* population had been displaced by the time of permanent European settlement of the area. By 1839 the *Bun wurrung* had been reduced to 83 people, with only 4 of 19 children under four years old, from a probable pre-contact population of greater than 300 people. By 1850 Protector William Thomas estimated just 28 *Bun wurrung* people remained (Rhodes & Bell 2004).. A camping reserve was established in 1841 at Mordialloc Creek where the *Bun wurrung* peoples were allocated 340 hectares. The rations for the reserve were managed by the Honorary Correspondents to the Boards of the Protection of Aborigines. The majority of the *Bun wurrung* population had moved to the Mordialloc reserve by 1856 which ceased operation in 1878.

Today the *Bun wurrung* people are represented by the Bunurong Land Council Aboriginal Corporation, Bunurong Land and Sea Association and the Boon Wurrung Foundation Ltd, who regard themselves as the traditional owners and custodians of this area.

4.2 Land-Use History of the Study Area

The study area is located within the margins of the former Carrum Swamp. The Carrum Swamp was an extensive fresh water wetlands area stretching from Mordialloc to Frankston. Drainage of the Carrum Swamp commenced in 1876 with the construction of Patterson Cut, now known as the Patterson River. In 1879 public works were conducted to create efficient drainage of the swamp to open up the land for settlement. By the early twentieth-century most of the original wetlands area had been drained and land was being used for market gardens and dairy farming. By the 1960s the former swamp area had been largely urbanised.




Figure 2: Map showing location of Edithvale-Seaford Wetlands in blue and extent of the former Carrum Swamp in grey.

Prior to European contact the Edithvale wetlands would have been one of the deeper parts of Carrum Swamp. Contour mapping shows that this area represents the lowest point in the surrounding landscape. An 1871 parish map shows the location of the swamp and the plan for division of land following drainage of the swamp.




Figure 3: 1871 Parish Map of Carrum Swamp (source: TROVE)

After Carrum Swamp was drained in the 1870s, land was opened up for farming, however it is likely that this area would have regularly been waterlogged and occasionally inundated when heavy rains occurred as contour mapping shows the area as being between 0-1m above sea level.

Historical aerial photography from 1945 shows the study area as cleared open farmland with minimal vegetation (Figure 4). In 1956 the study area has been partially inundated with pools of water forming in both the north and south sections of the wetlands. Residential subdivision of the Edithvale area has also developed rapidly since 1945 with large residential estates having been constructed to the west of the wetlands (Figure 5).

By 1963, some vegetation is appearing around the southwestern edge of the inundated sections of the study area (Figure 6). This vegetation has increased in density by 1972, and residential development is

present to the south of the study area (Figure 7). After 1980 (Figure 8) significant modifications have occurred to the northern section of the wetlands area, multiple pools have been constructed in the northern section along with a series of drains and weirs to regulate water flow and manage storm water. Along with these modifications, a series of trails have been constructed throughout the wetlands, along with birdwatching hides and a visitor centre (Figure 9).




Figure 4: 1945 map showing location of study area and surrounds



Figure 5: 1956 map showing location of study area and surrounds



Figure 6: 1963 map showing location of study area and surrounds




Figure 7: 1972 map showing location of study area and surrounds




Figure 8: 1980 map showing location of study area and surrounds



Figure 9: 2009 map showing location of study area and surrounds

5 Conclusions and Recommendations

5.1 Site Prediction Model

This desktop cultural heritage assessment for the study area has allowed a site prediction model to be developed. A site prediction model is intended for use as an indication of the types of archaeological sites that may occur in a given area.

The following statements can be made about the current study area:

- The location of previously registered Aboriginal Places within a 2km radius shows that Aboriginal sites can be located on elevated landforms around the edges of low-lying swampy areas, sites can also be located on sandy dune systems;
- Aboriginal site types are primarily low density artefact scatters and isolated artefacts, representing transitory use of the area rather than large camp sites;
- Raw materials are primarily quartz and quartzite, however some silcrete artefacts have also been located;
- While stone artefacts have been located underneath swamp deposits nearby, these were located within a remnant sand dune at a higher elevation than the study area;
- The study area is located in one of the deepest parts of the former Carrum Swamp and as such is unlikely to have been utilised by Aboriginal people in a manner that will leave cultural heritage material remains. While there is strong evidence pointing to the importance of Carrum Swamp as a resource for Aboriginal people, Aboriginal activity would have been focused around the margins and on elevated rises. As such, the study area is considered to contain extremely low archaeological potential; and
- In the post-contact period, the study area has been significantly modified through the drainage of Carrum Swamp, the clearance of native vegetation for agricultural purposes and use of the land for drainage and storm water management through the creation of a series of ponds, weirs and drainage channels. The majority of this modification has occurred in the northern section. The study area is also currently utilised for recreational purposes such as dog walking and bird watching and has been impacted by the construction of recreational structures such as walking trails, a bird hide and visitor centre.

The site prediction model concludes that there is an extremely low likelihood that Aboriginal cultural material will be present in the northern section of the study area, and a low likelihood in the southern section. If Aboriginal cultural heritage material is present it will most likely be in the form of low density stone artefact deposits.

5.2. Recommendations and Obligations under the Aboriginal Heritage Act

The Aboriginal Heritage Regulations 2007 state that:

A cultural heritage management plan is required for an activity if—

(a) all or part of the activity area for the activity is an area of cultural heritage sensitivity; and

(b) all or part of the activity is a high impact activity. (r. 6)

The study area is considered an area of cultural heritage sensitivity because it is a declared Ramsar wetland (Aboriginal Heritage Regulations (r. 26):

Declared Ramsar wetlands

(1) Subject to subregulation (2), a declared Ramsar wetland or land within 200 metres of a declared Ramsar wetland is an area of cultural heritage sensitivity.

(2) If part of a declared Ramsar wetland or part of the land within 200 metres of a declared Ramsar wetland has been subject to significant ground disturbance, that part is not an area of cultural heritage sensitivity.

For this reason if high impact activities are planned within the study area they would trigger a mandatory CHMP. A list of high impact activities is presented within Division 5 of the *Aboriginal Heritage Regulations 2007* (Appendix 1).

However, as regulation 26 (2) states, if the area has been subject to significant ground disturbance, that area is not an area of cultural heritage sensitivity and as such a mandatory CHMP would not be required. Some sections of the study area have clearly been subject to significant ground disturbance, for example, areas where water retaining ponds have been excavated. The Office of Aboriginal Affairs Victoria (OAAV) Practice Note on determining significant ground disturbance is attached as Appendix 3.

For this reason each project planned for the study area would need to be evaluated on an individual basis to determine if a CHMP would be required. The project manager would need to determine if the project is considered a high impact activity under the *Aboriginal Heritage Regulations 2007*, and if this is the case whether the location of the activity is in an area that could be considered to have undergone significant ground disturbance as defined in the *Regulations*. It should be noted that historical evidence shows that the study area has been substantially modified in the northern section, however, less disturbance is apparent in the southern section.

A field survey of the study area would be able to provide a more detailed archaeological assessment and identify areas of specific archaeological potential along with any surface Aboriginal sites within the study area. This information could then be used to determine whether mandatory or voluntary CHMP preparation is required for specific proposed works.

If there are any questions about the contents of this report or about the management of potential heritage issues within the wetlands, please don't hesitate to contact Heritage Insight P/L on 9376 6569.

References

Legislation

Aboriginal Heritage Act 2006

Aboriginal Heritage Regulations 2007

Organisations

Land Victoria Aerial Photography storage facility (Laverton)

Public Record Office Victoria

Internet Resources

Department of Environment and Primary Industries – Biodiversity Interactive Map
(<http://www.depi.vic.gov.au/environment-and-wildlife/biodiversity/biodiversity-interactive-map>)

Department of Environment and Primary Industries – EVC Benchmarks
(<http://www.depi.vic.gov.au/environment-and-wildlife/biodiversity/evc-benchmarks>)

Department of Environment and Primary Industries – GeoVic
(http://er-info.dpi.vic.gov.au/sd_weave/registered.htm)

National Library of Australia (TROVE)
(<http://trove.nla.gov.au/>)

Reports and Published Resources

Barwick, D 1984, 'Mapping the Past: an Atlas of Victorian Clans 1835-1904. Part 1', *Aboriginal History*, vol. 8, no. 1-2, pp. 100–131.

Bride, TF 1969, *Letters from Victorian Pioneers*, Robert Brain, Melbourne.

Cannon, M (ed.) 1983, *Historical Records of Victoria: Aborigines and Protectors 1838-1839*, Victorian Government Printing, Melbourne.

Clark, I 1990, *Aboriginal Languages and Clans: An Historical Atlas of Western and Central Victoria, 1800-1900*, Department of Geography and Environmental Science Monash University, Melbourne.

Coutts, PJF 1977, *Aboriginal Prehistory in North Western Victoria*, Victoria Archaeological Survey, Melbourne, Vic.

D'Arcy, J 2005, *Bunurong Land Council Historical and Genealogical Report*, Bunurong Land Council, Aboriginal Corporation.

DEPI 2013, *Edithvale-Seaford Wetlands Ramsar Site Boundary Description: Technical Report*, Department of Environment and Primary Industries, East Melbourne.

Dugay-Grist, L, Cowled, A & Maher, M 2012, *58 Fourth Avenue, Chelsea Heights: Voluntary CHMP*, Grist Archaeology Heritage Management.

- Gaughwin, D & Sullivan, H 1984, 'Aboriginal Boundaries and Movements in Western Port, Victoria', *Aboriginal History*, vol. 8, no. 1, pp. 80–98.
- Green, M 2013, *Springvale Road, Aspendale Gardens Nursery Development.*, Andrew Long & Associates, Fitzroy.
- KBR 2009, *Edithvale-Seaford Ramsar Site: Management Plan*, Kellogg Brown & Root Pty Ltd, St Kilda.
- Kennedy, S & Crocker, S 2013, *Perry Road and Mordialloc Main Drain: Drainage Infrastructure*, Keysborough, Archaeological & Heritage Management Solutions P/L (AHMS), Brunswick, Vic.
- Kennedy, S, Wheeler, J & Foley, E 2012, *44 First Avenue, Chelsea Heights*, Australian Cultural Heritage Management (ACHM), Brunswick, Vic.
- Kiddel, H, Watson, B & Howes, J 2011, *Residential Development 8-10 Maury Road, Chelsea*, Dr. Vincent Clark & Associates, Coburg & Woodend.
- Matic, A 2014, *40 Ella Grove, Chelsea: Residential Subdivision*, Pragmatic Cultural Heritage Services, Belgrave.
- Matic, A & Falvey, L 2012, *Sporting Pavilion and Early Years Hub, Edithvale*, Terra Culture Heritage Consultants Pty Ltd, Northcote.
- Matic, A, MacCulloch, J & Falvey, L 2013, *Drainage Works, Edithvale*, Terra Culture Heritage Consultants Pty Ltd, Northcote.
- McCrae, H (ed.) 1934, *Georgiana's Journal: Melbourne A Hundred Years Ago*, Angus & Robertson, Sydney.
- McCrae, H (ed.) 1992, *Georgiana's Journal Melbourne 1841-1865*, Angus & Robertson, Sydney.
- Mitchell, J 2014, *Proposed Residential Development at 58-60 Berry Avenue, Edithvale*, Alpha Archaeology Pty Ltd, Kensington.
- Murphy, A & Dugay-Grist, L 2008, *Edithvale-Seaford Wetland Discovery Centre. Cultural Heritage Management Plan 10081*, Tardis Enterprises Pty Ltd, Beaconsfield Upper.
- Murphy, A & Thomson, S 2013, *Residential Development, 85 Berry St Edithvale*, Tardis Enterprises Pty Ltd, Beaconsfield Upper.
- Murphy, A & Thomson, S 2015, *Subdivision Lot 1, 2, 3, TP49429 & Lot A PS 605114 Springvale Road Aspendale Gardens*, Archaeology at Tardis Pty Ltd, Beaconsfield.
- Plomley, NJB 1987, *Weep In Silence*, Blubber Head Press, Hobart.
- Presland, G (ed.) 1977, *Journals of George Augustus Robinson January - March 1840* Second Edition., Ministry for Conservation Publication, Melbourne.
- Rhodes, D 2003, *Victorian Channel Deepening Project EES: Aboriginal Cultural Heritage. Existing Conditions Report*, Terra Culture Heritage Consultants Pty Ltd, Fairfield.
- Rhodes, D & Bell, J 2004, *Shire of Cardinia Urban Growth Corridor Aboriginal Heritage Study*, Heritage Insight Pty Ltd, Richmond.
- Rogers, H 1957, *The Early History of the Mornington Peninsula*, Hallcraft Publishing Co. Pty. Ltd, Melbourne.
- Schultz, F & Donati, L 2011, *Proposed Residential Subdivision at 88 Berry Ave, Edithvale*, Australian Cultural Heritage Management (ACHM), Melbourne.

- Shillinglaw, J (ed.) 1972, *Historical records of Port Phillip, the first annals of the colony of Victoria* Facsimile edition of the 1879 edition., Heinemann Publishers Australia.
- Smith, LJ 1991, *Berwick - Pakenham Corridor Aboriginal Archaeology*, Victorian Archaeological Survey, Melbourne.
- Sullivan, H 1981, *Victorian Occasional Reports Series Number 6: An Archaeological Survey of the Mornington Peninsula, Victoria*, Victorian Archaeological Survey, Melbourne.
- Thomas, S & Howell-Meurs, J 2012, *142 - 143 Nepean Highway, Aspendale*, Andrew Long & Associates, Fitzroy.
- Webb, C 2000, *An Archaeological Assessment of a proposed subdivision at the Patterson River Country Club.*, Terra Culture Pty Ltd, Coburg.
- White, T 2015, *Residential Subdivision 1 Bristol Avenue, Edithvale 3196*, T.L. White Heritage Consulting Services, Berwick.

Appendix 1: Division 5 of the Aboriginal Heritage Regulations 2007

Division 5—High impact activities

42 Purpose

The purpose of this Division is to specify high impact activities.

Note

Under regulation 6, a cultural heritage management plan is required for an activity if all or part of the activity area is an area of cultural heritage sensitivity and if all or part of the activity is a high impact activity.

43 Buildings and works for specified uses

- (1) The construction of a building or the construction or carrying out of works on land is a high impact activity if the construction of the building or the construction or carrying out of the works—
 - (a) would result in significant ground disturbance; and
 - (b) is for or associated with the use of the land for any one or more of the following purposes—
 - (i) aquaculture;
 - (ii) a camping and caravan park;
 - (iii) a car park;
 - (iv) a cemetery;
 - (v) a child care centre;
 - (vi) a corrective institution;
 - (vii) a crematorium;
 - (viii) an education centre;
 - (ix) an emergency services facility;
 - (x) a freeway service centre;
 - (xi) a hospital;
 - (xii) an industry;
 - (xiii) intensive animal husbandry;
 - (xiv) a major sports and recreation facility;
 - (xv) a minor sports and recreation facility;
 - (xvi) a motor racing track;
 - (xvii) an office;
 - (xviii) a place of assembly;

- (xviii) a pleasure boat facility;
 - (xix) a research centre;
 - (xx) a retail premises;
 - (xxa) a retirement village;
 - (xxi) a service station;
 - (xxii) a transport terminal;
 - (xxiii) a utility installation, other than a telecommunications facility, if—
 - (A) the works are a linear project that is the construction of an overhead power line with a length exceeding one kilometre or for which more than 10 power poles are erected; or
 - (B) the works are a linear project that is the construction of a pipeline with a length exceeding 500 metres; or
 - (C) the works are a linear project with a length exceeding 100 metres (other than the construction of an overhead power line or a pipeline with a pipe diameter not exceeding 150 millimetres); or
 - (D) the works affect an area exceeding 25 square metres.;
 - (xxiv) a veterinary centre;
 - (xxv) a warehouse;
 - (xxvi) land used to generate electricity, including a wind energy facility.
- (2) The terms used in subregulation (1)(b) have the same meanings as they have in the VPP.
- (3) Despite subregulation (1), the construction of a building or the construction or carrying out of works on land is not a high impact activity if it is for or associated with a purpose for which the land was being lawfully used immediately before the commencement day.
- (4) In this regulation, ***linear project*** has the same meaning as in regulation 68.

44 Constructing specified items of infrastructure

- (1) The construction of any one or more of the following is a high impact activity if the construction would result in significant ground disturbance—
- (a) an airfield;
 - (b) a bicycle track with a length exceeding 100 metres;
 - (c) a helipad;
 - (d) rail infrastructure, other than—
 - (i) a railway track with a length of less than 100 metres; or
 - (ii) a railway track siding with a length of less than 100 metres; or
 - (iii) a cutting with a length of less than 100 metres; or
 - (iv) a tunnel with a length of less than 100 metres; or
 - (v) a bridge with a span of less than 100 metres; or

- (vi) a platform with a length of less than 100 metres; or
 - (vii) a service road with a length of less than 100 metres;"
 - (e) a road with a length exceeding 100 metres;
 - (f) a walking track with a length exceeding 100 metres;
 - (g) a telecommunications line consisting of an underground cable or duct with a length exceeding 500 metres.
- (2) In this regulation, **telecommunications line** has the same meaning as in the VPP.

45 Dwellings

- (1) The construction of three or more dwellings on a lot or allotment is a high impact activity.
- (2) The carrying out of works for three or more dwellings on a lot or allotment is a high impact activity.

Example

Constructing an apartment tower containing 50 dwellings is a high impact activity.
Constructing or extending only one or two dwellings on a lot or allotment is not a high impact activity.

Note

See regulation 9 in relation to the construction of a building, or the construction or carrying out of works, where the building or works are ancillary to an existing dwelling or the construction of one or two dwellings on a lot or allotment.

- (3) This regulation does not apply to the construction of or the carrying out of works for a retirement village within the meaning of the VPP.

46 Subdivision of land

- (1) The subdivision of land into three or more lots is a high impact activity if—
 - (a) the planning scheme that applies to the activity area in which the land to be subdivided is located provides that at least three of the lots may be used for a dwelling or may be used for a dwelling subject to the grant of a permit; and
 - (b) the area of each of at least three of the lots is less than eight hectares.
- (2) The subdivision of land into two or more lots in an industrial zone is a high impact activity.
- (3) In this regulation, **industrial zone** has the same meaning as in the VPP.

47 Alpine resorts

- (1) The construction of a building or the construction or carrying out of works in an alpine resort is a high impact activity if the construction of the building or the construction or carrying out of the works would result in significant ground disturbance.
- (2) In this regulation, **alpine resort** has the same meaning as in the **Alpine Resorts Act 1983**.

48 Activities requiring earth resource authorisations

An activity is a high impact activity if it is an activity—

- (a) for which an earth resource authorisation is required before the activity may be carried out; and
- (b) that would result in significant ground disturbance.

48A Extraction or removal of stone

- (1) The extraction or removal of stone (other than sand or sandstone) that does not require an earth resource authorisation is a high impact activity if—
 - (a) the primary purpose of the extraction or removal is—
 - (i) the sale or commercial use of the stone; or
 - (ii) the use of the stone in construction, building, road or manufacturing works; and
 - (b) the land from which the stone is extracted or removed is more than 2000 square metres; and
 - (c) the extraction or removal would result in significant ground disturbance.
- (2) In this regulation, **stone** has the same meaning as in the **Mineral Resources (Sustainable Development) Act 1990**.

49 Extraction or removal of sand or sandstone

- (1) The extraction or removal of sand or sandstone (other than extraction or removal that requires an earth resource authorisation) is a high impact activity if the extraction or removal would result in significant ground disturbance.
- (2) Subregulation (1) does not apply to the extraction or removal of sand or sandstone—
 - (a) from land that is a farm if the sand or sandstone is intended in good faith only to be used on that farm for the purposes of a dam or other farmworks and not for sale or any other commercial use; or
 - (b) undertaken by or on behalf of a Minister responsible for the administration of the **Conservation, Forests and Lands Act 1987** where the primary purpose of the extraction is for the footings or foundations of a building or structure, the construction of a carpark, road, track or other works or for any borrow pit adjacent to such an excavation; or
 - (c) if the extraction or removal, including dredging, constitutes works for marine navigational purposes or the establishment or renourishment of a beach; or
 - (d) if the extraction or removal constitutes works for the purpose of establishing a port facility, railway or tunnel; or
 - (e) if the primary purpose of the excavation or removal is for the construction of the footings or foundations of a building or structure.

50 Searching for stone

- (1) A search for stone is a high impact activity if it would result in significant ground disturbance.
- (2) In this regulation, *search for stone* has the same meaning as in the VPP.

51 Extraction or removal of loose stone on agricultural land on the Victorian Volcanic Plain

- (1) The extraction or removal of loose stone from the surface of land used for agriculture on the Victorian Volcanic Plain is a high impact activity if the extraction or removal—
 - (a) is for the primary purpose of land improvement, including pasture enhancement; and
 - (b) would result in significant ground disturbance.
- (1A) The crushing of loose stone on the surface of land used for agriculture on the Victorian Volcanic Plain is a high impact activity if the crushing is—
 - (a) by machinery; and
 - (b) for the primary purpose of land improvement, including pasture enhancement."
- (2) In regulation 51(2) of the Aboriginal Heritage Regulations 2007, for "Subregulation (1) does" **substitute** "Subregulations (1) and (1A) do".
- (3) In regulation 51(3) of the Aboriginal Heritage Regulations 2007, for "**Extractive Industries Development Act 1995;**" **substitute** "**Mineral Resources (Sustainable Development) Act 1990;**
- (2) Subregulation (1) does not apply if the land is used for crop raising or has been used for crop raising.
- (3) In this regulation—

agriculture and *crop raising* have the same meanings respectively as they have in the VPP;

stone has the same meaning as in the **Extractive Industries Development Act 1995;**

Victorian Volcanic Plain means the area comprised of the areas identified as "Qvh", "Qvn", "Qvs", "Qvs2" and "Qvt" on the following Geological Survey of Victoria 1:250 000 map series sheets—

 - (a) SJ54-8 entitled "Ballarat" (second edition, 1997);
 - (b) SJ55-1 entitled "Bendigo" (third edition, 2001);
 - (c) SJ54-12 entitled "Colac" (second edition, 1997);
 - (d) SJ54-7 entitled "Hamilton" (second edition, 1997);
 - (e) SJ55-5 entitled "Melbourne" (second edition, 1997);
 - (f) SJ54-11 entitled "Portland" (second edition, 1997);

(g) SJ55-9 entitled "Queenscliff" (second edition, 1997);

(h) SJ54-4 entitled "St Arnaud" (second edition, 1997).

52 Timber production

- (1) The use of an area of land greater than 40 hectares in size for timber production is a high impact activity if—
- (a) a permit is required under a planning scheme to use the land for timber production; and
 - (b) the use of the land for timber production would result in significant ground disturbance.
- (2) The construction of a building associated with timber production is a high impact activity if—
- (a) a permit is required under a planning scheme to construct the building; and
 - (b) the construction of the building would result in significant ground disturbance.

r. 53

- (3) In this regulation, *timber production* has the same meaning as in the VPP.

Note

A permit may not be required under a planning scheme to use an activity area for timber production if the timber production is ancillary to a particular agricultural enterprise (such as agroforestry).

53 Dams

The construction or alteration of a private dam, other than on a waterway, is a high impact activity if a licence is required under section 67(1A) of the **Water Act 1989** for the construction or alteration of the private dam.

54 Use of land

- (1) The use of land for a purpose specified in regulation 43(1) is a high impact activity if a statutory authorisation is required to use the land for that purpose.
- (2) The use of land for an extractive industry is a high impact activity if a statutory authorisation is required to use the land for the extractive industry.
- (3) The use of a lot or allotment for three or more dwellings is a high impact activity if a statutory authorisation is required to use the lot or allotment for three or more dwellings.
- (4) Despite subregulations (1), (2) and (3), if the whole of the activity area for an activity referred to in subregulation (1), (2) or (3) has been subject to significant ground disturbance, that activity is not a high impact activity.
- (5) In this regulation, *extractive industry* has the same meaning as in the VPP.

Example

A land owner proposes to change the use of his or her land from the grazing of animals to the storage of shipping containers. The land is flat and, in the first instance, no works are proposed, although the grass will first be cut and some non-indigenous shrubs removed. The use of the land for storing shipping containers is an industry and requires a statutory authorisation (a permit under the relevant planning scheme). The proposed use is a high impact activity. If, at a later date, the area is upgraded by works, including excavation for a concrete base on which to store the containers, the upgrade works would also be a high impact activity under regulation 43(1).

Appendix 2: Glossary

Adze A flake with stepped retouch along lateral margins that can be hafted for use as a tool.

Anvil A flat object on which a core was placed to flake material from. Anvils often have a small pit/groove, usually in the centre of the object, as a result of this action.

Archaeology The study of cultural remains from past cultures and generations.

Artefact Scatter The material remains of past Aboriginal peoples' activities. Usually contain stone artefacts, but other material may also be present, including charcoal, animal bone, shell and ochre. An artefact scatter is usually represented by a single stone flake or a concentration of flaked stone pieces (or fragments).

Assemblage A collection of artefacts that are derived from the same site.

Backed Blade Stone artefact associated with the Australian small tool tradition. They are characterised by unidirectional or bidirectional retouch found along a lateral margin, thought to be blunt for hafting (Holdaway & Stern 2004: 260).

Basalt A fine-grained rock occurring from lava flows.

Bifacially Flaked Flakes removed from two faces of an object such as a core.

Blade A flake that is twice as long as it is wide.

Bondi Point An asymmetrical blade with a point at one end with backing retouch. Part of the Australian Small Tool Tradition.

Burial Human Remains, normally found as concentrations of human bones or teeth, exposed by erosion or earthworks. They are sometimes associated with charcoal or ochre, although shell, animal bone and stone tools may also be present. Tend to be located in soft soils and sand, although can occur in rock shelters, caves and dead trees.

Burin A truncated flake formed by snapping or retouching along one lateral margin that then forms a platform from which small flakes are removed forming a triangular scar that acts as a working edge (Holdaway & Stern 2004: 241-243).

Ceramic A term used to identify wares made from either clay or fusible stone such as stoneware, earthenware, porcelain or terracotta (Davies & Buckley 1987: 186).

Chert A compact, fine-grained rock made of cryptocrystalline silica and can occur in a variety of colours, usually red, green or black.

Core A specimen of rock that has undergone a process of reduction through the removal of a number of flakes and as a result they have negative flake scars. Cores can contain a single platform, have two platforms or have had flakes removed in multiple directions.

Cortex The original surface of a mineral or rock subjected to weathering by the elements.

Cultural Material Any material remains which are produced by human activity.

Debitage Detached pieces of stone that are discarded during the reduction process.

Dry Stone Wall A wall formed of a number of courses of rock (usually basalt or limestone) with no bond or binding component. Walls are usually tapered, have two faces and can have hearting (packing), or plugging.

Earthenware A non-vitreous (porous) whiteware, usually used for domestic tablewares. Most earthenware is glazed and decorated, transfer printed or left plain (Davies & Buckley 1987: 186).

Earth Feature Collective term used to refer to mounds, rings, hearths, postholes and ovens.

Earth Mound Mounds generally appear as raised areas of darker soil. They are commonly found in the volcanic plains of western Victoria or on higher ground near water bodies. Mounds often contain charcoal, burnt clay or stone heat retainers from cooking ovens, animal bones, shells, stone tools and sometimes, Aboriginal burials.

Earth Ring Banked circles of soil often associated with stone arrangements, which had a ceremonial purpose for Aboriginal people in the past.

Excavation A controlled means of soil disturbance (digging) allowing for detailed recording of the soil profile, features and artefacts exposed.

Flake A stone artefact that contains characteristics such as the presence of a platform, bulb of percussion and termination which reveal that the stone has been struck from a core and is the result of stone working (Holdaway & Stern 2004: 5).

Flake Core A flake that has subsequently been used as a core and had other flakes removed from it.

Flaked Piece Small fragments of stone that have been removed from flakes resulting from tool maintenance or tool production (Holdaway & Stern 2004: 17). Flaked pieces do not display the characteristics evident in a complete flake.

Flint Similar to chert with a pale cortex and conchoidal fracture. Usually occurring in limestone (Roberts 1998: 65).

Footing The structural base/footprint from structures often built from bluestone, brick or wooden posts.

Geometric Microlith Part of the Australian small tool tradition. They are symmetrical in form, pointed at both ends and can be backed along a lateral margin (Holdaway & Stern 2004: 262).

Glaze A coating put over wares fired in a kiln. Glazes can come in a variety of colours and can also be transparent.

Greenstone A metamorphic rock derived from basalt containing feldspar and quartz and is made green by chlorite and epidote. Often used for the manufacture of hand axes.

Grindstone A flat slab of rock with central depression used to grind, crush or pound seeds, ochre, or sharpen tools, etc. Grindstones are usually made on sedimentary rocks with an abrasive surface and can be used in conjunction with a muller.

Ground Edge Axes A sharpening process – flaking, pecking and polishing, usually along a single lateral margin. The axes are generally hafted with the worked edge forming the tool edge.

Ground Surface Visibility The extent to which the natural soil surface below the vegetation on the ground is visible.

Hammerstone A hard rock or mineral used to flake fragments of stone from a core (Holdaway & Stern 2004: 4).

Hearth The remains of a fireplace containing charcoal and sometimes burnt earth, bone, stone artefacts or other organic material.

In situ An artefact or feature that remains in its original position, or where it was left.

Manuport A stone block that displays no attributes of being either a core or a flake.

Microblade Has the same characteristics as a blade but just of smaller proportions (Holdaway & Stern 2004: 17).

Ochre Earth varying in colour from yellow to red, used as a pigment.

Organic Compounds formed from living organisms (plants or animals).

Oven Mound Usually circular or oval in shape and often situated close to a water source. They were used for cooking and contain a rich greasy organic mix of soil and organic material. An oven mound is likely to contain charcoal, burnt clay or stone heat retainers, stone tools, bones, shell and on occasion, burials (AAV Mini Poster 4).

Platform The surface from which the flake was struck off the core – natural, flaked or abraded (Holdaway & Stern 2004: 120).

Point A flake that has two edges that form a point with retouch along one or both lateral margins (Holdaway & Stern 2004: 16).

Porcelain A non-porous ceramic with a glass-like appearance. Can be translucent, can be used for tableware or more decorative features such as ornaments.

Post-contact The period after contact between Aboriginal people and Europeans.

Pre-contact The period before contact between Aboriginal people and Europeans.

Quarry Outcrop of stone or ochre that has been quarried by Aboriginal people in the past. Generally

associated with a large amount of broken stone and flakes. The outcrop (cores) bear negative scars from flaking.

Quartz A mineral that commonly occurs in sedimentary, igneous and metamorphic rocks. Quartz can come in a number of forms including crystal, rose, and smoky.

Quartzite A metamorphic rock formed by the recrystallization of quartz. Quartz is rich in sandstone and limestone (Roberts 1998: 109).

Retouch A worked edge or modification of a flake formed by removing a number of small flakes along an edge. This can be done as a form of maintenance or to produce a tool.

Rock Art Paintings created on the rock surfaces of caves and rock shelters and engravings in limestone caves. Artwork includes stencils, prints and drawings. The paint consists of ochres, clays and charcoal mixed with fats.

Scarred Tree A tree which has had a slab of bark removed, exposing the sapwood on the trunk or branch of a tree. Aboriginal people used the bark to make shelters, containers (coolamons) and canoes.

Scraper A flake with at least one edge that has continuous retouch. Scraper types include steep-edged, end, side and nose scraper (Holdaway & Stern 2004: 16).

Shell Midden A surface and/or sub-surface deposit composed of shell and sometimes stone artefacts, charcoal and bone. Middens are normally found in association with coastlines, rivers, creeks and swamps – wherever coastal, riverine or estuarine shellfish resources were available and exploited.

Silcrete A fine-grained rock derived from shale or siltstone mixed with silica.

Spit A horizontal unit of soil removed during excavation. Spits can be arbitrary (dug to a depth of 50, 100, 200, 300mm, etc.) or can be confined to a particular soil type or context. The excavation of spits allows for greater understanding, analysis and interpretation of the soil profile.

Stone Feature Includes cairns, rock wells, stone arrangements, fish traps, stone structures and grinding

grooves. May be a natural feature, which was used or modified to be used by Aboriginal people in the past (rock well, stone arrangement), or a stone feature which has been deliberately constructed for a specific purpose (fish trap, stone structure, cairn), or is the result of a specific activity carried out by Aboriginal people in the past (grinding grooves).

Stoneware A vitreous (non-porous) ceramic, usually light brown in colour, used for drinking containers or used industrially. Often glazed or unglazed (salt glaze or slip applied) (Davies & Buckley 1987: 186).

Stratification The position of sediments and rocks in sequence throughout time.

Sub-surface Testing A method of excavation that involves ground disturbing works to identify the potential for cultural material. Sub-surface testing may comprise hand excavation and/or machine excavation.

Survey An inspection of land either by foot or by car (windscreen survey) noting conditions on surface visibility, landforms and the presence of cultural material.

Termination The shape of the distal end of a flake (Holdaway & Stern 2004: 129).

Terracotta A low-fired clay (ceramic), usually orange to red in colour and very porous. Often used for plumbing (drainage components) or garden ware.

Tool Modified flakes usually with retouch present along an edge (Holdaway & Stern 2004: 33).

Transect An excavated stretch of ground that can be of varying lengths in a straight line.

Transfer Printed A design is traced and engraved onto a copper plate on which ink and oil is then applied. The design is pressed onto tissue paper and then placed on an object and the paper removed. The object is then fired and glazed. Transfer printed ceramics come in a variety of colours and patterns and were mass produced.

Trench An area confined by excavation usually in the form of a square (e.g., 2x2m) or rectangular (e.g., 1.5x1m).

References

AAV Mini Posters (1-7)

Davies, M. & K. Buckley, 1987 *Port Arthur Conservation & Development Project: Archaeological Procedures Manual*. Occasional Paper No.13. Department of Lands, Parks and Wildlife, Tasmania.

Holdaway S. & N. Stern, 2004 *A Record in Stone: The Study of Australia's Flaked Stone Artefacts*, Museum Victoria, Melbourne.

Roberts, J. L., 1998 *A Photographic Guide to Minerals, Rocks and Fossils*, New Holland, London.

Appendix 3: OAAV Practice Note for Determining Significant Ground Disturbance

Edithvale-Seaford Ramsar Wetlands
Seaford Section
Desktop Cultural Heritage Assessment



Cultural Heritage Advisor: Bianca Di Fazio

Authors: Renee McAlister & Bianca Di Fazio

June 6, 2016

**Heritage
Insight**
Pty Ltd



Heritage Insight P/L
53 Stubbs Street Kensington VIC 3031

PHONE: +61 3 9376 6569 INQUIRIES: archaeology@heritageinsight.com
FAX: +61 3 9376 6526 WEB: www.heritageinsight.com

ABN: 73 116 621 884 ACN: 116 621 88

Acknowledgements

Heritage Insight Pty Ltd would like to acknowledge and thank the following people for their assistance and participation in this study:

Ecology Australia

Darren Quin

Jonathan Ricciardello

Heritage Insight

Shannah Anderson Background Research

Katarina Audy GIS Mapping

Traditional Owner Groups

Wurundjeri Tribe Land Compensation and Cultural Heritage Council Inc (WTCCHC)

Boon Wurrung Foundation (BWF)

Bunurong Land Council Aboriginal Corporation (BLCAC)

Bunurong Land and Sea Association (BLSA)

Heritage
Insight
Pty Ltd



Heritage Insight P/L
53 Stubbs Street Kensington VIC 3031

PHONE: +61 3 9376 6569 INQUIRIES: archaeology@heritageinsight.com
FAX: +61 3 9376 6526 WEB: www.heritageinsight.com

ABN: 73 116 621 884 ACN: 116 621 88

Disclaimer

The information contained in this Cultural Heritage Assessment has been compiled from the standard heritage database sources and is accurate as far as Heritage Insight Pty Ltd is aware. However, within the timeframes available for technical heritage reporting, it is not possible to carry out comprehensive research of all published or unpublished manuscripts, journals, maps or oral history which may pertain to the study area. No responsibility can be taken for errors or omissions in primary and secondary source material cited in this report. Any opinions expressed in this report are those of Heritage Insight Pty Ltd only.

Copyright Notice

This report is copyright. Apart from any fair dealing for the purposes of private study, research, criticism or review as permitted under the Copyright Act, no part may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, whether electronic, mechanical, photocopying, recording or otherwise without prior written permission. Enquiries should be made to Heritage Insight Pty Ltd.

Executive Summary

This report outlines the results of an investigation into the heritage values of the Edithvale-Seafood Ramsar Wetlands, and has been conducted as part of the larger Ramsar Management Plan undertaken by Ecology Australia for Melbourne Water.

The purpose of this report is to identify the heritage values associated with the Seafood section of the Edithvale-Seafood Ramsar wetlands, commonly known as the Seafood Swamp, formulate a site prediction model and provide recommendations for the ongoing management of cultural heritage issues within the area. A second report has been produced which deals with the Edithvale area.

Section 2 outlines information about the formation and hydrology of the Seafood section of the Edithvale-Seafood Ramsar wetlands. It discusses the current use of the area, as well as the geology and geomorphology, and provides information on the climate and available flora, fauna and stone resources prior to European contact. Seafood Swamp currently comprises a number of separate water bodies, which serve different purposes, including both fresh water and saline water deposits. The Seafood Swamp is utilised for both drainage purposes, stormwater management and as an open public space. The wetlands are home to a significant population of birdlife.

Section 3 presents a review of previous archaeological assessment within the study area and surrounds, this information is combined with the environmental data and the land use history to assist in the formulation of a site prediction model. A review of the Victorian Aboriginal Heritage Register (VAHR) data tells has shown that there are two registered Aboriginal places within the study area: a shell midden and an artefact scatter. The presence of two Aboriginal sites within the study area also illustrates that despite the Seafood Swamp having previously been part of Carrum Swamp and frequently inundated, fluctuating water levels would have provided access both around and through the swamp. Registered Aboriginal places within the study area are evidence of previous Aboriginal activity within the area. While two Aboriginal sites have been located within the study area, the study area has never been subject to detailed archaeological survey.

Section 4 details the land use history of the study area and provides an ethnography for the *Bun wurrung* people. A review of the land use history demonstrated that the study area is a landform that has undergone substantial modification and disturbance over the years, from being part of Carrum Swamp, through drainage and utilisation for farming, to its role in managing drainage and storm water today.

The site prediction model is contained in Section 5, and has been developed by utilising all of the available data. This model notes that two Aboriginal sites have been located within the study area, a shell midden and an artefact scatter. Both these sites are recorded as surface sites and are potentially much larger than recorded at present. Geological and EVC mapping shows that the north western section and the eastern margin of the study area contain elevated sandy landforms and these areas should be considered to contain moderate to high potential for the discovery of Aboriginal cultural material. Low lying areas within the study area and areas that have undergone substantial ground disturbance can be considered to contain little to no archaeological potential.

Section 5.2 outlines the responsibilities and requirements for future development under the *Aboriginal Heritage Regulations 2007*. It is noted that if any project is planned that constitutes a high impact activity as defined in the *Regulations* (Appendix 1), then that project will require that a mandatory CHMP be prepared.

It is also recommended that the land manager may wish to undertake a more detailed archaeological assessment of the study area in the form of a field survey to;

- Further investigate the registered Aboriginal places within the study area;
- Identify any surface deposits of Aboriginal cultural heritage material;
- Identify areas of archaeological potential; and
- Further investigate the extent of ground disturbance within the study area.

These results could then be utilised to assist with the future management of heritage issues within the Seaford wetlands area. This would be particularly useful for any activities that do not trigger a mandatory CHMP and could be used to prevent accidental harm to unrecorded Aboriginal cultural heritage material, which is an offence under the *Aboriginal Heritage Act 2006*.

Table of Contents

<i>Executive Summary</i>	<i>i</i>
<i>Part One: Assessment</i>	<i>1</i>
<i>1.0 Introduction</i>	<i>1</i>
<i>2. Existing Conditions, Environment and Geology</i>	<i>4</i>
2.1 Existing Conditions	4
2.2 Landforms and Geomorphology of the Study Area.....	6
2.3 Resources Available to Aboriginal People within the Study Area.....	8
<i>3. Review of Previous Archaeological Research</i>	<i>11</i>
3.1 Victorian Aboriginal Heritage Register Search.....	11
3.2 Previous Archaeological Assessment within 2km of the Study Area	15
<i>4. Land Use History and Ethnography</i>	<i>27</i>
4.1 Historical and Ethno-Historical Accounts in the Geographic Region	27
4.2 Post-Contact Land Use History of the Study Area	30
<i>5 Conclusions and Recommendations</i>	<i>40</i>
5.1 Site Prediction Model	40
5.2. Recommendations and Obligations under the Aboriginal Heritage Act	42
<i>References</i>	<i>43</i>
<i>Appendix 1: Division 5 of the Aboriginal Heritage Regulations 2007</i>	<i>48</i>
<i>Appendix 2: Glossary</i>	<i>56</i>
<i>Appendix 3: OAAV Practice Note for Determining Significant Ground Disturbance</i>	<i>61</i>

Maps

Map 1: Location of the study area Note: Coordinates provided are in GDA94 MGA (Zone 55).....	3
Map 2: Map showing current conditions	5
Map 3: Map showing geology of the study area and surrounds	7
Map 4: Map showing 1750s EVC mapping of the study area and surrounds.....	9
Map 5: Map showing location of registered Aboriginal places in the study area and within 200m of the study area.....	14

Figures

Figure 1. 1840 map by William Thomas indicating routes taken on his travels with <i>Bun wurrung</i> tribes. (Reproduced from Presland 1994, p. 74)	29
Figure 2: Map showing location of Edithvale-Seafood Wetlands in blue and extent of the former Carrum Swamp in grey.	31
Figure 3: 1871 Parish Map of Carrum Swamp (source: TROVE)	32
Figure 4: 1945 map showing location of study area and surrounds	34
Figure 5: 1956 map showing location of study area and surrounds	35
Figure 6: 1968 map showing location of study area and surrounds	36
Figure 7: 1972 map showing location of study area and surrounds	37
Figure 8: 1989 map showing location of study area and surrounds	38
Figure 9: 2009 map showing location of study area and surrounds	39

Tables

Table 1: Summary of registered Aboriginal Places within the geographic region	12
Table 2: Summary details for Aboriginal Places within 2km of the activity area (sites within the study area shown in green, sites within 200m shown in blue).....	12
Table 3: Relevant archaeological studies within 2km of the activity area	16

Abbreviations

ACHRIS – Aboriginal Cultural Heritage Register and Information System
ASL – Above Sea Level
BLCAC - Bunurong Land Council Aboriginal Corporation
BLSA – Bunurong Land and Sea Association
BWF – Boon Wurrung Foundation
CHA – Cultural Heritage Advisor
CHMP – Cultural Heritage Management Plan
DPC – Department of Premier and Cabinet
EMP – Environmental Management Plan
EVC – Ecological Vegetation Class
LDAD – Low Density Artefact Distribution
LGM – Last Glacial Maximum
OAAV – Office of Aboriginal Affairs Victoria
PAS – Potential Archaeological Sensitivity
PSP – Precinct Structure Plan
RTP – Radial Test Pit
STP – Shovel Test Pit
TP – Test Pit
UFZ – Urban Floodway Zone
VAHR – Victorian Aboriginal Heritage Register
WTLCHC – Wurundjeri Tribe Land Compensation and Cultural Heritage Council Inc

Please note that all maps and plans in this CHMP are prepared using Victorian Government Standard GDA94 MGA (Zone 55) coordinates.

Part One: Assessment

1.0 Introduction

This report outlines the results of an investigation into the heritage values of the Edithvale-Seaford Ramsar Wetlands, and has been conducted as part of the larger Ramsar Management Plan undertaken by Ecology Australia for Melbourne Water.

The purpose of this report is to identify the heritage values associated with the Seaford section of the Edithvale-Seaford Ramsar wetlands, commonly known as the Seaford Swamp, formulate a site prediction model and provide recommendations for the ongoing management of cultural heritage issues within the area. This section of the Edithvale-Seaford Wetlands is also commonly referred to as the Seaford Swamp.

The Edithvale-Seaford Wetland is the largest natural wetland of its type in the Port Phillip and Westernport basins. It is all that remains of Carrum Swamp, which once covered more than 5,000 hectares from Mordialloc in the north to Frankston in the south. This site comprises two separate wetland areas, Edithvale and Seaford. This report will deal with the Seaford section of the Edithvale-Seaford Wetlands.

The wetlands were listed on 29 August 2001 under the Ramsar Convention, as Ramsar site 1096, Australia's 11th site, in recognition of their international importance, and specifically because they:

- Are the last remaining examples of the Carrum Swamp, containing a variety of permanent and seasonal, freshwater and saline wetlands;
- Support populations of the Australasian bittern, considered to be of state significance and threatened in Victoria;
- Support more than 1% of the East Asian – Australasian Flyway population of sharp-tailed sandpipers (or over 2000 birds) in up to one year in three; and
- They are also considered to be of exceptional significance as examples of cost-effective management of wetlands in an urban setting to provide conservation benefits, manage storm water, and encourage environmental education and research

This assessment has utilised standard heritage databases and historic sources to provide a detailed land use history of the study area. A review of previous archaeological assessments has been completed and this along with geological and environmental information has been utilised to formulate a site prediction model and establish the heritage values of the study area.

Location of the Study Area

The Seaford Swamp site covers an area of 158 hectares and is located in the suburb of Seaford (section 1) immediately to the south of Eel Race Creek in the Parish of Lyndhurst, City of Frankston, approximately 35km south of the Melbourne CBD. The study area is bordered to the south by Austin Road, to the east by the Frankston Freeway, a BMX track and green space, to the north by Eel Race Creek and to the west by residential development.

Adjacent to the study area, in the north eastern corner of the study area is the area known as Downs Estate. This area is approximately 21 hectares and has recently been acquired by the council, with the

intention of incorporating this area into the Seaford Swamp area. This area is bordered by the Seaford Swamp site to the west and south, by Eel Race Creek to the north and Old Wells Road to the west. This area is shown outlined in green on Map 1.

Land Manager

The land is jointly managed by Melbourne Water and Frankston City Council, with the wetland areas managed by Melbourne Water and the dryland areas managed by Frankston City Council (KBR 2009)



Map 1: Location of the study area Note: Coordinates provided are in GDA94 MGA (Zone 55)

2. Existing Conditions, Environment and Geology

2.1 Existing Conditions

The study area is the Seaford section of the reclaimed Edithvale Seaford Wetlands and Downs Estate. These wetlands represent the remnants for the former Carrum Swamp that once extended over 5000 hectares, stretching along the eastern shore of Port Phillip Bay from Mordialloc to Frankston. Carrum Swamp was drained in the 1870s to reclaim the land for pastoral and agricultural purposes. Prior to the 1970s, the Seaford Swamp was used primarily for agricultural purposes and a series of excavations was undertaken to improve drainage of the area in the form of construction of drainage ponds and channels (GHD 2005)

Seaford Swamp currently comprises a number of separate water bodies, which serve different purposes, including both fresh water and saline water deposits. The system is highly regulated and requires significant intervention to manage and maintain. The swamp is divided into four sections, these are:

- Semi-permanent brackish pools in the south which receive local stormwater input;
- A deep, permanent, strongly saline pool in the south-western section which has been excavated into the peat layer exposing acid sulfate soils;
- Intermittently and shallowly inundated brackish herb/sedgeland areas in the south; and
- A number of small marshes on the eastern and northern side of the system which receive stormwater from small local catchments (KBR 2009)

While the Seaford swamp was originally a catchment area of Boggy Creek, this has been disconnected and inflows into Seaford Swamp now occur predominantly from the Wadsley Road Drain, to the north of the wetland, which obtains its water from Little Eumemmerring Creek. Additional flows from Eel Race Creek also occur during very high flow periods following significant rainfall events. The surrounding local catchment contributes inflows via stormwater drains while groundwater intrusions, resulting from historic excavations, also provide water to Seaford Swamp (KBR 2009)

The Seaford Swamp is utilised for both drainage purposes, stormwater management and as an open public space popular with local dog walkers, bird watchers and for other recreational uses. The wetlands are home to a significant population of birdlife.



Map 2: Map showing current conditions

2.2 Landforms and Geomorphology of the Study Area


The study area is located in the geomorphic region known as the Barrier Complexes – Discovery Bay, Gippsland Lakes. This is an extensive geomorphic region covering part of eastern Victoria. The landform of the wetlands area has been described as part of the Carrum Lowlands or Sunklands.

Geological mapping of the study area shows that the majority of the study area is located on Coastal Lagoon Deposits (Qg), which are described as dark grey to black silt and clay. There is a thin band of Inland dune deposits (Qd1) located along the eastern margin of the study area. This soil type is described as friable to consolidated sand, silt, and clay, including both lunette and dune deposits (Map 3). While the swampy lagoon deposits show a landform that was previously inundated, the sandy dune deposits located around the margins of the swamp represent a more elevated landform.

This area has undergone significant change overtime as a result of fluctuating sea levels, climate change and irrigation programs.

Between 14,000 and 10,000 years ago climatic conditions changed to become warmer and wetter resulting in rising sea levels. Port Phillip Bay flooded around 10,000 years ago, though formation of the current coastline did not stabilise until 4,000 years ago. Inundation occurred over several thousand years, with French and Phillip Islands not formed until the highest sea level of 5,000-6,000 years ago (Coutts 1977). As current dune barrier systems developed, previous discharge points for the region's rivers and creeks were blocked resulting in a series of freshwater swamps collectively known as Carrum Swamp.

The Carrum Swamp originally stretched from Port Phillip Bay across to Western Port Bay. The water from the Carrum Swamp and the Dandenong and Eumemmerring Creeks drained into Port Phillip Bay via Mordialloc and Kananook Creeks. In the 1870s works were undertaken to drain Carrum Swamp, these works included the creation of Pattersons Cut (today known as Patterson River) to drain Carrum Swamp into the bay. Other drainage points, such as Mordialloc and Kananook Creek also underwent dredging and channelization works to more effectively drain Carrum Swamp into Port Phillip Bay (Murphy & Dugay-Grist 2008).



Map 3: Map showing geology of the study area and surrounds

2.3 Resources Available to Aboriginal People within the Study Area

Plant Resources and Pre-Contact Vegetation

There are three pre-1750 Ecological Vegetation Communities (EVC) that have been identified within the study area. The dominant vegetation type that covered the majority of the study area has been identified as EVC 125 (Plains Grassy Wetland) (Department of Environment and Primary Industries Biodiversity Interactive Map, accessed 02/02/2016) (Map 4).

Plains Grassy Wetland is described as usually treeless, but in some instances can include sparse River Red Gum (*Eucalyptus camaldulensis*) or Swamp Gum (*Eucalyptus ovata*). A sparse shrub component may also be present. The characteristic ground cover is dominated by grasses and small sedges and herbs. The vegetation is typically species-rich on the outer verges but is usually species-poor in the wetter central areas. As the study area would have been either partially or completely inundated, non-aquatic vegetation would have been restricted to the edges of the swamp (Department of Environment and Primary Industries Biodiversity Interactive Map, accessed 02/02/2016).


Pre European contact the northwestern corner of the study area there would have been a small patch of EVC 2 (Coast Banksia Woodland. Coast Banksia Woodland was restricted to coastal or near coastal localities inland behind secondary or tertiary dunes or on sand sheets inland from Coastal Dune Scrub Mosaic. It was usually dominated by an overstorey of Coast Banksia (*Banksia integrifolia* var. *integrifolia*) over tall shrubs of Coast Tea-tree (*Leptospermum laevigatum*). Scramblers such as Bower Spinach (*Tetragonia implexicoma*) were common in the understorey with a groundcover of grasses, herbs and sedges (Department of Environment and Primary Industries Biodiversity Interactive Map, accessed 02/02/2016).

Along the eastern edge of the study area a thin strip of EVC 48 (Heathy Woodland) shown on Biodiversity Interactive Map (Map 4). 1750 EVC modelling is partially based on predictive modelling, however more recent ecological assessment has shown that this area has been incorrectly modelled and actually contains Damp Sands Herb Rich Woodland (EVC 3) (Australian Ecosystems Pty Ltd 2015). Vegetation in this area is considered to represent remnant vegetation. Damp Sands Herb Rich Woodland has been identified along the eastern edge of the study area and along the northern section. Damp Sands Herb Rich Woodland (EVC 3) is described as woodland with a grassy, heathy or bracken-dominated understorey and a ground layer rich in herbs, grasses, and orchids. This EVC occurs mainly on flat or undulating areas on moderately fertile, relatively well drained, deep sand or sandy loam (Department of Environment and Primary Industries Biodiversity Interactive Map, accessed 06/06/2016)..

A recent vegetation assessment of the Seaford Swamp area describes the study area as

“a series of water bodies which are surrounded by Tall Marsh, Brackish Wetland, Plains Sedge Wetland and areas devoid of native vegetation. Terrestrial areas of the site contain patches of Damp Sands Herb Rich Woodland of both remnant and revegetation origins.” (Australian Ecosystems Pty Ltd 2015)

While this EVC mapping shows that prior to European settlement it is likely most of the study area would have been inundated and therefore unsuitable for Aboriginal occupation prior to the draining of the swamp, there were bands of vegetation associated with dune landforms in both the northwestern section of the study area and at the back of raised dunes along the eastern edge of the study area.



Map 4: Map showing 1750s EVC mapping of the study area and surrounds

Information on Fauna of the Activity Area

A wide range of faunal resources would have been present within the study area and the wider region of Carrum Swamp and are likely to have been hunted by Aboriginal people. However, due to the inundated nature of the study area, Aboriginal activity would have been focused around the swamp margins and on elevated areas within the study area extent.

Animals recorded within the Ramsar site include; echidnas, antechinus, bandicoots, possums, wallabies, kangaroos and bats, these resources would also have been available pre-European settlement. Also present were a wide variety of frogs, geckos, skinks, snakes, eels and fish (KBR 2009).

Water Resources

Carrum Swamp would have been the closest fresh water source to the study area

Stone Resources

The lithology of the proposed activity area and surrounding region indicates that quartz would be a highly accessible stone material available to Aboriginal people for the manufacture of stone artefacts. Outcrops of basalt occur at Berwick and to the north siltstone and mudstone exists. Silcrete and chert, which were materials highly sought after by Aboriginal people, were probably obtained from coastal areas such as Mornington Peninsula (Smith 1991).

3. Review of Previous Archaeological Research

The Victorian Aboriginal Heritage Register (VAHR), accessed through Aboriginal Cultural Heritage Register and Information System (ACHRIS), was searched to identify any previously registered Aboriginal Places within a 2km radius of the study area. This search shows that two Aboriginal sites have been identified within the study area, a shell midden and an artefact scatter, and two culturally modified trees have been identified within 200m of the study area. The Seaford Swamp has not been subject to formal archaeological survey.

3.1 Victorian Aboriginal Heritage Register Search

A search of the Victorian Aboriginal Heritage Register identified that there are 33 registered Aboriginal Places within a 2km radius of the study area, comprising a total of 34 components (Table 1). These sites comprise artefact scatters (n=21), object collections (n=2), shell middens (n=4) and scarred trees (n=7) (Table 2). There are two sites located within the activity area and two sites within 200m of the activity area (Map 5).

It should be noted that historically, if it was unclear whether a scarred tree was cultural or not, it was still practice to register them as potential scarred trees. Five of the seven scarred trees are potentially culturally scarred trees. Two of the potentially scarred trees are located within 200m of the study area.

Artefact scatters are the most common site type in the local region, and these sites are generally located around the margins of the former Carrum Swamp within sandy dune deposits and Cranbourne Sands deposits. These sites are located in both surface and sub-surface contexts. Quartz is the most common raw material identified however quartzite, silcrete, chert, basalt and glass artefacts have all been recovered.

One artefact scatter is located within the study area itself. VAHR 7921-0187 is a scatter of stone artefacts located on a ridgeline around the edge of a swampy low-lying area. The recording archaeologist noted that it was likely that this site represented the remnants of a much larger scatter that had likely been collected previously.

There are four registered shell middens within the search region. Two of these shell middens are located on the foreshore dunes on Seaford Beach, one is associated with Kananook Creek and one is located within the study area. VAHR 7921-0372 comprises three dense exposures of mussel and pipi shell deposits eroding from the side of an access track on mudflats. There are a number of characteristics that can be utilised to determine the difference between a cultural and natural shell deposit. These include selection of specific shell species, for example a midden will often represent a collection of specific shell material indicative of selective exploitation of resources, rather than a mix of a wide variety of shell material. Cultural deposits of shell material are also often identified mixed in with charcoal and ash from the cooking process. Shell middens can also display stratification, from numerous episodes of shells being deposited over time. In this case, the archaeologist noted that only two species of shell material were present, suggesting a cultural deposit. The original site card also noted that this site was not in situ, suggested the shell deposit may have been imported into the area.

Within the wider Seaford/Frankston area, shell middens are commonly located along the coastal shoreline and deposits of stone artefacts are frequently located within sandy dune landforms and in deposits of Cranbourne Sands, on elevated ridge lines and in areas associated with the margins of Carrum Swamp. This archaeological record demonstrates that both Carrum Swamp and the coastal area were highly important resources for Aboriginal people and were frequently utilised.

The presence of two Aboriginal sites within the study area also show that despite the Seaford Swamp having previously been part of Carrum Swamp and frequently inundated, fluctuating water levels would have provided access both around and through the swamp. Registered Aboriginal places within the study area are evidence of previous Aboriginal activity within the area.

Table 1: Summary of registered Aboriginal Places within the geographic region

Component Type	Frequency (No.)	Frequency (%)
Artefact Scatter	21	62
Object Collection	2	6
Shell Midden	4	12
Scarred Tree	7	21
Total Components	34	
Total Registered Places	33	

Table 2: Summary details for Aboriginal Places within 2km of the activity area (sites within the study area shown in green, sites within 200m shown in blue).

VAHR ID	Site Name	Site Type	Description
7921-0187	EEL RACE RD 1	Artefact Scatter	Scatter of quartz, chert and ochre located on a sandy ridgeline around the edge of a low lying swampy area. Small scatter of artefacts remain, likely most have been previously removed by collectors.
7921-0242	SEAFORD 1	Shell Midden	Shell midden located on foreshore dune
7921-0294	KANANOOK CREEK 1	Scarred Tree	Banksia Tree in good health with two potentially cultural scars.
7921-0295	KANANOOK CREEK 2	Scarred Tree	Dead gum tree with five potentially cultural scars
7921-0296	SEAFORD 2	Shell Midden	Shell midden located on foreshore dune
7921-0297	DENBIGH RD 1	Scarred Tree	Gum tree in good health with five potentially cultural scars
7921-0298	DENBIGH RD 2	Scarred Tree	Banksia Tree in good health with two potentially cultural scars.
7921-0345	SEAFORD MIDDEN	Shell Midden	Shell midden located on edge of Kananook Creek.
7921-0346	SEAFORD AXES	Object Collection	Two greenstone axes held in private collection, axes found locally in sandy deposits
7921-0372	SEAFORD WETLANDS	Shell Midden	Three dense concentrations of pipi and mussel shell scatted along the edge of an access track.
7921-0530	CARRUM SWAMP 1	Artefact Scatter	Small scatter of silcrete and quartz flakes adjacent Boggy Creek drain
7921-0572	CARRUM SWAMP FOOTHOLD TREE	Scarred Tree	Healthy red gum tree with four scars, probably cultural in origin
7921-0574	WEDGE SAND RIDGE	Artefact Scatter	Large surface scatter of quartz and silcrete artefacts eroding from a sandy dune
7921-0577	COLEMANS	Artefact Scatter	Large scatter of silcrete and quartz flakes

7921-0598	BOGGY CREEK IA1	Artefact Scatter	Two silcrete, one quartz and one mudstone flake identified adjacent Boggy Creek drain.
7921-0639	BOGGY CREEK AS 1	Artefact Scatter	Scatter of quartz, silcrete, basalt and glass flakes on a low sandy rise
7921-0700	PALAROME 1	Scarred Tree	Healthy red gum tree with cultural scar
7921-0701	BOGGY CREEK EASTLINK 1	Artefact Scatter	Eight quartz and quartzite artefacts located on sandy deposits on a floodplain
7921-0702	CARRUM SWAMP 2	Artefact Scatter	Isolated quartzite flake on low sandy rise
7921-0722	BOGGY CREEK AS2	Artefact Scatter	Six quartz and quartzite artefacts located on sandy deposits on a floodplain
7921-0774	COLEMANS ROAD 1	Scarred Tree	Healthy red gum tree with large cultural scar
7921-0777	COLEMANS ROAD 4	Artefact Scatter	Six quartz, two silcrete and one basalt flake located on sandy dune deposits
7921-0792	COLEMANS ROAD 6	Artefact Scatter	Two quartz, three silcrete, two chert and one quartzite flake located on sandy deposits
7921-0793	COLEMANS ROAD 7	Artefact Scatter	Isolated silcrete flake on low sandy rise
7921-0953	PAGETT ROAD 4	Artefact Scatter	Two quartz and one silcrete flake located in disturbed sandy dune deposits
7921-1107	FRANKSTON BYPASS I1	Artefact Scatter	Two silcrete flakes located in a sub-surface context on a small rise.
7921-1108	FRANKSTON BYPASS I2	Artefact Scatter	Nine stone artefacts located in a sub-surface context on a small rise.
7921-1109	FRANKSTON BYPASS I3	Artefact Scatter	Eleven silcrete flakes located in a sub-surface context on a small rise.
7921-1110	FRANKSTON BYPASS I4	Artefact Scatter	No site card found
7921-1230	OLIPHANT WAY AS 1	Artefact Scatter/Object Collection	Three stone artefacts located within the cranbourne sands landform.
7921-1244	Lathams Road 2	Artefact Scatter	Isolated silcrete flake on a flat plain in highly disturbed context
7921-1401	Colemans Road 16	Artefact Scatter	Single artefact retrieved from a depth of 450mm
7921-1399	Colemans Road 17 IA	Artefact Scatter	Single artefact retrieved from a depth of 280mm



3.2 Previous Archaeological Assessment within 2km of the Study Area

A number of reports have been undertaken within the geographic region. The results of relevant regional and localised studies, including CHMPs are presented in Table 3. The results of these reports are utilised to assist in the formulation of the site prediction model.

Discussion of Previous Archaeological Work

A review of previous archaeological research within 2km of the study has noted the importance of Carrum Swamp to Aboriginal people. Drainage of the Carrum Swamp commenced in the 1870s with the creation of Patterson Cut to provide a drainage outlet into the bay. Flooding events widened the cut and by 1879 it was known as Patterson River. It should, however, be considered an artificially created drainage line. Archaeological research conducted within the margins of Carrum Swamp has shown this area is highly unlikely to contain Aboriginal cultural material (Rhodes 1990).

Carrum Swamp was extremely important to Aboriginal people. It provided fresh water and a wide range of flora and fauna resources. Archaeological research has shown that elevated land around the margins of the swamp contains high archaeological potential. The swamp margins would have been heavily utilised as prime camping land and also for hunting and gathering of resources. By contrast, land within the swamp would have been frequently inundated, boggy and low-lying and would therefore not have provided an attractive camping location for Aboriginal people. However, despite this, two Aboriginal sites, a shell midden and a stone artefact scatter have been located within the Seaford Swamp footprint. This shows that the area was utilised by Aboriginal people. Over the years the water levels of the swamp would have fluctuated providing more access in dry years.

Of the 39 CHMPs that have been conducted within a 2km radius of the study area, only six have located Aboriginal cultural material. Those that have located Aboriginal sites have shown that sites are located on relatively intact sandy dune formations and elevated rises associated with either coastal areas or the margins of the swamp. These recorded Aboriginal sites shows that the focus of Aboriginal activity was along the sandy rises around Carrum Swamp, along fresh water sources and along the coastline.

These studies also demonstrate that the impact of urban development has had a substantial impact on the landscape and as such Aboriginal sites are unlikely to be located within areas subject to intensive urbanisation.

Based on the results of previous studies, the most likely site types to occur are artefact scatters (which account for the vast majority of sites recorded in the geographic region), shell middens, and, where suitable trees remain, culturally modified trees.

Previous archaeological research has demonstrated that Aboriginal archaeological sites are likely to be located in close proximity to the coast and/or major waterways on elevated coastal landforms (dune ridges). While the study area is mapped as being within the former extent of Carrum Swamp, the discovery of two Aboriginal sites within the study area suggests that the study area may have archaeological potential where elevated rises and landforms that have not been impacted by substantial ground disturbance occur.

Table 3: Relevant archaeological studies within 2km of the activity area

Study Name	Investigation Type	Results
<p>Proposed Townhouse Development: 48-49 Nepean Highway Seaford</p> <p>(Dugay-Grist, Cowled & McAlister 2015)</p>	CHMP	This CHMP (13448) is a desktop-level report only. A previous CHMP (11837) was conducted for the same property in 2012, however the proposed activity had changed. See discussion of CHMP (11837) below for details.
<p>3CSE-3KEY- Communications Installation Between Pacific Drive and Perry Road to Station Street and Mascot Avenue, Keysborough, Victoria</p> <p>(Lever & Houghton 2015)</p>	CHMP	This CHMP (13295) was conducted for the installation of approximately 9km of telecommunications fibre cable within an activity area predominantly comprising road reserves. The activity area is located within the footprint of the former Carrum Swamp. The site prediction model noted that the majority of previously recorded Aboriginal places are located around the swamp margins and frequently on the sandy coastal plains with ridges and dune fields. The standard assessment noted poor ground visibility and much of the activity area had been significantly disturbed. The complex assessment comprised excavation of two 1m ² test pits and 28 shovel test pits. These all recorded high levels of previous disturbance and no cultural heritage material was located.
<p>Cultural Heritage Management Plan 95 East Road Seaford Townhouse Development</p> <p>(Walther 2015)</p>	CHMP	This CHMP (13430) was conducted for a multi-unit development on a small residential allotment. The desktop assessment shows that there is a moderate likelihood for low density artefact scatters to occur on the coastal plain, inland dune systems and sandy rises. The complex assessment comprised excavation of one 1m ² test pit and two shovel test pits excavated to a depth of 900mm. No Aboriginal cultural heritage material was located, and the upper 300mm of soil deposits were shown to be highly disturbed through previous land use history associated with urbanisation.
<p>5 Kipling St Carrum Residential Subdivision, Complex Cultural Heritage Management Plan</p> <p>(Tunn & Rae 2015)</p>	CHMP	This CHMP (13530) was conducted for a multi-unit development on a small residential allotment. The site prediction model noted that Aboriginal cultural sites were unlikely to be located in areas disturbed by residential development. The standard assessment noted extremely poor ground surface visibility. The complex assessment comprised excavation of two 1m ² test pits and five shovel test pits. Testing noted substantial levels of ground disturbance and no cultural heritage material was located.

Study Name	Investigation Type	Results
<p>Proposed Warehouse Construction at 27 Brett Drive, Carrum Downs (Mitchell & Heddell-Stevens 2015)</p>	CHMP	<p>This CHMP (13632) was conducted for a proposed warehouse. The activity area was located on the Cranbourne Sands landform. The site prediction model stated that there was low to moderate potential for the location of shell material and stone artefacts if intact sandy deposits were located. The complex assessment comprised excavation of a 1m² test pit and three shovel test pits. This testing revealed a highly disturbed sub-surface profile and no Aboriginal cultural heritage material was located.</p>
<p>473 Station Street, Bonbeach, Proposed 12 Unit Development (Barker 2015)</p>	CHMP	<p>This CHMP (13636) was conducted for a multi-unit development on a small residential allotment. The site prediction model noted that previous CHMPs on small urban commercial and residential allotments had demonstrated that it was unlikely Aboriginal cultural material would be located due to substantial ground disturbance caused by the process of urbanisation. Sub-surface testing comprised the excavation of a 1m² test pit and ten shovel test pits. No Aboriginal cultural heritage material was located.</p>
<p>644 Nepean Highway, Carrum Subdivision of Land (Stevens 2015)</p>	CHMP	<p>This CHMP (13730) was conducted for a small residential subdivision located on sandy dune deposits and within 200m of the coastline. The site prediction model stated that land near the coastline on sandy dune deposits contains potential for the discovery of shell middens and stone artefact scatters, however given the level of disturbance in this area from urbanisation, the archaeological potential was low. The complex assessment comprised excavation of a 1m² test pit and four shovel test pits. Sub-surface testing revealed a highly disturbed soil profile and no cultural heritage material was located.</p>
<p>42A Nepean Highway, Seaford: Residential Subdivision (Matic 2014)</p>	CHMP	<p>This CHMP (12936) was conducted for a small residential subdivision. It was considered that the activity area contained potential for the discovery of artefact scatters and shell middens, but that scarred trees were unlikely due to the land use history. The complex assessment comprised a 1m² test pit and four shovel test pits. No Aboriginal cultural heritage material was located.</p>
<p>Proposed Residential Development 34 Melaleuca Drive, Carrum (McAlister 2014)</p>	CHMP	<p>This CHMP (13083) was conducted for a small residential subdivision. The land-use history clearly demonstrated that the activity area had undergone significant ground disturbance through the draining of Carrum Swamp and the mechanical cutting, levelling and filling required to construct the residential development. This involved significant modification of the landform. The area has also been impacted through the construction of a residential dwelling and associated infrastructure. No areas of archaeological potential were identified. As such, a standard and complex assessment was not required.</p>

Study Name	Investigation Type	Results
<p>Proposed 24 Unit Development, 200-202 Nepean Highway, Seaford, Victoria (Cummins, Ward & Power 2014)</p>	CHMP	This CHMP (13189) was conducted for a multi-unit development located east of Kananook Creek. The desktop assessment noted that shell middens and artefact scatters were the most likely site type within the activity area. The complex assessment comprised the excavation of two 1m ² test pits and no Aboriginal sites were located.
<p>88 McLeod Road, Carrum Subdivision of Land (Stevens 2014)</p>	CHMP	This CHMP (13280) was conducted for a small residential subdivision on an allotment that had previously been impacted by construction of a residential dwelling and associated structures. The activity area is located at the transition of a barrier dune/swamp deposit and as such contained moderate archaeological potential due to being located along swamp margins. The standard assessment did not locate any sandy dune landforms, rather a floodplain. The complex assessment comprised the excavation of a 50x50cm test pit and four shovel test pits. Sub-surface testing revealed the soil profile was comprised of swamp deposits and no cultural heritage material was located.
<p>39 Mitchell Street, Seaford, Residential Subdivision (Tunn 2014)</p>	CHMP	This CHMP (13316) was conducted for a small residential subdivision on an allotment that had previously been impacted by construction of a residential dwelling and associated structures. The site prediction model noted that sites may be located in intact dune deposits however, the extent of urban development would negatively impact the potential for archaeological deposits. The complex assessment comprised excavation of two 1m ² test pits. Evidence for ground disturbance was noted in both test pits and no Aboriginal cultural material was located.
<p>23 Chevron Court, Seaford - Subdivision (Murphy & Morris 2013)</p>	CHMP	This CHMP (12437) was conducted for a residential subdivision. This CHMP was triggered by the proximity of the activity area to Kananook Creek. The complex assessment comprised two 1m ² test pits and one shovel test pit. No Aboriginal cultural material was located during this assessment.
<p>83 East Rd, Seaford - Residential Development (Barker 2013)</p>	CHMP	This CHMP (12493) was conducted for a residential development. The site prediction model stated that due to the proximity of Port Phillip Bay and the former Carrum Swamp, the activity area contained potential for the discovery of artefact scatters and shell middens. The complex assessment involved the excavation of a 1m ² test pit and six shovel test pits. No cultural material was located during testing. Significant ground disturbance was noted across the activity area.

Study Name	Investigation Type	Results
<p>178 and 178A Seaford Road, Seaford - Subdivision (Dugay-Grist, Cowled & Maher 2013)</p>	CHMP	This CHMP (12506) was conducted for a multi-dwelling development. The activity area was considered sensitive for the discovery of cultural heritage material due to its location on the Koo Wee Rup Plain. The complex assessment involved the excavation of a 1m ² test pit and five shovel test pits. No cultural material was located during testing. Significant ground disturbance was noted across the Activity Area.
<p>70 Armstrongs Road, Seaford: Residential Development (Laurinda Dugay-Grist, Cowled & Paynter 2013)</p>	CHMP	This CHMP (12667) was conducted for a residential subdivision. The activity area was considered to contain low archaeological potential due to its location within the former Carrum Swamp footprint. The complex assessment comprised the excavation of two 1m ² test pits and five shovel test pits. The sub-surface testing revealed the entire activity area has been subject to substantial ground disturbance and no Aboriginal cultural heritage material was identified.
<p>216 Nepean Highway, Seaford - Residential Development (O'Connor 2013)</p>	CHMP	This CHMP (12580) was conducted for a small residential subdivision. The site prediction model notes that sites are most likely to be located around swamp margins and on elevated rises and sandy dune landforms. The complex assessment comprised excavation of two 1m ² test pits and three shovel test pits. One of the 1m ² test pits was closed due to asbestos fragments. Testing showed disturbance across the activity area and no cultural heritage material was located.
<p>22 Valetta Street, Carrum: Residential Development (L Dugay-Grist, Cowled & Paynter 2013)</p>	CHMP	This CHMP (12667) was conducted for a residential subdivision. The desktop assessment noted the importance of the Carrum Swamp to Aboriginal people. Sandy rises around the swamp margins contain archaeological potential. The complex assessment comprised the excavation of a 1m ² test pit and six shovel test pits. Subsurface testing revealed modern rubbish throughout the top 300mm of the activity area. Soil deposits comprised sandy deposits which became increasingly moist with depth. No Aboriginal cultural heritage material was located during this assessment.
<p>43 Kananook Avenue, Seaford Three lot subdivision (Tunn 2013)</p>	CHMP	This CHMP (12679) was conducted for a small residential subdivision that is currently utilised as a residential dwelling. Geological mapping places the activity area on sandy dune deposits. The site prediction model stated the activity area contained potential for shell middens and stone artefact scatters. The complex assessment comprised a single 1m ² test pit that was excavated to a depth of 1000mm at which point ground water flooded the excavation. Four additional shovel test pits were excavated. Disturbance was noted to depths of 450mm and no Aboriginal cultural heritage material was located.

Study Name	Investigation Type	Results
<p>3 Canberra Street, Patterson Lakes: Residential Subdivision (Matic 2013)</p>	CHMP	<p>This CHMP (12831) was conducted for a residential subdivision. The desktop assessment noted that the activity area was highly unlikely to contain Aboriginal cultural material due to its location within the former Carrum Swamp and the construction methods used in the development of Patterson Lakes which included the stripping of swamp deposits and introduction of fill material to raise the level of the ground for construction. The standard assessment confirmed this conclusion and deemed a complex assessment was not required as the activity area contained no archaeological potential. No Aboriginal cultural heritage material was located during this assessment.</p>
<p>Commercial Subdivision, 28 to 40 Colemans Road, Carrum Downs (Mitchell, Burch & McFarlane 2012)</p>	CHMP	<p>This CHMP (11615) was conducted for a medium sized commercial subdivision. The desktop assessment concluded the area contained high archaeological potential for the discovery of stone artefact scatters which were commonly located nearby and these site types would be most likely to occur in dune and sandy rises landforms. The standard assessment identified three rises within the activity area considered to be areas of high potential. The complex assessment comprised the excavation of five 1m² test pits and 63 shovel test pits. Three Aboriginal sites were located during this assessment (VAHR 7921-1401, -1399, -1406). Two of these sites are isolated artefacts and the third comprised six stone artefacts.</p>
<p>176 to 178, 179 Nepean Highway and 1 Seaford Road, Seaford (Murphy & Thomson 2012)</p>	CHMP	<p>This CHMP (12214) was conducted for the construction of a retail premises located within 200m of the coastline. The standard assessment noted that the activity area had been impacted by the construction of subsequent demolition of two residential dwellings and associated structures. The complex assessment comprised excavation of a 1m² test pit and thirteen shovel test pits. No cultural heritage material was located and sub-surface deposits were found to be highly disturbed.</p>
<p>Eight Townhouse Development, 58 Nepean Highway, Seaford (Burch 2012)</p>	CHMP	<p>This CHMP (12396) was conducted for a multi-unit development. The site prediction model noted that the site types which may occur within the activity area are artefact scatters, shell middens and burials. The standard assessment noted that most of the activity area had undergone disturbance through the construction of a residential dwelling and associated works. However a dune landform was identified. The complex assessment comprised two 1m² test pits and five shovel test pits. Subsurface testing revealed disturbed deposits and no Aboriginal cultural heritage material was located.</p>

Study Name	Investigation Type	Results
<p>48 Nepean Highway, Seaford: Residential Subdivision (Dugay-Grist & McAlister 2011)</p>	CHMP	<p>This CHMP (11837) was conducted for a small residential subdivision. The desktop assessment noted that the land had previously been disturbed through the construction and subsequent demolition of a residential dwelling. The site prediction model noted that the activity area contains the potential for artefact scatters in areas that have been subject to minimal disturbance. The standard assessment noted poor ground surface visibility but did identify a sandy rise considered to contain archaeological potential. The Complex Assessment comprised a 1m² test pit, four mechanically excavated transects and four shovel test pits. The subsurface testing demonstrated disturbance across the Activity Area to depths of between 900-1200mm. No Aboriginal cultural material was located.</p>
<p>Lathams Road, Carrum Downs Industrial Development (Howell-Meurs 2011)</p>	CHMP	<p>This CHMP (11444) was conducted for an industrial development. The site prediction model stated that sites are likely to be located around the margins of swamps and on elevated sandy rises. The standard assessment identified two surface scatters of Aboriginal stone tools. The complex assessment comprised excavation of two 1m² test pits and 78 shovel test pits. A total of five Aboriginal places were located, all of which comprise either isolated artefacts or low density artefact scatters.</p>
<p>37 Fellowes Street, Seaford Residential Development (O'Reilly & Dudley 2011)</p>	CHMP	<p>This CHMP (11649) was conducted for a proposed residential subdivision potentially located within the Cranbourne Sands landform. The desktop assessment concluded that the most likely site type in the area would be low density stone artefact scatters of quartz and silcrete. The importance of determining whether the landform was Cranbourne Sands was also emphasised. The lack of cultural material identified during the standard assessment is considered to reflect the lack of ground surface visibility and past ground disturbance. During the complex assessment subsurface testing was conducted within the activity area in areas of archaeological sensitivity that were to be impacted by the proposed development. The complex assessment comprised the excavation of a 1m² test pit and four shovel test pits. Subsurface testing indicated that the Cranbourne Sands were present in the activity area, however that the area had undergone disturbance due to residential development and tree clearance. No Aboriginal cultural heritage material was located during this investigation and the activity area is considered to have low potential for Aboriginal cultural material.</p>

Study Name	Investigation Type	Results
78-83 Nepean Highway, Seaford: Proposed Residential Development (Matic 2011)	CHMP	This CHMP (11762) was conducted for a proposed residential development within a sandy dune landform. The complex assessment comprised excavation of a 1m ² test pit and five shovel test pits. Sub-surface testing revealed high levels of disturbance across the upper soil profile and no cultural heritage material was located.
75 Frankston Gardens Drive, Carrum Downs - Subdivision (Whincop 2011)	CHMP	This CHMP (11908) was conducted for an industrial subdivision. The site prediction model noted that the activity area is located on lands that encompassed the former margins of Carrum Swamp and as such contain moderate archaeological potential. The complex assessment comprised the excavation of a 1m ² test pit and twelve shovel test pits. Sub-surface testing revealed high levels of disturbance across the soil profile and no cultural heritage material was located.
Residential Development, 1-3 Attunga Crescent, Seaford (Dudley et al. 2010)	CHMP	This CHMP (10940) was conducted approximately 1.2km south of the current activity area. The desktop assessment indicated that the activity area was located on the natural edge of the former Carrum Swamp and as such may contain archaeological potential. The complex assessment comprised the excavation of a 1m ² test pit and six shovel test pits. Testing revealed sandy deposits and no Aboriginal cultural material was located.
Lot 2, 28 Oliphant Way, Seaford Factory, Warehouse and Office Development (Griffin & McAlister 2010)	CHMP	This CHMP (11343) was conducted for an industrial subdivision. The desktop assessment indicated that the activity area was located within the former swamp and dune area that encompasses both the old Carrum Swamp and the undulating sand and plain ridge complex known as the Cranbourne sands. The relatively undeveloped nature of the activity area suggested moderate to high potential for Aboriginal cultural material. The standard assessment noted some exposed areas but largely poor ground surface visibility. Two artefacts were located in an exposed area on the Cranbourne Sands landform and registered as VAHR 7921-1230. The complex assessment comprised excavation of a 1m ² test pit and forty 40x40cm shovel test pits laid out in a 10m x 10m grid across the activity area. The complex assessment identified one additional artefact associated with VAHR 7921-1230 and defined the site extent. The complex assessment also confirmed both the presence of the Cranbourne sands and former swamp landforms within the activity area.

Study Name	Investigation Type	Results
<p>95-96 Brunel Road, Seaford: Industrial Subdivision (Dugay-Grist & Dudley 2010)</p>	CHMP	<p>This CHMP (11382) was conducted for an industrial subdivision. The desktop assessment indicated the activity area was within the Cranbourne Sands landform and contained moderate potential for archaeological material. The standard assessment identified areas of disturbance within the western section of activity area. The complex assessment comprised a 1m² test pit and eleven 40x40cm shovel test pits. This testing revealed the area had been subject to significant ground disturbance and no Aboriginal archaeological material was identified during this investigation.</p>
<p>Proposed Retirement Village and Carpark, 45 Fellowes St, Seaford (Young & Prossor 2010)</p>	CHMP	<p>This CHMP was conducted for a proposed retirement village. The desktop assessment noted that archaeological sites are likely to occur on areas of well-drained soil and low rises which are less prone to flooding. The standard assessment noted poor surface visibility due to dense grass coverage, existing buildings and a gravel driveway. The complex assessment comprised excavation of a 1m² test pit and twenty-three 40x40cm shovel test pits. The complex assessment noted relatively undisturbed, deep sandy deposits; however no Aboriginal cultural material was identified.</p>
<p>Frankston Bypass EES, Mornington Peninsula (Long et al. 2009)</p>	CHMP	<p>This CHMP (10015) was conducted for the Peninsula Link Freeway Project. The Peninsula Link project involved the construction of 25km from the Mornington Peninsula Freeway/EastLink interchange at Carrum Downs to the Mornington Peninsula Freeway at Mt Martha. The desktop assessment concluded that the zones of highest Aboriginal heritage sensitivity within the activity area consisted of the margins of Carrum Swamp in the north and a zone surrounding Devil Bend Creek in the south. The standard assessment identified two new Aboriginal cultural heritage places, VAHR 7921-1097 and 7921-1100; both diffuse artefact scatters on a slope overlooking Devil Bend Creek. Limited ground surface visibility was noted across the activity area. The complex assessment comprised eleven 1m² test pits and 630 shovel test pits excavated to a maximum depth of 700mm. The subsurface testing program was designed to sample representative landscapes and soil profiles across the activity area with particular focus on landforms of predicted archaeological potential. Twenty-six stone artefact scatters were located over 21 separate testing areas, ranging from diffuse artefact scatters, to extensive artefact scatters containing a large number of artefacts. The majority of sites were located within the area of Devil Bend Creek and its associated valley and tributary, with a large number of diffuse and variable/high density sites being located in these areas. Diffuse scatters were also located around Boggy Creek, across the Baxter Sand Plain, and in the Tamarisk and Tuerong Creek valleys. Only stone and glass artefacts were retrieved, no other cultural heritage material (such as shell or bone) was identified.</p>

Study Name	Investigation Type	Results
<p>300 Frankston-Dandenong Road, Seaford (Schell & Barker 2009)</p>	CHMP	<p>This CHMP (10583) was conducted for an industrial subdivision located within Cranbourne Sands landform. The desktop assessment noted that the activity area had been subject to ground disturbance due to construction and industrial activities that had taken place on site however it was concluded that intact archaeological deposits may occur beneath the disturbed zones. The standard assessment noted poor ground surface visibility and failed to identify any landforms of archaeological sensitivity, it did however note locations that appeared to have been subject to fewer surface and subsurface impacts and focused subsurface testing in these areas. The complex assessment comprised two 1m² test pits and twenty-one shovel test pits. The assessment confirmed the presence of potentially sandy subsurface deposits and also located European debris across the site to an average depth of 500mm. No Aboriginal archaeological material was located during this assessment.</p>
<p>Seaford Village, Seaford (Dugay-Grist 2009)</p>	CHMP	<p>This CHMP (10637) was conducted for the Seaford Village. Seaford Village is located between Kananook Creek and the Nepean Highway. The desktop assessment indicated that the activity area had been subject to high levels of significant ground disturbance including significant work on the banks of Kananook Creek to redefine the flow of the creek, however the extent and method of the works was unknown. The standard assessment recorded poor ground surface visibility and high levels of disturbance within much of the activity area; however several sections were identified as retaining archaeological potential. The complex assessment comprised excavation of four 1m² test pits and nine shovel test pits. All subsurface testing was conducted in areas that had not been subject to prior significant disturbance and that would be impacted by the proposed activity. Excavation went to a maximum depth of 1m as further excavation in sandy deposits was considered unsafe. Subsurface testing revealed extensive previous ground disturbance and no Aboriginal archaeological materials were located within the activity area.</p>
<p>Construction of New Dwellings at 211-212 Nepean Highway, Seaford (Tucker & Marshall 2009)</p>	CHMP	<p>This CHMP (11036) was conducted for a small residential subdivision. The activity area is bounded to the east by Kananook Creek and underlain by coastal dune and beach deposits. The desktop assessment noted that areas along Kananook Creek showed high potential for archaeological material. Six test pits were excavated during the complex assessment; however no Aboriginal cultural material was identified. Evidence of prior disturbance was noted and the activity area was determined to be highly disturbed.</p>

Study Name	Investigation Type	Results
<p>8-10 Klauer Street (Industrial Subdivision), Seaford (Howell-Meurs 2008)</p>	CHMP	<p>This CHMP (10353) was conducted for an industrial subdivision located within the Cranbourne Sands landform. The standard assessment determined that the majority of the activity area had undergone a variety of disturbances associated with the construction of a paved car park area, golf driving range and the installation of a telecommunications tower and cable. The complex assessment comprised seventeen shovel test pits to a maximum depth of 1m in sandy soil. No Aboriginal material was located during this investigation.</p>
<p>Commercial Subdivision at 107 Boundary Road, Carrum Downs, Victoria (Mitchell & Richmond 2008)</p>	CHMP	<p>This CHMP (10591) was conducted for a commercial subdivision. Previous archaeological assessment had identified four Aboriginal sites, all stone artefact scatters within areas that had been impacted through earthmoving works. The complex assessment comprised excavation of one 1m x 50cm test pit, 24 auger holes and fifty shovel test pits. The complex assessment targeted the remains of the two sandy rises within the activity area that were considered to retain archaeological potential and were the home of three of the four sites. A fifth site was located on another small sandy rise and three of the sites were shown to be one larger site. The sites all comprised low densities of stone artefacts.</p>
<p>Boggy Creek Wetlands, Carrum Downs (Vines et al. 2007)</p>	CHMP	<p>This CHMP (10014) was conducted for works to a large wetlands area to the east of the current study area. This area had been subject to prior archaeological assessment with a survey in 2004 (Tulloch 2004) that identified two low density artefact scatters. Subsequent sub-surface testing revealed this site was a high density artefact scatter contained over 200 artefacts (Tulloch 2004). As such, recommendations were made to protect this site.</p>
<p>Frankston City Council Aboriginal Cultural Heritage Assessment (Marshall 1998)</p>	Field Survey	<p>This study involved a detailed desktop assessment and field survey. Its results were utilised to provide a site prediction model and provide assistance in determining archaeological potential for planners within the City of Frankston. The review showed that Frankston contained only a small number of archaeological sites and these results are not consistent with the wider area. This paucity of sites was attributed to the high degree of residential development in the Frankston area and may also reflect the lack of archaeological surveys. The site prediction model developed for this survey states that sites might be located along the bay and on the tops of inland sand dunes. These sites are likely to be artefact scatters and shell material.</p>

Study Name	Investigation Type	Results
<p>Dandenong Creek and Patterson River Melbourne: An Archaeological Survey</p> <p>(Rhodes 1990)</p>	Field Survey	<p>This report details the results of a survey along Dandenong Creek and Patterson River. It notes that land to the north and south of Patterson River comprises flat plains and drained swamp land associated with the former Carrum Swamp. Swamp reclamation works in the 1870s resulted in the channelization of southern sections of Dandenong and Eumemmerring Creeks and the cutting of a channel to drain the creeks and swamp into Port Phillip Bay. This channel was enlarged during 1879 by a storm and became the Patterson River. The report notes that there are very few archaeological sites within the boundary of the former Carrum Swamp, and those that do exist are scarred trees. Artefact scatters within the low-lying plains are located within the former Holocene dune belt, separating the swamp from Port Phillip Bay. The field survey did not inspect Patterson River due the Desktop Assessment noting that sites were extremely unlikely to be located adjacent to Patterson River. It is an artificially created river within the margins of a former swamp and as such it would have been inundated and unsuitable as a campsite for Aboriginal people.</p>
<p>Sites of Archaeological Significance in the Western Port Catchment</p> <p>(Gaughwin 1981)</p>	Field Survey	<p>Gaughwin recorded 266 Aboriginal archaeological sites, 13 of which were found within an area referred to as “Top of the Bay” (a landform that equates to Sullivan’s “northern plains”). Near the present Activity Area, the highest site and artefact densities were found to occur on the sandy ridges of the Cranbourne area, particularly those associated with water. Gaughwin determined that the sites located on these sand ridges are situated to take advantage of resources associated with swamp depressions. The site prediction model formulated by Gaughwin for the “Top of the Bay” landform is applicable to the current Activity Area.</p> <p>Gaughwin predicted artefact scatters and isolated artefacts are the most likely site type to occur within this unit. Most sites will be within 100m of a water source, including rivers, creeks, swamps, ponds, springs, coastline, lagoons and soaks. The highest site densities will be found in the Cranbourne Sands and high dry ground such as ridges and hummocks. Lowest site densities will be found along the foreshore and in low-lying areas such as past swamps. It is highly unlikely that scarred tree sites will be located within the region due to the lack of suitable trees. Surface scatters will be dominated by silcrete, quartz and chert artefacts.</p>

4. Land Use History and Ethnography

4.1 Historical and Ethno-Historical Accounts in the Geographic Region

Indigenous people have likely occupied and utilised the landscape around Melbourne for over 40,000 years, although there is very little known about human occupation and palaeo-environments in the Melbourne Region until 10-15,000 years ago.

Prior to European settlement, the Indigenous population of Victoria was divided into a number of groups. In central Victoria, the languages of each of these groups were dialectically similar, sharing up to 93% of their vocabularies. These dialect groups, named the Woiworung, Bunurong, Taungurong, Nguruiwurung, Wada wurrung and Dja dja wurrung, comprised a 'regional cultural confederacy known as the Kulin' (Barwick 1984, pp.104–105). Members of the Kulin confederacy shared economic and social relationships, common religious beliefs, creation legends and Dreamtime ancestors. Each person also belonged to one of two moieties named after the Dreamtime ancestors, Bunjil, the eaglehawk, and Waa, the crow. Moiety affiliation was determined at birth with the child adopting the moiety of the father's clan. Kulin people were exogamous, meaning they married outside their clan and to a person from the opposite moiety (Barwick 1984, p.106; Clark 1990).

The study area is located within the land of the *Bun wurrung* people. The *Bun wurrung* (Bunurong) were one of four 'dialectal tribes' comprising the East Kulin language group (Barwick 1984). William Thomas, Assistant Protector of Aborigines in the Western Port district (1839 – 1849), spent much of his time travelling with *Bun wurrung* people between his hut at Arthur's Seat, the Aboriginal reserve which he established at Narre Narre Warren during 1841 and the Aboriginal camps around Melbourne (Sullivan 1981, p.25; Cannon 1983). This extensive travel through *Bun Wurrung* territory enabled Thomas to argue that the *Bun Wurrung* claimed "all the country south of the Yarra River, whose creeks and inlets fall into the sea from the Werribee River west to the Tarwin River, east of Cape Patterson" (Thomas papers Vol.7 17/1/1860).

The study area was the country of the *Mayone buluk* meaning 'people of the swamp'. Their territory is thought to have been "Carrum Swamp, the coastal strip at the head of Western Port Bay and the upper portion of the Mornington Peninsular" (Barwick 1984: 177).. The *arweet* (clan head) was identified as Motrungo / Budgery Tom (c. 1797/8-1848). His heir was Buggup / Buckup (c. 1820/3-1848), a corporal in the Native Police Corps (Clark 1990, p.367; Barwick 1984, p.117). The moiety affiliation of the *Mayone bulluk* was Bunjil (Clark 1990, p.367).

Post-contact historical accounts

Much of the available information about traditional *Bun wurrung* culture has been provided by William Thomas, Assistant Protector of Aborigines for the Western Port District. Other sources of information are the journals and letters of G. A. Robinson, Chief Protector of Aborigines (1839 – 1850); A. W. Howitt and early settlers such as Georgiana McCrae and Maurice Meyrick (Bride 1969; Presland 1977; McCrae 1992).

First contact between *Bun wurrung* people and Europeans was with whalers and sealers prior to 1803 (sealing began in Bass Strait in 1798), as European huts were discovered by Grimes at Boneo in January of that year (Rogers 1957). The fact that whalers and sealers had arrived early in the history of the colony is further substantiated by entries in Robinson's journal (26 December 1836 cited in (Plomley 1987, p.405), where he records an account of the early kidnapping of *Bun wurrung* women:

Matilda the VDL native woman pointed out the spot a few miles down the harbour at Point Nepean where she said George Meredith and his crew of sealers stole the native women. The men's names were Brown, Mr. West the master of the schooner, and a man named Billy...said there was plenty of black fellows, some on the Port Phillip side some outside, sea coast. Said the sealers were afraid of the Port Phillip natives. Said they employed her to entice them. George Meredith stole the, I think she said, four women, took them...and then sold them to the [other] sealers there. I am informed that Munro bought one. (G. A. Robinson Journal entry, 26 December 1836, in (Plomley 1987, p.405).

During 1839, the British Colonial Government established an Aboriginal Protectorate in what is now Victoria. A Chief Protector, G.A. Robinson and four Assistant Protectors were appointed to administer the Protectorate. William Thomas was appointed Assistant Protector for the Port Phillip and Westernport Districts and had extensive contact with traditional *Bun wurrung* people during the early years of European settlement (Cotter 2005; 1-2). Thomas travelled with groups of *Bun wurrung* and related Woiworong people on seasonal movements around the Mornington Peninsula and Westernport, producing a map in 1841, showing the locations of *Bun wurrung* campsites and routes of movement.

Thomas also noted the frequent exploitation of the Nepean Peninsula coast. Sullivan ascertains that some of the places mentioned by Thomas could be base camps including *Boniong* and *Turtguruk* located in the hinterland of the Nepean Peninsula (1981, p.29).

Shellfish gathering was observed by Thomas in Port Phillip Bay, near Melbourne, when he reported that a group of women went at least three times a week to collect shellfish (Thomas cited in (Sullivan 1981, p.25). Cockle and mussel shells were observed in Aboriginal huts on the Nepean Peninsula (Knopwood cited in (Sullivan 1981, p.36). Rough waves pound the Bass Strait coastline of the Nepean Peninsula, particularly during winter, and this may have made the collection of shellfish difficult at this time of year and restricted activities to the warmer months (Sullivan 1981, p.8). Thomas also observed shellfish being collected by women diving in creeks and rivers (Thomas cited in (Sullivan 1981, p.28).

In 1840, Thomas (cited in (Sullivan 1981, p.30) observed that about 44 people were still exploiting the country between Mount Martha and Cape Schanck, divided into small groups of about six to seven people. Larger groups of 20 to 30 people would have gathered where resources were plentiful or concentrated (Sullivan 1981, p.32).

Thomas considered that the *Bun wurrung* economy was predominantly terrestrial in orientation despite their close association with the coast (Gaughwin & Sullivan 1984, p.89). However, it is possible that this was partially a reflection of the time of year that he made his observations. He noted that favourite foods were kangaroo and possum, and that they had the 'greatest abhorrence' for snakes (Thomas cited in (Sullivan 1981, p.22). Robinson was told by an eyewitness that in about 1834, "there was a tribe of Natives on the Point hunting kangaroo" (Robinson cited in (D'Arcy 2005, p.28). Women caught many smaller creatures such as bandicoots, rats and lizards (Thomas cited in (Sullivan 1981, p.22). It is probable that many women's subsistence and other activities were not seen or recorded by Thomas, either through his lack of interest or because the women carried them out away from the presence of men.

Fires were commonly lit by Aboriginal people in the coastal area of Port Phillip and were seen by early explorers (Sullivan 1981, p.23). While in the vicinity of Arthurs Seat in January 1803, James Flemming noted that the area was 'all newly burnt' (Shillinglaw 1972, p.24). In December 1836, George Augustus Robinson also noted at Point Nepean that "the bush was on fire, as indeed it was in all parts of the country" (Plomley 1987, p.405). Fires were mainly lit to maintain pathways through dense scrub, to

increase the fertility of the land, to drive game and quite probably as a smoke screen to hide behind or to warn off or confuse intruders.

Further historical accounts of *Bun wurrung* traditions were recorded by Georgiana McCrae. Andrew McCrae and his wife Georgiana built a homestead at Arthur's Seat. Georgiana recounts that *Bun wurrung* people camped on the "edge of Cape Schanck Road" (McCrae 1934, p.209) and also mentions that large groups of *Bun wurrung* people made seasonal camps between the Cape Schanck Road and the bay beneath tea trees or coastal banksia (McCrae 1934, p.194). In 1851, Georgiana observed a *Bun wurrung* burial outside the paddock fence on the edge of the Cape Schanck Road:

I watched the grave being dug by some, while others wrapped a possum-rug about the corpse, which they interred in a sitting position, the elbows on the knees, the chin supported in the left hand, and the opposite one laid, with the fingers open, along the angle of the jaw. Cords were drawn tightly across the shoulders and around the waist, then a new pannikin and the last of the bottle of medicine I had sent him having been put into the grave, the father, and (fifth) stepmother filled the hole with sand. (McCrae 1992, p.213)

Indigenous travel routes were located throughout Bunurong country. Thomas' 1840 map below (Fig. 1) shows some of these tracks, it also shows Carrum Swamp, labelled 'lagoon'.




Figure 1. 1840 map by William Thomas indicating routes taken on his travels with *Bun wurrung* tribes. (Reproduced from Presland 1994, p. 74)

There is considerable emphasis in the ethnographic literature on the use of creeks and waterways as corridors of movement. Gunson (1974: 10) stated that members of the *Mayune balug* usually camped beside waterholes, creeks, and at coastal locations. Early settlers of the Western Port region also noted that Aboriginal campsites containing huts were often found beside rivers and creeks (Sullivan 1981: 33). Carrum Swamp would have been an important resource for Aboriginal people and frequently visited.

Just prior to, and overlapping, the period of British exploration and settlement, the *Bun wurrung* were involved in a long-running dispute with the *Gunai/Kurnai* people from Gippsland. According to William Barak, the conflict was a dispute over resources, which resulted in heavy casualties being suffered by the *Bun wurrung*. Many *Gunai/Kurnai* raids occurred to abduct *Bunurong* women. According to Barwick (1984), the *Yowengerra* had almost been completely annihilated by 1836, largely as a result of attacks from the *Gunai/Kurnai*. During 1833 - 1834 around 60-70 *Bunurong* people were killed in a raid by *Gunai/Kurnai* while they were camped to the north of Carrum Swamp (Rhodes 2003).

Warfare, disease and kidnapping all conspired to precipitate a rapid decline in the population of *Bun wurrung* peoples inhabiting their traditional lands. These numerous impacts would also have disrupted the economic and social ties binding the *Bun wurrung* to neighbouring *Kulin* peoples. Consequently, much of the *Bun wurrung* population had been displaced by the time of permanent European settlement of the area. By 1839 the *Bun wurrung* had been reduced to 83 people, with only 4 of 19 children under four years old, from a probable pre-contact population of greater than 300 people. By 1850 Protector William Thomas estimated just 28 *Bun wurrung* people remained (Rhodes & Bell 2004).. A camping reserve was established in 1841 at Mordialloc Creek where the *Bun wurrung* peoples were allocated 340 hectares. The rations for the reserve were managed by the Honorary Correspondents to the Boards of the Protection of Aborigines. The majority of the *Bun wurrung* population had moved to the Mordialloc reserve by 1856 which ceased operation in 1878.

Today the *Bun wurrung* people are represented by the *Bunurong Land Council Aboriginal Corporation*, *Bunurong Land and Sea Association* and the *Boon Wurrung Foundation Ltd*, who regard themselves as the Traditional Owners and custodians of this area.

4.2 Post-Contact Land Use History of the Study Area

The study area is located within the footprint of the former Carrum Swamp. The Carrum Swamp was an extensive fresh water wetlands area stretching from Mordialloc to Frankston. Drainage of the Carrum Swamp commenced in 1876 with the construction of Patterson Cut, now known as the Patterson River. In 1879 public works were conducted to create efficient drainage of the swamp to open up the land for settlement. By the early twentieth-century most of the original wetlands area had been drained and land was being used for market gardens and dairy farming. By the 1960s the former swamp area had been largely urbanised.




Figure 2: Map showing location of Edithvale-Seafood Wetlands in blue and extent of the former Carrum Swamp in grey.

The study area has been significantly impacted by its land use history. Prior to European contact the study area was located at the southern end of the Carrum Swamp. An 1871 parish map shows the location of the swamp and the plan for division of land following drainage of the swamp (Figure 3).




Figure 3: 1871 Parish Map of Carrum Swamp (source: TROVE)

After Carrum Swamp was drained in the 1870s, land was opened up for farming, however it is likely that this area would have regularly been waterlogged and frequently inundated when heavy rains occurred as Google Earth shows the area as being between -1m to 1m above sea level.

Historical aerial photography from 1945 shows the study area as cleared open farmland with minimal vegetation. However in this image it can be seen that the area is still frequently waterlogged and there are pools of water visible in the southern end of the study area (Figure 4).

In 1956 the study area has been partially inundated with pools of water forming in both the south sections of the wetlands. The northwest section appears to contain what is a large sand dune. Residential subdivision is occurring along the Nepean Highway and to the southwest of the study area (Figure 5).

By 1968, water is pooling in both the northern and southern end of the wetland area and revegetation is happening along the eastern edge. Residential subdivision to the west has proceeded rapidly (Figure 6).

By 1972 water is still pooling in the paddocks, however the paddocks are clearly being ploughed and utilised with crop lines and furrows clearly visible (Figure 7).

By 1989 the study area is clearly being used as a drainage and wetlands area and vegetation cover around the pools of water is green. The area around the study area has largely all been converted to either residential or industrial development (Figure 8).

Since 1989 (Figure 8) significant modifications have occurred to the wetlands area, multiple pools have been constructed along with a series of drains and weirs to regulate water flow and manage storm water. Along with these modifications, a series of trails have been constructed throughout the wetlands. The northern end of the wetlands has undergone a program of revegetation (Figure 9).




Figure 4: 1945 map showing location of study area and surrounds



Figure 5: 1956 map showing location of study area and surrounds




Figure 6: 1968 map showing location of study area and surrounds




Figure 7: 1972 map showing location of study area and surrounds




Figure 8: 1989 map showing location of study area and surrounds



Figure 9: 2009 map showing location of study area and surrounds

5 Conclusions and Recommendations

5.1 Site Prediction Model

This desktop cultural heritage assessment for the study area has allowed a site prediction model to be developed. A site prediction model is intended for use as an indication of the types of archaeological sites that may occur in a given area.

The following statements can be made about the current study area:

- The location of previously registered Aboriginal Places within a 2km radius shows that Aboriginal sites can be located on elevated landforms around the edges of low-lying swampy areas, sites can also be located on sandy dune systems;
- Two Aboriginal sites have been located within the study area, one a large scatter of shell material and the second a scatter of stone artefacts;
- A shell midden site has been located within the study area, this site was located on a mud flat. The recording archaeologist noted the site was not in situ. Further archaeological investigation of this site would be recommended to determine if the site represents a natural or cultural deposit of shell material;
- One artefact scatter has been located within the study area, this site is a scatter of surface artefacts on a ridgeline considered to be the remainder of a much larger deposit;
- Aboriginal site types are primarily low density artefact scatters and isolated artefacts, representing transitory use of the area rather than significant camping events;
- Raw materials are primarily quartz and quartzite, however some silcrete artefacts have also been located;
- These two sites suggest that the study area was not completely inundated by Carrum Swamp or subject to fluctuations in sea level, and as such the study area may contain archaeological potential;
- Geological and EVC mapping shows that both the northwestern section of the study area and the eastern margin of the study area were once elevated ridgelines around the margins of low lying swampy areas and as such these areas should be considered to contain moderate to high archaeological material;
- The lower lying sections of the study area should be considered to contain low archaeological potential;
- The study area contains potential for the discovery of deposits of shell material and/or stone artefacts;
- Historical mapping shows that the majority of the study area has been cleared of native vegetation, with the exception of a small stand of remnant native vegetation in the north

western section. Areas of remnant native vegetation should be considered to contain the potential for culturally scarred trees;

- Post Contact, the study area has been significantly modified through the drainage of Carrum Swamp, the clearance of native vegetation for agricultural purposes and use of the land for drainage and stormwater management through the creation of a series of ponds, weirs and drainage channels. The study area is also currently utilised for recreational purposes such as dog walking and bird watching and has been impacted by the construction of recreational structures such as walking trails.

The site prediction model concludes that there is potential for further deposits of Aboriginal cultural material within the study area. Areas at higher elevations, such as land along the eastern boundary and in the northwestern corner should be considered to contain moderate to high Aboriginal archaeological potential. Lower-lying areas should be considered to contain lower archaeological potential. The archaeological potential will also be dependent on the level of ground disturbance which has occurred within any given area.

A field survey of the study area would be able to provide a more detailed archaeological assessment and identify areas of archaeological potential along with any surface Aboriginal sites within the study area.

5.2. Recommendations and Obligations under the Aboriginal Heritage Act

Seaford Swamp has not previously been subject to detailed archaeological assessment. While two Aboriginal sites have been located, these have never been thoroughly investigated and their current condition is unknown. Further archaeological assessment of these site can help determine their extent and significance. Additionally, while the desktop assessment can identify areas of higher archaeological potential, this can be confirmed by a field survey which would allow for a more detailed archaeological assessment that is outside the scope of this desktop report.

It is recommended that the land manager may wish to undertake a more detailed archaeological assessment of the study area in the form of a field survey in order to;

- Further investigate the registered Aboriginal places within the study area;
- Identify any surface deposits of Aboriginal cultural heritage material;
- Identify areas of archaeological potential; and
- Further investigate the extent of ground disturbance within the study area.

These results could then be utilised to assist with the future management of heritage issues within the Seaford wetlands area. This would be particularly useful for any activities that do not trigger a mandatory CHMP and could prevent accidental harm to unrecorded Aboriginal cultural heritage material.

For any planned activities within the study area that constitute a high impact activity, a CHMP will be required.

The Aboriginal Heritage Regulations 2007 state that:

A cultural heritage management plan is required for an activity if—

- (a) all or part of the activity area for the activity is an area of cultural heritage sensitivity; and*
- (b) all or part of the activity is a high impact activity. (r. 6)*

The study area is considered an area of cultural heritage sensitivity because it is a declared Ramsar wetland (Aboriginal Heritage Regulations (r. 26)).

Declared Ramsar wetlands

(1) Subject to subregulation (2), a declared Ramsar wetland or land within 200 metres of a declared Ramsar wetland is an area of cultural heritage sensitivity.

(2) If part of a declared Ramsar wetland or part of the land within 200 metres of a declared Ramsar wetland has been subject to significant ground disturbance, that part is not an area of cultural heritage sensitivity.

As such, if works that involve a high impact activity within the study area are planned, they would trigger a mandatory CHMP. The definition of a high impact activity is presented within Division 5 of the *Aboriginal Heritage Regulations 2007* (Appendix 1).

References

Legislation

Aboriginal Heritage Act 2006

Aboriginal Heritage Regulations 2007

Organisations

Land Victoria Aerial Photography storage facility (Laverton)

Public Record Office Victoria

Internet Resources

Department of Environment and Primary Industries – Biodiversity Interactive Map
(<http://www.depi.vic.gov.au/environment-and-wildlife/biodiversity/biodiversity-interactive-map>)

Department of Environment and Primary Industries – EVC Benchmarks
(<http://www.depi.vic.gov.au/environment-and-wildlife/biodiversity/evc-benchmarks>)

Department of Environment and Primary Industries – GeoVic
(http://er-info.depi.vic.gov.au/sd_weave/registered.htm)

National Library of Australia (TROVE)
(<http://trove.nla.gov.au/>)

Department of Environment and Primary Industries – Victorian Resources Online
<http://vro.agriculture.vic.gov.au/dpi/vro/>

Reports and Published Resources

Australian Ecosystems Pty Ltd 2015, *Melbourne Water: Sites of Biodiversity Significance: Vegetation Assessments 2014-2015 DRAFT REPORT*, Australian Ecosystems Pty Ltd, Port Melbourne.

Barker, M 2013, *83 East Road, Seaford – Residential Development*, Benchmark Heritage Management, Kinglake.

Barker, M 2015, *473 Station Street, Bonbeach, Proposed 12 Unit Development: Desktop, Standard and Complex Assessments*, Benchmark Heritage Management Pty Ltd, Kinglake.

Barwick, D 1984, 'Mapping the Past: an Atlas of Victorian Clans 1835-1904. Part 1', *Aboriginal History*, vol. 8, no. 1-2, pp. 100–131.

Bride, TF 1969, *Letters from Victorian Pioneers*, Robert Brain, Melbourne.

Burch, J 2012, *Eight Townhouse Development, 58 Nepean Highway, Seaford, Victoria*, Gem Archaeology, Melbourne.

Cannon, M (ed.) 1983, *Historical Records of Victoria: Aborigines and Protectors 1838-1839*, Victorian Government Printing, Melbourne.

- Clark, I 1990, *Aboriginal Languages and Clans: An Historical Atlas of Western and Central Victoria, 1800-1900*, Department of Geography and Environmental Science Monash University, Melbourne.
- Coutts, PJF 1977, *Aboriginal Prehistory in North Western Victoria*, Victoria Archaeological Survey, Melbourne, Vic.
- Cummins, D, Ward, B & Power, R 2014, *Proposed 24 Unit Development, 200-202 Nepean Highway, Seaford, Victoria: Aboriginal Cultural Heritage Management Plan*, Ecology Partners Pty Ltd, Ascot Vale.
- D'Arcy, J 2005, *Bunurong Land Council Historical and Genealogical Report*, Bunurong Land Council, Aboriginal Corporation.
- Dudley, N, Marcus, D, Mann, K & Stradwick, J 2010, *Residential Development, 1-3 Attunga Crescent, Seaford Cultural Heritage Management Plan Number: 10940*, Kyandel Archaeological Services, Kew.
- Dugay-Grist, L 2009, *Seaford Village, Seaford: Cultural Heritage Management Plan No. 10637: Proposed Redevelopment of Seaford Village Precinct.*, Grist Archaeology Heritage Management, Unknown.
- Dugay-Grist, L, Cowled, A & Maher, M 2013, *178 & 178A Seaford Road, Seaford: 6 Unit Development*, Grist Archaeology Heritage Management, Frankston South.
- Dugay-Grist, L, Cowled, A & McAlister, R 2015, *Proposed Townhouse Development: 48-49 Nepean Highway, Seaford*, Grist Archaeology Heritage Management.
- Dugay-Grist, L, Cowled, A & Paynter, N 2013, *70 Armstrongs Road, Seaford: Residential Development*, Grist Archaeology Heritage Management.
- Dugay-Grist, L, Cowled, A & Paynter, N 2013, *22 Valetta Street, Carrum: Residential Development*, Grist Archaeology Heritage Management, Frankston South.
- Dugay-Grist, L & Dudley, N 2010, *95-96 Brunel Road, Seaford: Industrial Subdivision Cultural Heritage Management Plan 11382*, Urban Colours Arts Cultural Heritage Consultants, Longwarry.
- Dugay-Grist, L & McAlister, A 2011, *48 Nepean Highway, Seaford: Residential Subdivision*, Grist Archaeology Heritage Management, Frankston South.
- Gaughwin, D 1981, *Sites of Archaeological Significance in the Western Port Catchment Volume 1*, La Trobe University, Bundoora.
- Gaughwin, D & Sullivan, H 1984, 'Aboriginal Boundaries and Movements in Western Port, Victoria', *Aboriginal History*, vol. 8, no. 1, pp. 80-98.
- GHD 2005, *Edithvale Seaford Wetlands Review of Wetlands Operations*, GHD, Victoria.
- Griffin, D & McAlister, R 2010, *Lot 2, 28 Oliphant Way, Seaford Factory, Warehouse and Office Development Cultural Heritage Management Plan 11343*, Urban Colours Arts Cultural Heritage Consultants, Longwarry.
- Howell-Meurs, J 2008, *8-10 Klauer Street (Industrial Subdivision), Seaford Aboriginal Cultural Heritage Management Plan 10353*, Andrew Long & Associates and Archaeological Consulting Services, Fitzroy.
- Howell-Meurs, J 2011, *Lathams Road, Carrum Downs Industrial Development*, Andrew Long & Associates, Fitzroy.
- KBR 2009, *Edithvale-Seaford Ramsar Site: Management Plan*, Kellogg Brown & Root Pty Ltd, St Kilda.

- Lever, M & Houghton, K 2015, *3CSE-3KEY- Communications installation between Pacific Drive and Perry Road to Station Street and Mascot Avenue, Keysborough, Victoria*, Biosis Research Pty Ltd, Keysborough.
- Long, A, Mathews, D, O'Reilly, S, Feldman, R & Howell-Meurs, J 2009, *Frankston Bypass EES Mornington Peninsula, Victoria Cultural Heritage Management Plan Volume 1 – Cultural Heritage Management Plan 10015*, Andrew Long and Associates, Fitzroy.
- Marshall, B 1998, *Frankston City Council Aboriginal Cultural Heritage Assessment: A Report Commissioned by the Frankston City Council*, Austral Heritage Consultants, Coburg.
- Matic, A 2011, *78-83 Nepean Highway, Seaford: Proposed Residential Development*, Terra Culture Heritage Consultants Pty Ltd, Northcote.
- Matic, A 2013, *3 Canberra Street, Patterson Lakes: Residential Subdivision*, Pragmatic Cultural Heritage Services, Belgrave.
- Matic, A 2014, *42A Nepean Highway, Seaford: Residential Subdivision*, Pragmatic Cultural Heritage Services, Belgrave.
- McAlister, R 2014, *Proposed Residential Development 34 Melaleuca Drive, Carrum Cultural Heritage Management Plan*, Heritage Insight Pty Ltd, Collingwood.
- McCrae, H (ed.) 1934, *Georgiana's Journal: Melbourne A Hundred Years Ago*, Angus & Robertson, Sydney.
- McCrae, H (ed.) 1992, *Georgiana's Journal Melbourne 1841-1865*, Angus & Robertson, Sydney.
- Mitchell, J, Burch, J & McFarlane, E 2012, *Commercial Subdivision, 28-40 Colemans Road, Carrum Downs*, Alpha Archaeology Pty Ltd, Kensington.
- Mitchell, J & Heddell-Stevens, P 2015, *Proposed Warehouse Construction at 27 Brett Drive, Carrum Downs*, Alpha Archaeology Pty Ltd, Kensington.
- Mitchell, J & Richmond, J 2008, *Commercial Subdivision at 107 Boundary Road, Carrum Downs, Victoria Cultural Heritage Management Plan 10591*, Alpha Archaeology Pty Ltd, Kensington.
- Murphy, A & Dugay-Grist, L 2008, *Edithvale-Seaford Wetland Discovery Centre. Cultural Heritage Management Plan 10081*, Tardis Enterprises Pty Ltd, Beaconsfield Upper.
- Murphy, A & Morris, A 2013, *23 Chevron Court: Seaford, Residential Subdivision*, Archaeology at Tardis Pty Ltd, Beaconsfield.
- Murphy, A & Thomson, S 2012, *Retail Premises, 176-178, 179 Nepean Highway & 1 Seaford Road, Seaford*, Archaeology at Tardis Pty Ltd, Beaconsfield.
- O'Connor, A 2013, *216 Nepean Highway, Seaford Residential Development Cultural Heritage Management Plan*, Andrew Long & Associates Pty Ltd, Fitzroy.
- O'Reilly, S & Dudley, N 2011, *37 Fellowes Street, Seaford: Residential Development*, Urban Colours Arts Cultural Heritage Consultants, Lara.
- Plomley, NJB 1987, *Weep In Silence*, Blubber Head Press, Hobart.
- Presland, G (ed.) 1977, *Journals of George Augustus Robinson January - March 1840* Second Edition., Ministry for Conservation Publication, Melbourne.

- Rhodes, D 1990, *Dandenong Creek and Patterson River Melbourne: An Archaeological Survey, Occasional Report 33*, Department of Conservation and Environment, Melbourne.
- Rhodes, D 2003, *Victorian Channel Deepening Project EES: Aboriginal Cultural Heritage. Existing Conditions Report*, Terra Culture Heritage Consultants Pty Ltd, Fairfield.
- Rhodes, D & Bell, J 2004, *Shire of Cardinia Urban Growth Corridor Aboriginal Heritage Study*, Heritage Insight Pty Ltd, Richmond.
- Rogers, H 1957, *The Early History of the Mornington Peninsula*, Hallcraft Publishing Co. Pty. Ltd, Melbourne.
- Schell, P & Barker, A 2009, *300 Frankston - Dandenong Road, Seaford Cultural Heritage Management Plan 10583*, Ochre Imprints, Abbotsford.
- Shillinglaw, J (ed.) 1972, *Historical records of Port Phillip, the first annals of the colony of Victoria* Facsimile edition of the 1879 edition., Heinemann Publishers Australia.
- Smith, LJ 1991, *Berwick - Pakenham Corridor Aboriginal Archaeology*, Victorian Archaeological Survey, Melbourne.
- Stevens, J 2014, *Cultural Heritage Management Plan. 88 McLeod Road, Carrum*, Urban Colours Arts Cultural Heritage Consultants, Carrum.
- Stevens, J 2015, *Cultural Heritage Management Plan 644 Nepean Highway, Carrum Subdivision of land*, Urban Colours Cultural Resource Managers.
- Sullivan, H 1981, *Victorian Occasional Reports Series Number 6: An Archaeological Survey of the Mornington Peninsula, Victoria*, Victorian Archaeological Survey, Melbourne.
- Tucker, C & Marshall, B 2009, *Construction of new Dwellings at 211-212 Nepean Highway, Seaford Cultural Heritage Management Plan 11036*, Terra Culture Heritage Consultants Pty Ltd, Northcote.
- Tulloch, J 2004, *An Archaeological sub-surface testing program east of Boggy Creek, Carrum Downs: Final Report for Grogan Richards Pty Ltd*, Biosis Research Pty Ltd., Port Melbourne.
- Tunn, J 2013, *43 Kananook Avenue, Seaford Three lot subdivision Cultural Heritage Management Plan*, Ironbark Heritage and Environment.
- Tunn, J 2014, *39 Mitchell St, Seaford Residential Subdivision*, AHMS Archaeological & Heritage Management Solutions, Melbourne.
- Tunn, J & Rae, E 2015, *5 Kipling St, Carrum Residential Subdivision, Complex Cultural Heritage Management Plan*, AHMS Archaeological & Heritage Management Solutions, Melbourne.
- Vines, G, Fiddian, J, Thomson, M & Lawler, M 2007, *Boggy Creek Wetlands, Carrum Downs, Victoria: Cultural Heritage Management Plan 10014*, Biosis Research Pty Ltd, Port Melbourne.
- Walther, E 2015, *Cultural Heritage Management Plan 95 East Road Seaford Townhouse development*, ACHM, Kilmore.
- Whincop, M 2011, *75 Frankston Gardens Drive, Carrum Downs City of Frankston Industrial Subdivision Cultural Heritage Management Plan*, Andrew Long & Associates Pty Ltd, Fitzroy.
- Young, J & Prossor, L 2010, *Proposed Retirement Village and Carpark, 45 Fellowes St, Seaford Cultural Heritage Management Plan 11423*, Heritage Insight Pty Ltd, Richmond.

Appendix 1: Division 5 of the *Aboriginal Heritage Regulations 2007*

Division 5—High impact activities

42 Purpose

The purpose of this Division is to specify high impact activities.

Note

Under regulation 6, a cultural heritage management plan is required for an activity if all or part of the activity area is an area of cultural heritage sensitivity and if all or part of the activity is a high impact activity.

43 Buildings and works for specified uses

- (1) The construction of a building or the construction or carrying out of works on land is a high impact activity if the construction of the building or the construction or carrying out of the works—
 - (a) would result in significant ground disturbance; and
 - (b) is for or associated with the use of the land for any one or more of the following purposes—
 - (i) aquaculture;
 - (ii) a camping and caravan park;
 - (iii) a car park;
 - (iv) a cemetery;
 - (v) a child care centre;
 - (vi) a corrective institution;
 - (vii) a crematorium;
 - (viii) an education centre;
 - (ix) an emergency services facility;
 - (x) a freeway service centre;
 - (xi) a hospital;
 - (xii) an industry;
 - (xiii) intensive animal husbandry;
 - (xiv) a major sports and recreation facility;
 - (xv) a minor sports and recreation facility;
 - (xvi) a motor racing track;
 - (xvii) an office;
 - (xviii) a place of assembly;
 - (xix) a pleasure boat facility;

- (xix) a research centre;
 - (xx) a retail premises;
 - (xxa) a retirement village;
 - (xxi) a service station;
 - (xxii) a transport terminal;
 - (xxiii) a utility installation, other than a telecommunications facility, if—
 - (A) the works are a linear project that is the construction of an overhead power line with a length exceeding one kilometre or for which more than 10 power poles are erected; or
 - (B) the works are a linear project that is the construction of a pipeline with a length exceeding 500 metres; or
 - (C) the works are a linear project with a length exceeding 100 metres (other than the construction of an overhead power line or a pipeline with a pipe diameter not exceeding 150 millimetres); or
 - (D) the works affect an area exceeding 25 square metres.;
 - (xxiv) a veterinary centre;
 - (xxv) a warehouse;
 - (xxvi) land used to generate electricity, including a wind energy facility.
- (2) The terms used in subregulation (1)(b) have the same meanings as they have in the VPP.
- (3) Despite subregulation (1), the construction of a building or the construction or carrying out of works on land is not a high impact activity if it is for or associated with a purpose for which the land was being lawfully used immediately before the commencement day.
- (4) In this regulation, ***linear project*** has the same meaning as in regulation 68.

44 Constructing specified items of infrastructure

- (1) The construction of any one or more of the following is a high impact activity if the construction would result in significant ground disturbance—
- (a) an airfield;
 - (b) a bicycle track with a length exceeding 100 metres;
 - (c) a helipad;
 - (d) rail infrastructure, other than—
 - (i) a railway track with a length of less than 100 metres; or
 - (ii) a railway track siding with a length of less than 100 metres; or
 - (iii) a cutting with a length of less than 100 metres; or
 - (iv) a tunnel with a length of less than 100 metres; or
 - (v) a bridge with a span of less than 100 metres; or
 - (vi) a platform with a length of less than 100 metres; or

(vii) a service road with a length of less than 100 metres;"

(e) a road with a length exceeding 100 metres;

(f) a walking track with a length exceeding 100 metres;

(g) a telecommunications line consisting of an underground cable or duct with a length exceeding 500 metres.

(2) In this regulation, *telecommunications line* has the same meaning as in the VPP.

45 Dwellings

(1) The construction of three or more dwellings on a lot or allotment is a high impact activity.

(2) The carrying out of works for three or more dwellings on a lot or allotment is a high impact activity.

Example

Constructing an apartment tower containing 50 dwellings is a high impact activity.

Constructing or extending only one or two dwellings on a lot or allotment is not a high impact activity.

Note

See regulation 9 in relation to the construction of a building, or the construction or carrying out of works, where the building or works are ancillary to an existing dwelling or the construction of one or two dwellings on a lot or allotment.

(3) This regulation does not apply to the construction of or the carrying out of works for a retirement village within the meaning of the VPP.

46 Subdivision of land

(1) The subdivision of land into three or more lots is a high impact activity if—

(a) the planning scheme that applies to the activity area in which the land to be subdivided is located provides that at least three of the lots may be used for a dwelling or may be used for a dwelling subject to the grant of a permit; and

(b) the area of each of at least three of the lots is less than eight hectares.

(2) The subdivision of land into two or more lots in an industrial zone is a high impact activity.

(3) In this regulation, *industrial zone* has the same meaning as in the VPP.

47 Alpine resorts

(1) The construction of a building or the construction or carrying out of works in an alpine resort is a high impact activity if the construction of the building or the construction or carrying out of the works would result in significant ground disturbance.

(2) In this regulation, *alpine resort* has the same meaning as in the **Alpine Resorts Act 1983**.

48 Activities requiring earth resource authorisations

An activity is a high impact activity if it is an activity—

- (a) for which an earth resource authorisation is required before the activity may be carried out; and
- (b) that would result in significant ground disturbance.

48A Extraction or removal of stone

- (1) The extraction or removal of stone (other than sand or sandstone) that does not require an earth resource authorisation is a high impact activity if—
 - (a) the primary purpose of the extraction or removal is—
 - (i) the sale or commercial use of the stone; or
 - (ii) the use of the stone in construction, building, road or manufacturing works; and
 - (b) the land from which the stone is extracted or removed is more than 2000 square metres; and
 - (c) the extraction or removal would result in significant ground disturbance.
- (2) In this regulation, **stone** has the same meaning as in the **Mineral Resources (Sustainable Development) Act 1990**.

49 Extraction or removal of sand or sandstone

- (1) The extraction or removal of sand or sandstone (other than extraction or removal that requires an earth resource authorisation) is a high impact activity if the extraction or removal would result in significant ground disturbance.
- (2) Subregulation (1) does not apply to the extraction or removal of sand or sandstone—
 - (a) from land that is a farm if the sand or sandstone is intended in good faith only to be used on that farm for the purposes of a dam or other farmworks and not for sale or any other commercial use; or
 - (b) undertaken by or on behalf of a Minister responsible for the administration of the **Conservation, Forests and Lands Act 1987** where the primary purpose of the extraction is for the footings or foundations of a building or structure, the construction of a carpark, road, track or other works or for any borrow pit adjacent to such an excavation; or
 - (c) if the extraction or removal, including dredging, constitutes works for marine navigational purposes or the establishment or renourishment of a beach; or
 - (d) if the extraction or removal constitutes works for the purpose of establishing a port facility, railway or tunnel; or
 - (e) if the primary purpose of the excavation or removal is for the construction of the footings or foundations of a building or structure.

50 Searching for stone

- (1) A search for stone is a high impact activity if it would result in significant ground disturbance.
- (2) In this regulation, **search for stone** has the same meaning as in the VPP.

51 Extraction or removal of loose stone on agricultural land on the Victorian Volcanic Plain

- (1) The extraction or removal of loose stone from the surface of land used for agriculture on the Victorian Volcanic Plain is a high impact activity if the extraction or removal—
 - (a) is for the primary purpose of land improvement, including pasture enhancement; and
 - (b) would result in significant ground disturbance.
- (1A) The crushing of loose stone on the surface of land used for agriculture on the Victorian Volcanic Plain is a high impact activity if the crushing is—
 - (a) by machinery; and
 - (b) for the primary purpose of land improvement, including pasture enhancement."
- (2) In regulation 51(2) of the Aboriginal Heritage Regulations 2007, for "Subregulation (1) does" **substitute** "Subregulations (1) and (1A) do".
- (3) In regulation 51(3) of the Aboriginal Heritage Regulations 2007, for "**Extractive Industries Development Act 1995;**" **substitute** "**Mineral Resources (Sustainable Development) Act 1990;**
- (2) Subregulation (1) does not apply if the land is used for crop raising or has been used for crop raising.
- (3) In this regulation—

agriculture and **crop raising** have the same meanings respectively as they have in the VPP;

stone has the same meaning as in the **Extractive Industries Development Act 1995;**

Victorian Volcanic Plain means the area comprised of the areas identified as "Qvh", "Qvn", "Qvs", "Qvs2" and "Qvt" on the following Geological Survey of Victoria 1:250 000 map series sheets—

 - (a) SJ54-8 entitled "Ballarat" (second edition, 1997);
 - (b) SJ55-1 entitled "Bendigo" (third edition, 2001);
 - (c) SJ54-12 entitled "Colac" (second edition, 1997);
 - (d) SJ54-7 entitled "Hamilton" (second edition, 1997);
 - (e) SJ55-5 entitled "Melbourne" (second edition, 1997);
 - (f) SJ54-11 entitled "Portland" (second edition, 1997);
 - (g) SJ55-9 entitled "Queenscliff" (second edition, 1997);
 - (h) SJ54-4 entitled "St Arnaud" (second edition, 1997).

52 Timber production

- (1) The use of an area of land greater than 40 hectares in size for timber production is a high impact activity if—
 - (a) a permit is required under a planning scheme to use the land for timber production; and
 - (b) the use of the land for timber production would result in significant ground disturbance.
- (2) The construction of a building associated with timber production is a high impact activity if—
 - (a) a permit is required under a planning scheme to construct the building; and
 - (b) the construction of the building would result in significant ground disturbance.
- (3) In this regulation, *timber production* has the same meaning as in the VPP.

r. 52

Note

A permit may not be required under a planning scheme to use an activity area for timber production if the timber production is ancillary to a particular agricultural enterprise (such as agroforestry).

53 Dams

The construction or alteration of a private dam, other than on a waterway, is a high impact activity if a licence is required under section 67(1A) of the **Water Act 1989** for the construction or alteration of the private dam.

54 Use of land

- (1) The use of land for a purpose specified in regulation 43(1) is a high impact activity if a statutory authorisation is required to use the land for that purpose.
- (2) The use of land for an extractive industry is a high impact activity if a statutory authorisation is required to use the land for the extractive industry.
- (3) The use of a lot or allotment for three or more dwellings is a high impact activity if a statutory authorisation is required to use the lot or allotment for three or more dwellings.
- (4) Despite subregulations (1), (2) and (3), if the whole of the activity area for an activity referred to in subregulation (1), (2) or (3) has been subject to significant ground disturbance, that activity is not a high impact activity.
- (5) In this regulation, *extractive industry* has the same meaning as in the VPP.

Example

A land owner proposes to change the use of his or her land from the grazing of animals to the storage of shipping containers. The land is flat and, in the first instance, no works are

proposed, although the grass will first be cut and some non-indigenous shrubs removed. The use of the land for storing shipping containers is an industry and requires a statutory authorisation (a permit under the relevant planning scheme). The proposed use is a high impact activity. If, at a later date, the area is upgraded by works, including excavation for a concrete base on which to store the containers, the upgrade works would also be a high impact activity under regulation 43(1).

Appendix 2: Glossary

Adze A flake with stepped retouch along lateral margins that can be hafted for use as a tool.

Anvil A flat object on which a core was placed to flake material from. Anvils often have a small pit/groove, usually in the centre of the object, as a result of this action.

Archaeology The study of cultural remains from past cultures and generations.

Artefact Scatter The material remains of past Aboriginal peoples' activities. Usually contain stone artefacts, but other material may also be present, including charcoal, animal bone, shell and ochre. An artefact scatter is usually represented by a single stone flake or a concentration of flaked stone pieces (or fragments).

Assemblage A collection of artefacts that are derived from the same site.

Backed Blade Stone artefact associated with the Australian small tool tradition. They are characterised by unidirectional or bidirectional retouch found along a lateral margin, thought to be blunt for hafting (Holdaway & Stern 2004: 260).

Basalt A fine-grained rock occurring from lava flows.

Bifacially Flaked Flakes removed from two faces of an object such as a core.

Blade A flake that is twice as long as it is wide.

Bondi Point An asymmetrical blade with a point at one end with backing retouch. Part of the Australian Small Tool Tradition.

Burial Human Remains, normally found as concentrations of human bones or teeth, exposed by erosion or earthworks. They are sometimes associated with charcoal or ochre, although shell, animal bone and stone tools may also be present. Tend to be located in soft soils and sand, although can occur in rock shelters, caves and dead trees.

Burin A truncated flake formed by snapping or retouching along one lateral margin that then forms a platform from which small flakes are removed forming a triangular scar that acts as a working edge (Holdaway & Stern 2004: 241-243).

Ceramic A term used to identify wares made from either clay or fusible stone such as stoneware, earthenware, porcelain or terracotta (Davies & Buckley 1987: 186).

Chert A compact, fine-grained rock made of cryptocrystalline silica and can occur in a variety of colours, usually red, green or black.

Core A specimen of rock that has undergone a process of reduction through the removal of a number of flakes and as a result they have negative flake scars. Cores can contain a single platform, have two platforms or have had flakes removed in multiple directions.

Cortex The original surface of a mineral or rock subjected to weathering by the elements.

Cultural Material Any material remains which are produced by human activity.

Debitage Detached pieces of stone that are discarded during the reduction process.

Dry Stone Wall A wall formed of a number of courses of rock (usually basalt or limestone) with no bond or binding component. Walls are usually tapered, have two faces and can have hearting (packing), or plugging.

Earthenware A non-vitreous (porous) whiteware, usually used for domestic tablewares. Most earthenware is glazed and decorated, transfer printed or left plain (Davies & Buckley 1987: 186).

Earth Feature Collective term used to refer to mounds, rings, hearths, postholes and ovens.

Earth Mound Mounds generally appear as raised areas of darker soil. They are commonly found in the volcanic plains of western Victoria or on higher ground near water bodies. Mounds often contain charcoal, burnt clay or stone heat retainers from cooking ovens, animal bones, shells, stone tools and sometimes, Aboriginal burials.

Earth Ring Banked circles of soil often associated with stone arrangements, which had a ceremonial purpose for Aboriginal people in the past.

Excavation A controlled means of soil disturbance (digging) allowing for detailed recording of the soil profile, features and artefacts exposed.

Flake A stone artefact that contains characteristics such as the presence of a platform, bulb of percussion and termination which reveal that the stone has been struck from a core and is the result of stone working (Holdaway & Stern 2004: 5).

Flake Core A flake that has subsequently been used as a core and had other flakes removed from it.

Flaked Piece Small fragments of stone that have been removed from flakes resulting from tool maintenance or tool production (Holdaway & Stern 2004: 17). Flaked pieces do not display the characteristics evident in a complete flake.

Flint Similar to chert with a pale cortex and conchoidal fracture. Usually occurring in limestone (Roberts 1998: 65).

Footing The structural base/footprint from structures often built from bluestone, brick or wooden posts.

Geometric Microlith Part of the Australian small tool tradition. They are symmetrical in form, pointed at both ends and can be backed along a lateral margin (Holdaway & Stern 2004: 262).

Glaze A coating put over wares fired in a kiln. Glazes can come in a variety of colours and can also be transparent.

Greenstone A metamorphic rock derived from basalt containing feldspar and quartz and is made green by chlorite and epidote. Often used for the manufacture of hand axes.

Grindstone A flat slab of rock with central depression used to grind, crush or pound seeds, ochre, or sharpen tools, etc. Grindstones are usually made on sedimentary rocks with an abrasive surface and can be used in conjunction with a muller.

Ground Edge Axes A sharpening process – flaking, pecking and polishing, usually along a single lateral margin. The axes are generally hafted with the worked edge forming the tool edge.

Ground Surface Visibility The extent to which the natural soil surface below the vegetation on the ground is visible.

Hammerstone A hard rock or mineral used to flake fragments of stone from a core (Holdaway & Stern 2004: 4).

Hearth The remains of a fireplace containing charcoal and sometimes burnt earth, bone, stone artefacts or other organic material.

In situ An artefact or feature that remains in its original position, or where it was left.

Manuport A stone block that displays no attributes of being either a core or a flake.

Microblade Has the same characteristics as a blade but just of smaller proportions (Holdaway & Stern 2004: 17).

Ochre Earth varying in colour from yellow to red, used as a pigment.

Organic Compounds formed from living organisms (plants or animals).

Oven Mound Usually circular or oval in shape and often situated close to a water source. They were used for cooking and contain a rich greasy organic mix of soil and organic material. An oven mound is likely to contain charcoal, burnt clay or stone heat retainers, stone tools, bones, shell and on occasion, burials (AAV Mini Poster 4).

Platform The surface from which the flake was struck off the core – natural, flaked or abraded (Holdaway & Stern 2004: 120).

Point A flake that has two edges that form a point with retouch along one or both lateral margins (Holdaway & Stern 2004: 16).

Porcelain A non-porous ceramic with a glass-like appearance. Can be translucent, can be used for tableware or more decorative features such as ornaments.

Post-contact The period after contact between Aboriginal people and Europeans.

Pre-contact The period before contact between Aboriginal people and Europeans.

Quarry Outcrop of stone or ochre that has been quarried by Aboriginal people in the past. Generally

associated with a large amount of broken stone and flakes. The outcrop (cores) bear negative scars from flaking.

Quartz A mineral that commonly occurs in sedimentary, igneous and metamorphic rocks. Quartz can come in a number of forms including crystal, rose, and smoky.

Quartzite A metamorphic rock formed by the recrystallization of quartz. Quartz is rich in sandstone and limestone (Roberts 1998: 109).

Retouch A worked edge or modification of a flake formed by removing a number of small flakes along an edge. This can be done as a form of maintenance or to produce a tool.

Rock Art Paintings created on the rock surfaces of caves and rock shelters and engravings in limestone caves. Artwork includes stencils, prints and drawings. The paint consists of ochres, clays and charcoal mixed with fats.

Scarred Tree A tree which has had a slab of bark removed, exposing the sapwood on the trunk or branch of a tree. Aboriginal people used the bark to make shelters, containers (coolamons) and canoes.

Scraper A flake with at least one edge that has continuous retouch. Scraper types include steep-edged, end, side and nose scraper (Holdaway & Stern 2004: 16).

Shell Midden A surface and/or sub-surface deposit composed of shell and sometimes stone artefacts, charcoal and bone. Middens are normally found in association with coastlines, rivers, creeks and swamps – wherever coastal, riverine or estuarine shellfish resources were available and exploited.

Silcrete A fine-grained rock derived from shale or siltstone mixed with silica.

Spit A horizontal unit of soil removed during excavation. Spits can be arbitrary (dug to a depth of 50, 100, 200, 300mm, etc.) or can be confined to a particular soil type or context. The excavation of spits allows for greater understanding, analysis and interpretation of the soil profile.

Stone Feature Includes cairns, rock wells, stone arrangements, fish traps, stone structures and grinding

grooves. May be a natural feature, which was used or modified to be used by Aboriginal people in the past (rock well, stone arrangement), or a stone feature which has been deliberately constructed for a specific purpose (fish trap, stone structure, cairn), or is the result of a specific activity carried out by Aboriginal people in the past (grinding grooves).

Stoneware A vitreous (non-porous) ceramic, usually light brown in colour, used for drinking containers or used industrially. Often glazed or unglazed (salt glaze or slip applied) (Davies & Buckley 1987: 186).

Stratification The position of sediments and rocks in sequence throughout time.

Sub-surface Testing A method of excavation that involves ground disturbing works to identify the potential for cultural material. Sub-surface testing may comprise hand excavation and/or machine excavation.

Survey An inspection of land either by foot or by car (windscreen survey) noting conditions on surface visibility, landforms and the presence of cultural material.

Termination The shape of the distal end of a flake (Holdaway & Stern 2004: 129).

Terracotta A low-fired clay (ceramic), usually orange to red in colour and very porous. Often used for plumbing (drainage components) or garden ware.

Tool Modified flakes usually with retouch present along an edge (Holdaway & Stern 2004: 33).

Transect An excavated stretch of ground that can be of varying lengths in a straight line.

Transfer Printed A design is traced and engraved onto a copper plate on which ink and oil is then applied. The design is pressed onto tissue paper and then placed on an object and the paper removed. The object is then fired and glazed. Transfer printed ceramics come in a variety of colours and patterns and were mass produced.

Trench An area confined by excavation usually in the form of a square (e.g., 2x2m) or rectangular (e.g., 1.5x1m).

References

AAV Mini Posters (1-7)

Davies, M. & K. Buckley, 1987 *Port Arthur Conservation & Development Project: Archaeological Procedures Manual*. Occasional Paper No.13. Department of Lands, Parks and Wildlife, Tasmania.

Holdaway S. & N. Stern, 2004 *A Record in Stone: The Study of Australia's Flaked Stone Artefacts*, Museum Victoria, Melbourne.

Roberts, J. L., 1998 *A Photographic Guide to Minerals, Rocks and Fossils*, New Holland, London.

Appendix 3: OAAV Practice Note for Determining Significant Ground Disturbance